

Sampling Feed For Analysis

J.W. Schroeder
Livestock Specialist-Dairy


Kevin Sedivec
Rangeland Management Specialist

Warning

A chemical analysis can be no better than the sample submitted. Make sure the sample is representative of a selected lot of feed. Remember: Know what analysis is needed and inform the laboratory. See the county Extension office for a current listing of available labs.

Proper handling of the sample between the farm and laboratory ensures best results. General rules are:

- Sample size: minimum of one quart (be sure the sample is representative). All forages should be chopped to a length of 3 inches or less to make handling easier.
- Pack tightly to exclude air. Seal air tight. Use plastic bags in all cases except very dry samples.
- Send samples to the laboratory as quickly as possible. Indicate analysis wanted by letter or with appropriate form, if part of an organized program.
- All letters, instructions, checks or money orders should be put in first class mail envelope and attached to mailing container. The sample container can go as fourth class mail.
- Address both letter and sample container with correct laboratory address and return address



Equipment For Sampling

- Ratchet brace and square shank adaptor
- Electric drill and round shank adaptor
- Penn State forage sampler with dowel plunger to remove moist or packed samples
- Sack thief
- Grain probe
- Plastic bags for submitting samples

Baled Hay

With a commercial forage sampler (for example the Penn State sampler), core sub-samples from 15-20 bales should be used for the composite lab sample. The bales should be selected at random for each lot of hay. If the hay will be fed by its cutting (first cutting alfalfa, etc.) each cutting should be analyzed. If a mixture of cuttings or classes of hay will be fed, a proportional number of bales can be used from each class to approximate the feed offered to the animals. For square bales, sample from the end, the full length of the sampler tube. Place each core in the container. For round bales, sample across the bale at the center. You can sample baled hay without a forage sampler by removing a small section from each bale (15-20 bales) and cutting the hay into 3 inch length with shears or hatchet. This is a less desirable technique, since you're almost sure to lose leaves. Make every effort to include the appropriate combination of leaves and stem. Mix and then randomly grab handfuls of the chopped sample. Place them in the container to be sent to the laboratory.

Loose Hay

Using the commercial forage sampler or equivalent, place the hay sample beneath your feet to compress the hay and sample between your feet the full depth of the sampler. Random samples from the side of the mow or stack should be taken where the hay is compressed enough for good sampling. An extension shaft on the sampler can be used to sample deeper into the hay. At least 15 core samples should be taken. You can sample loose hay without a forage sampler by removing a handful of hay from many (15-20) positions from each lot of hay. Be careful to save all leafy material belonging to the sample. Cut into 3 inch length, mix and send representative sample to the laboratory.

Pasture

Pasture sampling is most difficult. Fertility and moisture differences in a single pasture add to the problem. Sample by picking at random 8-10 locations. Remove the forage from a square foot area at grazing height. Mix all the collected forage and take a representative sample for analysis.

A second method is to take forage being selectively grazed by the animals at several locations for the sample. This is a preferred method in unimproved pasture where selective grazing is evident. However, to accurately determine how much of which forage to sample can be difficult. With a little practice, an experienced manager can accurately identify the species being consumed at the time of sampling.

Precaution

Green pasture samples should be immediately dried or frozen to prevent marked chemical changes. The most practical alternative is to pack the sample tightly in a plastic bag, exclude all possible air, freeze, and promptly mail to the laboratory. With good container insulation, the sample will arrive in a cool condition or with a minimum of a silage-like fermentation. When choosing to dry a sample, take care to

control the temperature and time the sample is exposed to heat. Drying for too long or at too high a temperature will significantly alter the composition of your sample. The microwave can be an excellent tool for drying samples.

Forage Crops in the Field

Remove the plants from a square foot area at normal harvesting heights at several (8 to 10) locations in the field. Whenever possible, chop the forage into small pieces (1 to 2 inch), mix and remove a representative sample. This sample should be handled as noted in the precaution section under pasture. Where you want nitrate and cyanide analysis, a field check should be made or the sample quickly frozen (use dry ice when possible) and transported to the laboratory in a frozen state. The time from chopping to freezing should be less than 15 minutes for cyanide. However, the whole plant can be transported to the laboratory within two hours with little loss.

Green Chopped Forages

A plastic bag can be filled by randomly grabbing a handful of the chopped material from the cutter, blower or several locations in a wagon or truck. Since the material will be fed as such, or ensiled, the plastic bag should be packed tightly, sealed with air excluded and sent to the laboratory immediately. The fermentation will reduce nitrate content and slightly modify other components but this change can be considered minor and in the direction of its ultimate feeding value. When more exact values are required, the sample should be frozen before transporting.

Silage

All silage samples should represent several locations within the silo to ensure a representation of the silage. The representative sample must be tightly packed in a plastic bag, with all air excluded, and sealed. The sample should be sent to the laboratory as soon as possible.

Trench Silo

If silo is open, the face of the silage should be cleaned off in the center. A column of silage 6 inches by 12 inches should be removed from top to bottom, mixed thoroughly and then a representative sample taken.

Where the silo is not opened, a series of (four to six) holes can be dug from the top with a post hole digger or suitable equipment. Spoiled silage should be placed beside the hole to be returned after sampling. The samples of good silage from each hole are mixed and a representative sample taken. Be sure holes are packed tightly with the silage that has been removed to avoid undue spoilage.

Upright Silo

Representative sampling is also difficult. An open silo can be sampled by taking grab samples from a given day(s) feed. If unopened, a procedure similar to that suggested for the unopened trench can be used.

Silage and haylage moisture (percent dry matter content) can vary considerably from one level to another in both upright and horizontal silos. Due to many reasons such as date of chopping, rainfall, differences in fields, etc, this factor is important when allowing for both the nutrient and fiber content in the diet.

Because silages and haylage are more variable in their dry matter content than feeds stored in the dry form, it's important to continue monitoring their moisture level on a regular basis during the feeding period. The testing interval, however, will depend on how fast the silo is emptied. For instance, if the silo is fed out quickly, samples taken whenever 20-25 percent of the contents are removed should be sufficient to monitor variations in moisture and nutrient content. Even if the silage remains in the silo for extended periods, its composition changes with the fermentation process. In this case, samples taken every four to six weeks of the feeding period are advisable. Regardless, a silo should be sampled more than once during the feeding period, especially when more than one field or kind of forage is stored in the silo. Both the microwave method and the commercial moisture testers can provide quick and reliable results to help you make the appropriate adjustments for a balanced ration.

Sacked Feed

Most sacked feed is thoroughly mixed. However, it is recommended that at least five to six sacks be sampled (1 to 2 pounds), mixed and a representative sample (1 pound) submitted for analysis. Either a small probe or sack thief should be used when available. Be aware that settling is quite common, even in sacked feed, making thorough sampling important.

Bulk Concentrates

Commodity feeds should be analyzed as a composite of at least 10-15 areas of a given lot of feed. When mixing the composite, avoid segregation by particle size or the true sample value may be distorted. At least 1 pound, or a quart of material, should be sent to the laboratory. Be advised that these results represent bulk averages, and will not give you information on the uniformity of nutrient content within the mix. If you're experiencing inconsistent herd performance using these feeds, the uniformity of the mix may be questionable and a different technique should be employed. Taking several samples of the commodity and having each analyzed will allow the producer or feed nutritionist to do a better job of adjusting the ration to accommodate the feed differences.

Grain in Bin

It is highly desirable to use a grain probe to obtain the sample. However if one is not available random grab-samples from 10-15 areas of the bin can be mixed and a representative sample (1 quart) sent in for analysis.

General

Getting a representative sample is as important as procedural and technical expertise. When sampling, consider the intended use of the results and the time and cost involved. When forage analysis reveals a quality too low to meet your herd or flock needs, you may want to reexamine the cropping practices (fertilizer used, variety selection of forage, harvest time and method of harvest, etc.) and provide supplements to meet livestock production demands. Those who find their feed supply to be above average can save money by reducing their supplements. Even when an analysis reveals feed supplies that are of average quality, this information serves as a basis for sound crop and livestock decisions that will help maximize economic farm returns.

When forage quality is low, testing is the most important investment you can make. This is a good time to consider a 'wet chemistry' lab analysis. When producers find no choice but to utilize inferior quality feeds, experience shows that it's prudent to discount the expected nutrient value. Even though you're trying to save on feed inputs, poor quality feeds generally don't elicit the animal performance expected. NIRS (near infrared spectrophotometry), although quick and inexpensive, should not be relied upon in these cases unless duplicate samples of some of the feeds/forages have also been double checked with those results of a reliable lab using more extensive analysis.

Who Will Test Your Samples?

Forage and grain testing for nutrient content can be performed via NIRS or through wet chemistry. Wet chemistry can be used for any forage type or mixture of forages, grains and other plant species. While it is preferable to test individual diet components, wet chemistry is the method of choice when considering mixed feeds. NIRS is used only for "pure" samples due to set calibrations for specific plant species or grain type. Examples of "pure" samples are 100 percent barley or 100 percent alfalfa, without grasses or weedy species.

NIRS is a less expensive method of analysis that provides similar results to wet chemistry when the sample is pure. Some local elevators and feed companies will test forages by NIRS testing, with most elevators testing grain samples through NIRS. Wet chemistry is only conducted by laboratories capable of performing tests on feedstuffs. The National Forage Testing Association (1993) listed 98 laboratories capable of conducting both NIRS and wet chemistry analysis. Eleven known labs conduct both NIRS and wet chemistry in the North Dakota, Minnesota and South Dakota area. For a copy of these lists, please contact your local Extension agent.

Toxicology verses nutrient analysis –

The techniques described in this publication apply to samples taken for either nutrient content or toxicological tests. Refer to the lab request form to determine the analytical services offered.

Feed Sampling Checklist (✓)

This page is provided as a sampling reminder and feed analysis record. DO NOT use it to substitute for the appropriate lab analysis request forms supplied by the testing laboratory. Use this reference to track your samples and as a worksheet for transferring the collected information to the laboratory request sheet.

Laboratory Information —

Refer to the lab of choice for charges. (Costs will vary with lab, type of tests and combination of tests requested.)

Name of Lab _____ Address _____

City _____ State _____ Zip _____

Phone (_____) _____ FAX (_____) _____

Type of feed (example — 50% alfalfa:50% brome) _____

Analysis requested (check those requested)

_____ WET CHEMISTRY

- moisture (dry matter)
- fat
- pH
- crude protein
- soluble protein
- degradable protein
- crude fiber
- ADF (acid detergent fiber)
- NDF (neutral detergent fiber)
- ash

Minerals

- aluminum (Al)
- calcium (Ca)
- cobalt (Co)
- copper (Cu)
- iron (Fe)
- magnesium (Mg)
- manganese (Mn)
- molybdenum (Md)
- phosphorus (P)
- potassium (K)
- selenium (Se)
- sodium (Na)
- sulfur (S)
- zinc (Z)

_____ NIRS (Near Infrared Spectrophotometry)

- moisture
- protein (dry matter)
- available crude protein
- ADF (acid detergent fiber)
- NDF (neutral detergent fiber)
- calcium (Ca)
- magnesium (Mg)
- phosphorus (P)
- potassium (K)

Calculations requested (from either wet lab or NIRS above) NEL (net energy for lactation) RFV (relative feed value)

Others (not included above, example — vitamins, toxicology) _____

Background information — You are encouraged to record the following information for your reference. The lab may not require it, but often times additional information can be useful to evaluate your request.

Acres being sampled (example — 50 acres field peas and oats) _____

Livestock to be fed (example — yearling Holstein heifers, dry cows) _____

Description of feed _____

Date harvested _____ Date sampled _____ Date sent _____

Weather conditions at harvest _____

How will this feed be used? _____

(Make extra copies for future use)

For more information on this and other topics, see: www.ag.ndsu.edu

This publication may be copied for noncommercial, educational purposes in its entirety with no changes. Requests to use any portion of the document (including text, graphics or photos) should be sent to NDSU.permission@ndsu.edu. Include exactly what is requested for use and how it will be used.

North Dakota State University does not discriminate on the basis of race, color, national origin, religion, sex, gender identity, disability, age, status as a U.S. veteran, sexual orientation, marital status, or public assistance status. Direct inquiries to the Vice President for Equity, Diversity and Global Outreach, 205 Old Main, (701) 231-7708.

County Commissions, NDSU and U.S. Department of Agriculture Cooperating.

This publication will be made available in alternative formats for people with disabilities upon request, (701) 231-7881.

3M-9-93; 1.5M-3-97; W-7-10