

North Dakota durum wheat variety descriptions, agronomic traits, 2009.

| Variety | Agent or Origin ¹ | Year Released | Chaff Color | Height | Straw Strength | Maturity | Reaction to Disease ² | | | |
|---------------|------------------------------|---------------|-------------|--------|----------------|----------|----------------------------------|-----------|----------------|------|
| | | | | | | | Stem Rust | Leaf Rust | Foliar Disease | Scab |
| AC Avonlea | Can. | 1997 | White | Med. | Med. | Med. | R | R | MS | S |
| AC Commander | Can. | 2002 | White | S.dwf. | Med. | Med. | R | R | MS | NA |
| AC Melita | Can. | 1995 | White | Tall | Med. | Med. | R | NA | NA | S |
| AC Morse | Can. | 1996 | White | S.dwf. | Strong | Med. | R | R | M | NA |
| AC Napoleon | Can. | 2001 | White | S.dwf. | Med. | Med. | R | R | S | NA |
| AC Navigator | Can. | 1999 | White | S.dwf. | Weak | Med. | R | R | M | S |
| AC Pathfinder | Can. | 1999 | White | Med. | Weak | Med. | R | R | M | S |
| Alkabo | ND | 2005 | White | Med. | V.strong | Med. | R | R | M | MS |
| Alzada | WB | 2004 | White | S.dwf. | Strong | Early | R | R | S | VS |
| Belzer | ND | 1997 | White | Tall | Med. | Late | R | R | M | MR |
| Ben | ND | 1996 | White | Med. | Strong | Med. | R | R | MR | S* |
| Cando | ND | 1975 | Tan | S.dwf. | V.strong | Med. | R | R | M | VS |
| DG Max | DGP | 2008 | White | Med. | Strong | Med. | R | MR | MR | MS |
| DG Star | DGP | 2007 | White | Med. | Strong | Early | R | R | M | NA |
| Dilse | ND | 2002 | White | Med. | Strong | Late | R | R | M | MS |
| Divide | ND | 2005 | White | Med. | Strong | Med. | R | R | M | MR |
| Dressler | AgriPro | 1996 | White | Tall | Med. | Med. | R | MR | NA | VS |
| Fjord | AgriPro | 1986 | White | Tall | Strong | M.early | R | R | M | S |
| Grande D'Oro | WB/DGP | 2005 | White | Med. | Strong | Med. | R | R | M | NA |
| Grenora | ND | 2005 | White | Med. | Strong | Med. | R | R | M | MS |
| Kari | AgriPro | 1998 | White | Med. | Strong | Med. | R | R | M | S |
| Kyle | Can. | 1984 | White | Tall | Weak | Med. | R | MR | M | NA |
| Laker | WB | 1985 | White | S.dwf. | Strong | Med. | R | MR | S | S |
| Lebsock | ND | 1999 | White | Med. | Strong | Med. | R | R | M | MS |
| Lloyd | ND | 1983 | White | S.dwf. | V.strong | Med. | R | MR | S | VS |
| Maier | ND | 1998 | White | Med. | Strong | M.late | R | R | M | S* |
| Medora | Can. | 1983 | White | Tall | Strong | M.early | R | R | MS | VS |
| Monroe | ND | 1985 | White | Tall | Med. | Early | R | R | M | VS |
| Mountrail | ND | 1998 | White | Med. | Strong | Late | R | R | M | S* |
| Munich | ND | 1995 | White | Med. | V.strong | Med. | R | R | MR | S* |
| Pierce | ND | 2001 | White | Med. | M.strong | Med. | R | R | MS | S |
| Plaza | ND | 1999 | White | S.dwf. | V.strong | Late | R | R | M | MS |
| Plenty | Can. | 1990 | White | Tall | Weak | Late | R | R | MR | MS |
| Primo D'Oro | WB/DGP | 2004 | White | Tall | Med. | M.early | R | R | MS | NA |
| Renville | ND | 1988 | White | Tall | Med. | Med. | R | R | M | S* |
| Rugby | ND | 1973 | Tan | Tall | Strong | M.early | R | R | MR | S* |
| Strongfield | Can | 2004 | White | Med. | Med. | Med. | R | R | MS | NA |
| Vic | ND | 1979 | White | Tall | Med. | M.early | R | R | MR | S* |
| Voss | AgriPro | 1994 | White | S.dwf. | V.strong | Med. | R | MR | MS | S |
| Wales | Westbred | 2008 | White | Med. | M.strong | Med. | R | R | M | S* |

¹Refers to agent or developer: Can = Agriculture Canada, WB = Westbred, ND = North Dakota State University, DGP = Dakota Growers Pasta.

²R = resistant; MR = moderately resistant (slow rusters); M = intermediate; MS = moderately susceptible; S = susceptible; VS = very susceptible; Foliar Disease = reaction to tan spot and septoria leaf spot complex. Letter ratings for head blight (scab) based on visual head symptoms. * Indicates yields and/or quality often have been higher than would be expected based on visual symptoms. NA = Not adequately tested.