

# Management of Insecticide-resistant Soybean Aphids

Robert Koch, University of Minnesota  
Erin Hodgson, Iowa State University  
Janet Knodel, North Dakota State University  
Adam Varenhorst, South Dakota State University


Soybean aphids, *Aphis glycines*, (Figure 1) are the most significant insect pest of soybean in Minnesota, Iowa, North Dakota and South Dakota. Development of insecticide resistance in this pest creates new challenges for effective soybean pest management and profitable soybean production.

Resistance is defined as a decrease in susceptibility of a pest population to an insecticide that may result in failures when the product is used according to label recommendations for that pest.


Failures of certain pyrethroid insecticides for management of some soybean aphid populations have been observed in commercial fields (Figure 2), and resistance to bifenthrin and lambda-cyhalothrin has been documented through small-plot research and laboratory bioassays. Because of the mobility of winged soybean aphids, the challenges posed by insecticide-resistant populations of the pest could spread to soybean fields in other parts of the region.




**Figure 1.** Soybean aphids infesting a soybean leaf.  
(Photo courtesy of R. Koch, University of Minnesota)




**2015**



**2016**



**2017**



**2018**

**Figure 2.** Counties with reported failures of pyrethroids for control of soybean aphid. Red-shaded counties indicate those from which Extension entomologists received reports of failures. (Maps courtesy of B. Potter, University of Minnesota)

## Management Strategies for Resistant Soybean Aphids

In response to the challenge that insecticide-resistant soybean aphids pose, we encourage growers, consultants and applicators to evaluate their soybean aphid management practices carefully. Best management practices include:

- Treat fields only when needed to reduce insecticide exposure to soybean aphids. This will reduce the selection pressure for further development of resistance.
  - ◆ Fields should be scouted on a regular schedule (every seven to 10 days).
  - ◆ Use the economic threshold (250 aphids per plant, with greater than 80 percent of plants infested) to determine when to apply insecticides.
  - ◆ Treat within five to seven days of exceeding the economic threshold to protect yield.
- If a field exceeds the threshold, make sure the insecticide is applied correctly.
  - ◆ Use an effective (and labeled) insecticide at full labeled rate.
  - ◆ Use proper nozzles, spray volume (15 to 20 gallons per acre by ground; 3 to 5 gallons per acre by air) and pressure (40 pounds per square inch).
- ◆ Spray under favorable environmental conditions to promote efficacy and reduce drift.
- ◆ After applications, scout fields again after three to five days to ensure the product provided the level of management expected.
- If a field needs to be retreated due to a failure, **rotate to a different insecticide group** for the follow-up application. Tables 1 and 2 list insecticide groups, active ingredients and example trade names of products available for soybean aphid management.
  - ◆ For example, if a field was treated with a pyrethroid (Group 3A) and a follow-up insecticide application is needed, then an insecticide from different insecticide group, such as an organophosphate (Group 1B), should be selected.
  - ◆ Mixtures of insecticides (premixes or formulated mixtures) may be effective for pest suppression but generally are not preferred for insecticide-resistance management.
  - ◆ Insecticide seed treatments are not a viable answer to managing insecticide resistant aphid populations.

**Table 1.** Foliar insecticides with a single active ingredient that are labeled for soybean aphid management.

Group #	Group	Active Ingredient	Product Examples (Trade Names)
1A	Carbamates	methomyl	Lannate LV, Nudrin LV, others
1B	Organophosphates	acephate	Acephate 97, Orthene 97, others
		chlorpyrifos	Govern 4E, Hatchet, Lorsban 4E, Warhawk, others
		dimethoate	Dimethoate 4E, Dimate 4E, others
3A	Pyrethroids and Pyrethrins	alpha-cypermethrin	Fastac CS
		beta-cyfluthrin	Baythroid XL
		bifenthrin	Bifender FC, Beifenture EC, Brigade 2EC, Discipline 2EC, Sniper, Tundra EC, others
		cyfluthrin	Tombstone Helios
		deltamethrin	Delta Gold
		esfenvalerate	Asana, Zyrate, others
		gamma-cyhalothrin	Declare, Proaxis, Proaxis-Insecticide
		lambda-cyhalothrin	Grizzly Too, Karate, Lambda-Cy AG, LambdaStar, Province, Silencer, Warrior II, others
		permethrin	Permethrin, Perm-UP 3.2 EC, Arctic 3.2 EC, others
		zeta-cypermethrin	Mustang Maxx
4A	Neonicotinoids	acetamiprid	
		cholthianadin	Belay
		imidacloprid	Admire Pro, Alisa 4F, Nuprid 4F Max, others
		thiamethoxam	
4C	Sulfoxamines	sulfoxaflor	Transform
4B	Butenolides	flupyradifurone	Sivanto Prime
9D	Pyropenes	afidopyropen	Sefina

## Management Strategies (continued)

- Report suspected cases of insecticide-resistant soybean aphids to a local/regional Extension educator or Extension entomologist (see contact information on back). Before assuming resistance, try to rule out other potential causes for an insecticide failure (such as incorrect rate or application method, or unfavorable environmental conditions).

Until aphid-resistant soybean varieties and other management tactics become more widely available, cost-effective management of soybean aphid will continue to rely on scouting and threshold-based insecticide applications of the few labeled insecticide groups (Table 1).

This short list of insecticide groups is under threat of becoming even shorter through continued development of pest resistance to insecticides and potential regulatory actions. The agricultural community would be wise to work together to preserve the effectiveness of and continued access to these important tools for protection of crops from insect pests.

**Table 2.** Foliar insecticides with two or more active ingredients (often called premixes or formulated mixtures) labeled for soybean aphid management. Mixtures may be effective for pest suppression, but generally not preferred for insecticide resistance management.

IRAC Mode-of-Action Group #	Group	Active Ingredient	Premix (Trade Names)
28	Diamides	chlorantraniliprole	Besiege
3A	Pyrethroid	lambda-cyhalothrin	
3A	Pyrethroid	bifenthrin	Brigadier
4A	Neonicotinoid	imidacloprid	
1B	Organophosphate	chlorpyrifos	Cobalt Advanced
3A	Pyrethroid	lambda-cyhalothrin	
3A	Pyrethroid	lambda-cyhalothrin	Endigo ZC
4A	Neonicotinoid	thiamethoxam	
3A	Pyrethroid	bifenthrin	Hero
3A	Pyrethroid	zeta-cypermethrin	
4A	Neonicotinoid	acetamiprid	Justice
3A	Pyrethroid	bifenthrin	
3A	Pyrethroid	beta-cyfluthrin	Leverage 360
4A	Neonicotinoid	imidacloprid	
1B	Organophosphate	chlorpyrifos	Match-Up
3A	Pyrethroid	bifenthrin	
3A	Pyrethroid	bifenthrin	Skyraider, Swagger
4A	Neonicotinoid	imidacloprid	
1B	Organophosphate	chlorpyrifos	Stallion
3A	Pyrethroid	zeta-cypermethrin	
3A	Pyrethroid	bifenthrin	Tundra Supreme
1B	Organophosphate	chlorpyrifos	

## Insecticide Resistance Factors

Factors that likely led to the development of insecticide resistance in soybean aphids:

- Frequent infestations – Economically threatening infestations of soybean aphid have occurred more frequently in portions of Minnesota and neighboring states than in other parts of the soybean aphid's range, resulting in a long history of selection pressure for development of resistance.
- Limited number of insecticide groups for soybean aphid management (Table 1) – Management of soybean aphid has relied on foliar applications of only a few insecticide groups, mainly pyrethroids (Group 3A) and organophosphates (Group 1B), for more than 15 years.
- Misuse of insecticides can result in pests being exposed to insecticides more frequently, which further increases selection pressure. Examples of this include:
  - Application of insecticides when pest populations are below economic threshold.
  - Tank mixing insecticide with herbicide applications regardless of pest populations.
  - Applying insecticides below the labeled rate.
  - Repeated application of the same active ingredient or insecticide group within a single season.



## Contact Your Extension Entomologists:

- Minnesota: Robert Koch, University of Minnesota, email: koch0125@umn.edu, phone: 612-624-6771 [www.extension.umn.edu/agriculture/soybean/pest](http://www.extension.umn.edu/agriculture/soybean/pest)
- Iowa: Erin Hodgson, Iowa State University, email: ewh@iastate.edu, phone: 515-294-2847 [www.ent.iastate.edu/soybeanresearch/content/extension](http://www.ent.iastate.edu/soybeanresearch/content/extension)
- North Dakota: Janet Knodel, North Dakota State University, email: janet.knodel@ndsu.edu, phone: 701-231-7915 [www.ag.ndsu.edu/extensionentomology/field-crops-insect-pests/soybean](http://www.ag.ndsu.edu/extensionentomology/field-crops-insect-pests/soybean)
- South Dakota: Adam Varenhorst, South Dakota State University, email: adam.varenhorst@sdsu.edu, phone: 605-688-6854 <http://igrow.org/up/resources/03-2007-2017.pdf>




Photo courtesy of P. Beauzay, NDSU

## Other Resources:

Insecticide options for resistant soybean aphids, University of Minnesota Extension. <https://extension.umn.edu/soybean-pest-management/insecticide-options-resistant-soybean-aphid>

North Dakota Field Crop Insect Management Guide. NDSU Extension Service, E1143 (revised). [www.ag.ndsu.edu/pubs/plantsci/pests/e1143.pdf](http://www.ag.ndsu.edu/pubs/plantsci/pests/e1143.pdf)

South Dakota Pest Management Guide: Soybeans. South Dakota State University Extension and South Dakota Department of Agriculture. <http://igrow.org/up/resources/03-3042-2017.pdf>

Management Recommendations for Soybean Aphid (Hemiptera: Aphididae) in the United States. *Journal of Integrated Pest Management*. 3(1): 2012. <http://dx.doi.org/10.1603/IPM11019>

Biology and Economics of Recommendations for Insecticide-Based Management of Soybean Aphid. *Plant Health Progress* 17(4): 265-269. <http://dx.doi.org/10.1094/PHP-RV-16-0061>

**Always read, understand and follow the Environmental Protection Agency-approved label directions on the product container.**

Disclaimer: Insecticides are given as examples only and do not imply endorsement of one product versus another nor discrimination against any product not mentioned by the authors or the universities.

The NDSU Extension Service does not endorse commercial products or companies even though reference may be made to tradenames, trademarks or service names.

NDSU encourages you to use and share this content, but please do so under the conditions of our Creative Commons license. You may copy, distribute, transmit and adapt this work as long as you give full attribution, don't use the work for commercial purposes and share your resulting work similarly. For more information, visit [www.ag.ndsu.edu/agcomm/creative-commons](http://www.ag.ndsu.edu/agcomm/creative-commons).

**For more information on this and other topics, see [www.ndsu.edu/extension](http://www.ndsu.edu/extension)**

County commissions, North Dakota State University and U.S. Department of Agriculture cooperating. NDSU does not discriminate in its programs and activities on the basis of age, color, gender expression/identity, genetic information, marital status, national origin, participation in lawful off-campus activity, physical or mental disability, pregnancy, public assistance status, race, religion, sex, sexual orientation, spousal relationship to current employee, or veteran status, as applicable. Direct inquiries to Vice Provost for Title IX/ADA Coordinator, Old Main 201, NDSU Main Campus, 701-231-7708, [ndsu.eoaa@ndsu.edu](mailto:ndsu.eoaa@ndsu.edu). This publication will be made available in alternative formats for people with disabilities upon request, 701-231-7881.

Printing is being supported by



Published with support from:

