

**NDSU: Carrington Research Extension Center
2002 Variety Trial Data**

Oat - Dryland						Carrington
Variety	Days to Head	Plant Lodge 0-9	Plant Height inch	Leaf Disease %	Test Weight lb/bu	Grain Yield bu/ac
AC Assiniboia	56.8	3.0	38.1	6.5	32.9	75.6
AC Medallion	55.3	2.8	36.2	11.3	32.2	63.9
AC Ronald	54.3	2.5	32.8	9.3	34.5	68.9
Brawn	55.5	5.8	30.8	60.5	29.3	55.9
Buff	50.5	6.8	31.7	31.5	44.0	58.9
Ebeltoft	57.5	3.5	32.8	6.3	32.6	71.0
HiFi	55.3	2.5	33.6	5.5	33.1	72.5
Hyttest	52.8	6.8	35.6	76.5	37.4	58.3
Jerry	52.0	7.5	31.4	68.0	35.9	55.5
Jud	55.8	2.8	37.0	9.5	33.7	80.7
Killdeer	52.8	4.8	29.4	50.0	32.1	68.9
Loyal	56.8	2.5	35.8	5.0	35.2	68.9
Morton	54.3	1.3	38.6	11.5	34.2	64.6
Otana	54.8	7.3	37.0	83.0	29.2	54.7
Paul	55.8	3.0	36.6	30.0	36.0	43.4
Reeves	49.5	6.0	30.9	12.8	36.5	72.5
Sesqui	55.0	6.5	34.8	10.0	33.5	70.1
Triple Crown	58.3	3.8	38.2	13.3	24.5	40.9
Vista	53.5	6.0	33.6	6.8	35.4	72.2
Wabasha	50.5	8.0	32.2	10.5	34.4	67.9
Youngs	55.8	2.8	37.7	8.0	30.1	59.8
MEAN	54.3	4.0	34.9	20.9	34.1	66.5
C.V.%	1.3	29.8	6.7	20.0	3.8	12.5
LSD.05	1.0	1.7	3.3	5.9	1.8	11.6

Planting Date = May 10; Harvest Date = August 13; Previous Crop = Soybean

Leaf Disease = percent of flag leaf with disease symptoms (primarily crown rust) at medium milk.