

**NDSU Carrington Research Extension Center  
2004 Variety Trial Data**

<b>Dry Edible Bean - Irrigated</b>	<b>Carrington</b>
------------------------------------	-------------------

Variety	Market Class	----- Seeds per Pound ----			----- Seed Yield -----		
		2004	3-yr. Avg.	Seed Weight gram/100	Test Weight lb/bu	2004	3-yr. Avg. lb/ac
Buster	Pinto	1307	1334	34.9	59.8	1815	2062
Eclipse	Black	2610	--	17.5	62.7	1649	--
Erimo	Adzuki	3267	--	13.9	.	483	--
GTS 900	Pinto	1388	--	33.0	60.0	1471	--
Hokkai	Kintoki	571	--	79.5	59.0	2600	--
Maverick	Pinto	1358	1276	33.6	58.5	1868	2060
Navigator	Navy	2522	2493	18.0	63.7	1579	1793
Norstar	Navy	2574	2675	17.6	65.7	1459	1493
Othello	Pinto	1279	1269	35.7	60.1	1437	1800
Ralley	Pinto	1270	--	35.8	59.7	1635	--
Red Hawk	Dark Red Kidney	725	937	62.8	58.0	2155	1278
Seahawk	Navy	1922	--	23.7	65.0	1588	--
T-39	Black	2333	2407	19.5	62.0	1581	1775
Topaz	Pinto	1327	1325	34.3	57.5	1684	1377
Vista	Navy	2492	2622	18.2	63.5	1538	1835
Winchester	Pinto	1174	--	38.7	61.7	2165	--
MEAN		1738	--	31.4	61.1	1719	--
C.V.%		5.7	--	5.3	1.1	21.7	--
LSD.05		140	--	2.3	1.0	528	--

**Planting Date = May 28; Harvest Date = October 6; Previous Crop = Corn**

\*\* All varieties were significantly damaged by the frost that occurred on August 20th.

\*\* The variety Arthur was planted but no data is reported due to poor stand.