

# SUMMARY: 1983 COOPERATIVE CUTWORM PHEROMONE PROJECT

Department of Entomology

North Dakota State University

John D. Busacca and Dennis D. Kopp

| OBS | SITE       | DATE | EM  | EO | ET | EA | PS  | ST  | AO | LC  | PU | FJ | HE |
|-----|------------|------|-----|----|----|----|-----|-----|----|-----|----|----|----|
| 1   | BISMARCK   | 508  | 0   | 0  | 0  | 0  | 0   | 0   | 0  | 1   | 0  | 0  | .  |
| 2   | BISMARCK   | 515  | 0   | 0  | 0  | 0  | 0   | 0   | 0  | 3   | 0  | 0  | .  |
| 3   | BISMARCK   | 522  | 5   | 0  | 1  | 0  | 4   | 5   | 1  | 10  | 1  | 0  | .  |
| 4   | BISMARCK   | 529  | 10  | 0  | 1  | 1  | 15  | 6   | 0  | 10  | 3  | 0  | .  |
| 5   | BISMARCK   | 605  | 0   | 0  | 0  | 0  | 5   | 5   | 0  | 20  | 4  | 0  | .  |
| 6   | BISMARCK   | 612  | 1   | 0  | 1  | 1  | 3   | 4   | 0  | 20  | 1  | 0  | .  |
| 7   | BISMARCK   | 619  | 0   | 0  | 0  | 1  | 2   | 4   | 1  | 35  | 0  | 0  | .  |
| 8   | BISMARCK   | 626  | 0   | 0  | 8  | 0  | 6   | 1   | 1  | 67  | 0  | 0  | .  |
| 9   | BISMARCK   | 703  | 0   | 0  | 7  | 0  | 7   | 3   | 0  | 56  | 0  | 0  | .  |
| 10  | BISMARCK   | 710  | 0   | 0  | 5  | 0  | 8   | 0   | 0  | 60  | 0  | 0  | .  |
| 11  | BISMARCK   | 717  | 0   | 2  | 9  | 0  | 0   | 2   | 0  | 40  | 0  | 0  | .  |
| 12  | BISMARCK   | 724  | 0   | 6  | 7  | 1  | 0   | 5   | 0  | 20  | 0  | 0  | .  |
| 13  | BISMARCK   | 731  | 0   | 3  | 8  | 0  | 0   | 1   | 0  | 20  | 0  | 0  | .  |
| 14  | BISMARCK   | 807  | 0   | 1  | 6  | 0  | 0   | 2   | 0  | 14  | 0  | 0  | .  |
| 15  | BISMARCK   | 814  | 0   | 0  | 1  | 0  | 0   | 1   | 0  | 10  | 0  | 0  | .  |
| 16  | BISMARCK   | 821  | 0   | 1  | 0  | 0  | 0   | 2   | 0  | 5   | 0  | 0  | .  |
| 17  | BISMARCK   | 828  | 0   | 0  | 0  | 0  | 0   | 0   | 0  | 1   | 0  | 0  | .  |
| 18  | BISMARCK   | 904  | 0   | 0  | 0  | 0  | 0   | 0   | 0  | 2   | 0  | 0  | .  |
| 19  | BISMARCK   | 911  | 0   | 0  | .  | 0  | 0   | 0   | 0  | 0   | 1  | 0  | 0  |
| 20  | CARRINGTON | 515  | 0   | 0  | 0  | 0  | 100 | 0   | 0  | 35  | 0  | 0  | .  |
| 21  | CARRINGTON | 609  | 0   | 0  | 0  | 0  | 15  | 26  | 0  | 0   | 2  | 2  | .  |
| 22  | CARRINGTON | 620  | 0   | 0  | 0  | 0  | 7   | 53  | 0  | 5   | 2  | 0  | .  |
| 23  | CARRINGTON | 701  | 1   | 0  | 0  | 0  | 19  | 66  | 1  | 17  | 6  | 0  | .  |
| 24  | CARRINGTON | 715  | 8   | 1  | 23 | 0  | 6   | 16  | 1  | 4   | 3  | 4  | .  |
| 25  | CARRINGTON | 729  | 1   | 25 | 20 | 0  | 10  | 25  | 0  | 76  | 0  | 1  | .  |
| 26  | CARRINGTON | 805  | 3   | 43 | 2  | 0  | 0   | 10  | 0  | 7   | 0  | 1  | .  |
| 27  | CARRINGTON | 1006 | 487 | 54 | 0  | 2  | 42  | 112 | 0  | 103 | 0  | 10 | .  |
| 28  | COGSWELL   | 503  | 0   | 0  | 0  | 0  | 8   | 2   | 0  | 3   | 13 | 2  | .  |
| 29  | COGSWELL   | 509  | 0   | 0  | 0  | 0  | 15  | 0   | 0  | 9   | 4  | 1  | .  |
| 30  | COGSWELL   | 516  | 1   | 0  | 0  | 0  | 14  | 0   | 0  | 5   | 1  | 0  | .  |
| 31  | COGSWELL   | 523  | 0   | 0  | 0  | 0  | 2   | 0   | 0  | 3   | 0  | 1  | .  |

**Cutworm Project Continued:**

| OBS | SITE      | DATE | EM | EO | ET | EA | PS | ST | AO | LC  | PU | FJ | HE |
|-----|-----------|------|----|----|----|----|----|----|----|-----|----|----|----|
| 32  | COGSWELL  | 601  | 0  | 0  | 0  | 0  | 2  | 3  | 0  | 2   | 0  | 0  | .  |
| 33  | COGSWELL  | 606  | 0  | 0  | 0  | 0  | 1  | 1  | 0  | 2   | 0  | 0  | .  |
| 34  | COGSWELL  | 613  | 0  | 0  | 0  | 1  | 2  | 15 | 0  | 5   | 1  | 0  | .  |
| 35  | COGSWELL  | 620  | 0  | 0  | 0  | 0  | 1  | 2  | 2  | 3   | 0  | 0  | .  |
| 36  | COGSWELL  | 627  | 0  | 0  | 0  | 0  | 3  | 8  | 0  | 8   | 0  | 1  | .  |
| 37  | COGSWELL  | 705  | 2  | 0  | 9  | 0  | 4  | 7  | 0  | 10  | 0  | 0  | .  |
| 38  | COGSWELL  | 715  | 5  | 0  | 0  | 0  | 9  | 10 | 0  | 62  | 2  | 4  | .  |
| 39  | COGSWELL  | 718  | 2  | 0  | 41 | 0  | 1  | 2  | 0  | 123 | 0  | 0  | .  |
| 40  | COGSWELL  | 725  | 0  | 0  | 9  | 0  | 1  | 1  | 0  | 12  | 0  | 0  | .  |
| 41  | COGSWELL  | 801  | 2  | 1  | 9  | 0  | 0  | 3  | 0  | 35  | 0  | 0  | .  |
| 42  | COGSWELL  | 808  | 0  | 2  | 1  | 0  | 1  | 15 | 0  | 15  | 0  | 0  | .  |
| 43  | COGSWELL  | 817  | 8  | 0  | 0  | 0  | 1  | 7  | 0  | 9   | 0  | 0  | .  |
| 44  | COGSWELL  | 822  | 5  | 0  | 0  | 2  | 1  | 4  | 0  | 5   | 0  | 1  | .  |
| 45  | COGSWELL  | 829  | 18 | 0  | 0  | 1  | 2  | 4  | 0  | 3   | 1  | 2  | .  |
| 46  | COGSWELL  | 907  | 11 | 0  | 0  | 0  | 4  | 3  | 0  | 2   | 0  | 0  | .  |
| 47  | COGSWELL  | 911  | 2  | 0  | 0  | 1  | 1  | 0  | 0  | 8   | 1  | 0  | .  |
| 48  | COGSWELL  | 918  | 0  | 0  | 0  | 0  | 4  | 0  | 0  | 4   | 0  | 0  | .  |
| 49  | COGSWELL  | 1003 | 13 | 0  | 1  | 12 | 8  | 0  | 0  | 14  | 1  | 0  | .  |
| 50  | DICKINSON | 509  | 0  | 0  | 0  | 0  | 5  | 0  | 0  | 7   | 0  | 0  | 0  |
| 51  | DICKINSON | 516  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0   | 0  | 0  | 0  |
| 52  | DICKINSON | 523  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 5   | 0  | 1  | 0  |
| 53  | DICKINSON | 531  | 0  | 0  | 0  | 0  | 0  | 29 | 0  | 8   | 0  | 0  | 0  |
| 54  | DICKINSON | 606  | 2  | 0  | 0  | 0  | 0  | 27 | 0  | 3   | 0  | 0  | 0  |
| 55  | DICKINSON | 613  | 2  | 0  | 0  | 0  | 0  | 7  | 1  | 3   | 1  | 2  | 0  |
| 56  | DICKINSON | 620  | 11 | 1  | 0  | 0  | 0  | 17 | 0  | 17  | 0  | 27 | 0  |
| 57  | DICKINSON | 627  | 17 | 2  | 0  | 0  | 2  | 21 | 3  | 26  | 7  | 29 | 2  |
| 58  | DICKINSON | 705  | 15 | 2  | 37 | 0  | 3  | 47 | 12 | 115 | 13 | 16 | 6  |
| 59  | DICKINSON | 711  | 0  | 0  | 18 | 0  | 0  | 0  | 0  | 111 | 18 | 0  | 1  |
| 60  | DICKINSON | 718  | 0  | 0  | 66 | 0  | 0  | 41 | 0  | 357 | 4  | 0  | 2  |
| 61  | DICKINSON | 725  | 2  | 3  | 42 | 2  | 0  | 45 | 0  | 281 | 2  | 0  | 2  |
| 62  | DICKINSON | 801  | 0  | 3  | 36 | 0  | 1  | 8  | 0  | 122 | 0  | 0  | 0  |
| 63  | DICKINSON | 808  | 9  | 0  | 14 | 0  | 0  | 1  | 0  | 47  | 0  | 0  | 0  |
| 64  | DICKINSON | 815  | 45 | 0  | 0  | 0  | 0  | 27 | 0  | 3   | 0  | 0  | 2  |
| 65  | DICKINSON | 822  | 48 | 0  | 0  | 0  | 0  | 38 | 0  | 3   | 1  | 2  | 2  |
| 66  | DICKINSON | 830  | 39 | 0  | 0  | 0  | 1  | 25 | 2  | 6   | 0  | 2  | 3  |
| 67  | DICKINSON | 906  | 51 | 0  | 0  | 0  | 1  | 13 | 2  | 4   | 3  | 2  | 0  |
| 68  | DICKINSON | 913  | 20 | 2  | 0  | 0  | 1  | 10 | 0  | 3   | 12 | 0  | 1  |
| 69  | DICKINSON | 922  | 23 | 0  | 0  | 0  | 0  | 3  | 1  | 1   | 10 | 0  | 0  |
| 70  | HETTINGER | 602  | 0  | 0  | 0  | 0  | 23 | 5  | 0  | 14  | 0  | 0  | .  |
| 71  | HETTINGER | 613  | 0  | 0  | 0  | 0  | 5  | 1  | 0  | 3   | 1  | 0  | .  |

**Cutworm Project Continued:**

| OBS | SITE      | DATE | EM | EO  | ET | EA | PS | ST  | AO | LC  | PU | FJ | HE |
|-----|-----------|------|----|-----|----|----|----|-----|----|-----|----|----|----|
| 72  | HETTINGER | 620  | 1  | 0   | 0  | 0  | 7  | 5   | 0  | 4   | 4  | 0  | .  |
| 73  | HETTINGER | 627  | 4  | 0   | 0  | 0  | 3  | 2   | 0  | 1   | 6  | 9  | .  |
| 74  | HETTINGER | 705  | 13 | 0   | 0  | 0  | 4  | 7   | 0  | 2   | 10 | 4  | .  |
| 75  | HETTINGER | 711  | 2  | 0   | 2  | 0  | 5  | 1   | 3  | 7   | 7  | 3  | .  |
| 76  | HETTINGER | 718  | 4  | 0   | 3  | 0  | 8  | 3   | 5  | 7   | 6  | 5  | .  |
| 77  | HETTINGER | 725  | 4  | 0   | 15 | 0  | 10 | 3   | 2  | 13  | 3  | 3  | .  |
| 78  | HETTINGER | 801  | 3  | 7   | 13 | 0  | 3  | 0   | 0  | 9   | 0  | 0  | .  |
| 79  | HETTINGER | 807  | 1  | 7   | 8  | 0  | 4  | 0   | 0  | 5   | 0  | 1  | .  |
| 80  | HETTINGER | 906  | 25 | 3   | 0  | 2  | 4  | 8   | 0  | 6   | 1  | 2  | .  |
| 81  | HETTINGER | 913  | 1  | 0   | 0  | 1  | 2  | 1   | 2  | 2   | 0  | 0  | .  |
| 82  | HETTINGER | 921  | 0  | 0   | 0  | 3  | 0  | 0   | 0  | 1   | 0  | 0  | .  |
| 83  | HETTINGER | 927  | 0  | 0   | 1  | 9  | 0  | 0   | 0  | 0   | 0  | 0  | .  |
| 84  | LANGDON   | 509  | 0  | 0   | 0  | 0  | 0  | 0   | 0  | 0   | 0  | 0  | 0  |
| 85  | LANGDON   | 516  | 2  | 0   | 0  | 0  | 1  | 0   | 0  | 0   | 0  | 0  | 0  |
| 86  | LANGDON   | 523  | 3  | 0   | 0  | 0  | 3  | 0   | 0  | 0   | 0  | 0  | 0  |
| 87  | LANGDON   | 531  | 0  | 0   | 0  | 0  | 5  | 0   | 0  | 5   | 0  | 0  | 0  |
| 88  | LANGDON   | 606  | 0  | 0   | 0  | 0  | 5  | 3   | 0  | 2   | 0  | 0  | 0  |
| 89  | LANGDON   | 614  | 2  | 0   | 0  | 0  | 3  | 0   | 0  | 3   | 0  | 1  | 0  |
| 90  | LANGDON   | 621  | 0  | 0   | 0  | 0  | 2  | 7   | 0  | 3   | 0  | 4  | 0  |
| 91  | LANGDON   | 627  | 7  | 1   | 0  | 0  | 7  | 2   | 0  | 1   | 0  | 0  | 0  |
| 92  | LANGDON   | 705  | 11 | 0   | 1  | 0  | 0  | 0   | 0  | 0   | 8  | 10 | 0  |
| 93  | LANGDON   | 711  | 5  | 0   | 13 | 0  | 4  | 9   | 0  | 0   | 0  | 0  | 0  |
| 94  | LANGDON   | 718  | 0  | 0   | 28 | 0  | 0  | 2   | 0  | 0   | 0  | 0  | 0  |
| 95  | LANGDON   | 725  | 0  | 4   | 3  | 0  | 0  | 2   | 0  | 0   | 0  | 0  | 0  |
| 96  | LANGDON   | 801  | 0  | 30  | 72 | 0  | 0  | 8   | 0  | 0   | 0  | 0  | 0  |
| 97  | LANGDON   | 808  | 11 | 126 | 47 | 0  | 0  | 111 | 0  | 0   | 0  | 0  | 0  |
| 98  | LANGDON   | 815  | 29 | 32  | 33 | 0  | 0  | 35  | 0  | 0   | 0  | 0  | 0  |
| 99  | LANGDON   | 822  | 24 | 25  | 3  | 0  | 0  | 8   | 0  | 0   | 0  | 0  | 0  |
| 100 | LANGDON   | 829  | 30 | 58  | 0  | 0  | 2  | 0   | 0  | 0   | 0  | 0  | 0  |
| 101 | LANGDON   | 906  | 23 | 39  | 0  | 0  | 3  | 14  | 0  | 0   | 0  | 0  | 0  |
| 102 | LANGDON   | 912  | 3  | 7   | 0  | 0  | 0  | 0   | 0  | 0   | 0  | 0  | 0  |
| 103 | LANGDON   | 919  | 0  | 0   | 0  | 0  | 2  | 0   | 0  | 0   | 0  | 0  | 0  |
| 104 | LANGDON   | 926  | 1  | 0   | 0  | 0  | 5  | 0   | 0  | 0   | 0  | 0  | 0  |
| 105 | LANGDON   | 1003 | 0  | 0   | 0  | 0  | 0  | 1   | 0  | 0   | 0  | 0  | 0  |
| 106 | MINOT     | 527  | 1  | 0   | 0  | 0  | 16 | 0   | 0  | 12  | 1  | 0  | .  |
| 107 | MINOT     | 602  | 0  | 0   | 0  | 0  | 2  | 2   | 0  | 6   | 0  | 0  | .  |
| 108 | MINOT     | 608  | 0  | 1   | 0  | 0  | 6  | 0   | 0  | 2   | 0  | 0  | .  |
| 109 | MINOT     | 615  | 0  | 1   | 0  | 0  | .3 | 3   | 0  | 4   | 0  | 0  | .  |
| 110 | MINOT     | 623  | 3  | 1   | 0  | 0  | .2 | 1   | 0  | 11  | 7  | 6  | .  |
| 111 | MINOT     | 630  | 8  | 0   | 2  | 1  | 5  | 1   | 0  | 111 | 1  | 8  | .  |
| 112 | MINOT     | 708  | 13 | 0   | 4  | 0  | 8  | 2   | 0  | 100 | 6  | 8  | .  |

**Cutworm Project Continued:**

| OBS | SITE      | DATE | EM | EO | ET | EA | PS | ST | AO | LC  | PU | FJ | HE |
|-----|-----------|------|----|----|----|----|----|----|----|-----|----|----|----|
| 113 | MINOT     | 713  | 0  | 0  | 10 | 0  | 9  | 1  | 0  | 95  | 2  | 2  | 0  |
| 114 | MINOT     | 728  | 1  | 0  | 65 | 0  | 51 | 4  | 0  | 196 | 0  | 4  | 1  |
| 115 | MINOT     | 805  | 1  | 0  | 10 | 0  | 0  | 2  | 0  | 96  | 0  | 1  | 1  |
| 116 | MINOT     | 819  | 56 | 0  | 6  | 0  | 0  | 14 | 0  | 14  | 0  | 3  | 0  |
| 117 | MINOT     | 825  | 4  | 0  | 0  | 0  | 1  | 1  | 0  | 0   | 0  | 0  | 2  |
| 118 | MINOT     | 831  | 32 | 0  | 0  | 0  | 0  | 3  | 0  | 0   | 0  | 0  | 0  |
| 119 | WILLISTON | 505  | 0  | 0  | 0  | 0  | 2  | 0  | 0  | 0   | 0  | 0  | 0  |
| 120 | WILLISTON | 513  | 0  | 0  | 0  | 0  | 3  | 0  | 0  | 1   | 0  | 0  | 0  |
| 121 | WILLISTON | 520  | 0  | 0  | 0  | 0  | 1  | 0  | 0  | 0   | 0  | 0  | 0  |
| 122 | WILLISTON | 603  | 0  | 0  | 0  | 0  | 4  | 8  | 0  | 4   | 0  | 0  | 0  |
| 123 | WILLISTON | 609  | 0  | 0  | 0  | 0  | 2  | 11 | 0  | 3   | 0  | 0  | 0  |
| 124 | WILLISTON | 616  | 1  | 0  | 0  | 0  | 1  | 6  | 0  | 1   | 8  | 1  | 0  |
| 125 | WILLISTON | 623  | 3  | 0  | 0  | 0  | 0  | 6  | 0  | 0   | 2  | 0  | 0  |
| 126 | WILLISTON | 630  | 4  | 0  | 5  | 0  | 6  | 7  | 0  | 20  | 8  | 6  | 0  |
| 127 | WILLISTON | 707  | 1  | 1  | 8  | 0  | 2  | 4  | 1  | 49  | 2  | 2  | 0  |
| 128 | WILLISTON | 714  | 2  | 1  | 18 | 0  | 6  | 3  | 2  | 46  | 0  | 0  | 0  |
| 129 | WILLISTON | 725  | 1  | 6  | 90 | 0  | 2  | 16 | 1  | 49  | 1  | 0  | 0  |
| 130 | WILLISTON | 728  | 1  | 3  | 22 | 0  | 0  | 10 | 0  | 11  | 0  | 0  | 0  |
| 131 | WILLISTON | 804  | 13 | 4  | 34 | 0  | 1  | 5  | 0  | 30  | 2  | 0  | 0  |
| 132 | WILLISTON | 812  | 31 | 5  | 14 | 0  | 0  | 5  | 0  | 2   | 1  | 3  | 1  |
| 133 | WILLISTON | 819  | 62 | 1  | 0  | 0  | 0  | 28 | 0  | 1   | 0  | 3  | 0  |
| 134 | WILLISTON | 826  | 47 | 3  | 0  | 0  | 2  | 12 | 0  | 3   | 0  | 1  | 2  |
| 135 | WILLISTON | 901  | 77 | 1  | 2  | 1  | 2  | 28 | 3  | 4   | 1  | 0  | 0  |
| 136 | WILLISTON | 1003 | 79 | 5  | 0  | 0  | 5  | 21 | 0  | 6   | 12 | 1  | 0  |

| Trap | Species Name                 | Common Name          |
|------|------------------------------|----------------------|
| Em   | <i>Euxoa messoria</i>        | darksided cutworm    |
| Eo   | <i>Euxoa ochrogaster</i>     | redbacked cutworm    |
| Et   | <i>Euxoa tessellata</i>      | striped cutworm      |
| Ea   | <i>Euxoa auxiliaris</i>      | army cutworm         |
| Ps   | <i>Peridroma saucia</i>      | variegated cutworm   |
| Sc   | <i>Scotogramma trifolii</i>  | clover cutworm       |
| Ao   | <i>Agrotis orthogonia</i>    | pale-western cutworm |
| Lc   | <i>Leucania commoides</i>    |                      |
| Pu   | <i>Pseudaletia unipuncta</i> | armyworm             |
| Fj   | <i>Feltia jaculifera</i>     | dingy cutworm        |
| He   | <i>Homoeosoma electellum</i> | sunflower moth       |

**Annual Trap Catch and Average Trap Catch of Various Moths Attracted to  
Pheromone Baited Traps in North Dakota**

|                    |           | Em   | Eo  | Et  | Ea  | Ps  | St  | Ao | Lc   | Pu  | Fj | He |
|--------------------|-----------|------|-----|-----|-----|-----|-----|----|------|-----|----|----|
| <b>Straubville</b> | 82        | 152  | 1   | 12  | 3   | --  | 41  | 1  | 293  | 34  | -- | 7  |
|                    | 83        | 77   | 3   | 66  | 5   | 89  | 97  | 6  | 338  | 33  | 12 | -- |
|                    | $\bar{X}$ | 115  | 2   | 39  | 4   | 89  | 69  | 4  | 316  | 34  | 12 | 7  |
| <b>Williston</b>   | 80        | 768  | 157 | 191 | 164 | 31  | 112 | 9  | 330  | --  | -- | -- |
|                    | 81        | 334  | 72  | 182 | 25  | 90  | 146 | 7  | 235  | 28  | -- | -- |
|                    | 82        | 79   | 34  | 144 | 20  | --  | 103 | 52 | 151  | 72  | -- | 2  |
|                    | 83        | 322  | 30  | 183 | 1   | 39  | 169 | 6  | 230  | 37  | 17 | 3  |
|                    | $\bar{X}$ | 376  | 73  | 175 | 53  | 53  | 133 | 19 | 236  | 46  | 17 | 3  |
| <b>Dickinson</b>   | 80        | 723  | 123 | 331 | 232 | 65  | 195 | 4  | 329  | --  | -- | -- |
|                    | 81        | 244  | 70  | 203 | 100 | 224 | 216 | 36 | 371  | 43  | -- | -- |
|                    | 82        | 634  | 24  | 173 | 6   | --  | 296 | 9  | 384  | 130 | -- | 1  |
|                    | 83        | 284  | 13  | 213 | 2   | 14  | 359 | 21 | 1122 | 62  | 81 | 21 |
|                    | $\bar{X}$ | 471  | 57  | 230 | 85  | 101 | 267 | 18 | 552  | 78  | 81 | 11 |
| <b>Langdon</b>     | 80        | 199  | 613 | 523 | 21  | 117 | 134 | 23 | 501  | --  | -- | -- |
|                    | 81        | 250  | 463 | 436 | 26  | 186 | 55  | 15 | 605  | 144 | -- | -- |
|                    | 82        | 165  | 198 | 117 | 0   | --  | 55  | 5  | 109  | 30  | -- | 3  |
|                    | 83        | 151  | 322 | 200 | 0   | 42  | 202 | 0  | 14   | 8   | 15 | 0  |
|                    | $\bar{X}$ | 191  | 399 | 319 | 12  | 115 | 112 | 11 | 307  | 61  | 15 | 3  |
| <b>Hettinger</b>   | 80        | 831  | 94  | 324 | 287 | 50  | 122 | 17 | 294  | --  | -- | -- |
|                    | 81        | 117  | 27  | 61  | 332 | 243 | 48  | 25 | 356  | 76  | -- | -- |
|                    | 82        | 98   | 9   | 101 | 12  | --  | 56  | 5  | 216  | 96  | -- | 18 |
|                    | 83        | 58   | 17  | 42  | 15  | 78  | 36  | 12 | 74   | 38  | 27 | -- |
|                    | $\bar{X}$ | 276  | 37  | 132 | 162 | 124 | 66  | 15 | 235  | 70  | 27 | 18 |
| <b>Bismarck</b>    | 80        | 57   | 4   | 56  | 0   | 6   | 27  | 0  | 57   | --  | -- | -- |
|                    | 81        | 151  | 21  | 225 | 10  | 159 | 179 | 7  | 307  | 23  | -- | -- |
|                    | 82        | 137  | 20  | 134 | 3   | --  | 102 | 2  | 432  | 37  | -- | 3  |
|                    | 83        | 16   | 13  | 54  | 4   | 50  | 41  | 2  | 395  | 9   | 0  | -- |
|                    | $\bar{X}$ | 90   | 15  | 117 | 4   | 72  | 87  | 2  | 298  | 23  | 0  | 3  |
| <b>Carrington</b>  | 82        | 834  | 174 | 166 | 8   | --  | 127 | 1  | 161  | 50  | -- | 1  |
|                    | 83        | 490  | 123 | 45  | 2   | 199 | 308 | 2  | 247  | 13  | 18 | -- |
|                    | $\bar{X}$ | 662  | 149 | 106 | 5   | 199 | 218 | 2  | 204  | 32  | 18 | 1  |
| <b>Minot</b>       | 80        | 1322 | 627 | 710 | 20  | 95  | 241 | 9  | 865  | --  | -- | -- |
|                    | 81        | 303  | 158 | 147 | 3   | 263 | 57  | 14 | 256  | 50  | -- | -- |
|                    | 82        | 235  | 49  | 238 | 1   | --  | 13  | 2  | 237  | 59  | -- | 0  |
|                    | 83        | 119  | 3   | 97  | 1   | 87  | 34  | 0  | 647  | 17  | 4  | 4  |
|                    | $\bar{X}$ | 495  | 212 | 298 | 8   | 148 | 86  | 8  | 501  | 42  | 4  | 2  |

## SUMMER FALLOW PRACTICES DEMONSTRATION

By

Duane D. Hauck

Assoc. Ext. Eg. Eng.

### INTRODUCTION:

Summer Fallow Practices Demonstration trials were conducted during the summer of 1983 in cooperation with the Dickinson and Williston Branch Experiment Stations. The objective of the project was to demonstrate different fallowing practices stressing the importance of maintaining a protective residue cover on the soil surface. Individual fields were four acres each allowing for the use of regular sized field equipment. Parameters monitored included fuel and labor requirements, soil moisture, soil fertility, and residue levels.

### PRACTICES DEMONSTRATED:

Four practices were demonstrated at each location. These were identified as Conventional Fallow, Stubble Mulch, Reduced Tillage, and Chemical Fallow. The treatments used on each practice are shown in Table 1.

**Table 1. Treatments Used in Fallowing Practices**

| PRACTICES           | TREATMENTS   |                                 |
|---------------------|--|---------------------------------|
|                     | WILLISTON  | DICKINSON                       |
| Conventional Fallow | Chisel Plow with sweeps (2x)<br>Rod Weeder (1x)                            | Field Cultivator (3x)           |
| Stubble Mulch       | Undercutter with 30" sweeps (3x)<br>Chisel Plow with sweeps (1x)           | Undercutter with 8' sweeps (3x) |
| Reduced Tillage     | Roundup & Broadleaf Herbicide & Surf. (1x)<br>Chisel Plow with sweeps (2x) |                                 |
| Chemical Fallow     | Roundup & Broadleaf Herbicide & Surf. (3x)                                 |                                 |

Spring started out cool causing slow weed growth initially. The first treatments were performed May 26 at Williston and June 8 at Dickinson. High temperatures and adequate moisture contributed to vigorous weed growth throughout the summer. The second treatments were performed the first week of July at both locations with the exception of the Stubble Mulch practice at Williston. Here, due to excessive weed growth, the treatment was performed a week earlier.

The third treatments at Williston involving Tillage were performed the last week of July with the Chemical Fallow being treated a week later. The Stubble Mulch practice at Williston required a fourth treatment on September 6.

At Dickinson, the third and final treatments on all four practices were performed on August 8. At this time, the Stubble Mulch and Reduced Tillage practices were very heavily infested with pigeon grass from 10-12 inches tall and heading. Treating these two practices a week to ten days earlier would have been desirable.

### RESIDUE LEVELS:

Residue levels were measured at both locations in the spring prior to any treatment and again in the fall after all treatments were completed by collecting the residue from a given area and weighing it. Percent surface cover was estimated at several locations on each practice by walking 100 steps and counting the number of times a designated point struck residue.

The Dickinson site was cropped with oats the previous year and had the straw baled off. This left about 750 lbs. of residue per acre in the spring prior to fallowing treatments, which was about a 60% surface cover.

The Williston site was cropped with spring wheat the previous year with the straw spread after harvest. This left about 2,000 lbs. of residue per acre in the spring, which was a 90-95% surface cover.

Residue levels were again measured in the individual practices at the end of the season (Table 2). In some cases due to additional weed growth, residue levels were actually higher at the end of the season than they were at the start.

**Table 2. Residue Levels on Fallow Practices**

|                                  | Williston |         | Dickinson |         |
|----------------------------------|-----------|---------|-----------|---------|
|                                  | Lbs/Ac.   | % Cover | Lbs/Ac.   | % Cover |
| <b>Spring – Before Treatment</b> | 2,000     | 90-95   | 750       | 60      |
| <b>Fall – After Treatment:</b>   |           |         |           |         |
| Conventional Fallow              | 750       | 50      | 140       | 8       |
| Stubble Mulch                    | 1,200     | 60      | 800       | 45      |
| Reduced Tillage                  | 1,000     | 58      | 600       | 30      |
| Chemical Fallow                  | 2,500     | 90      | 1,100     | 65      |

## SOIL FERTILITY:

Soil samples were taken at both locations in the spring and again in the fall after treatments were completed. The samples were fertility tested by the NDSU Soil Testing Lab. The results indicate that Nitrogen levels on all the fallow sites increased substantially from last spring and are also considerably higher than that on adjacent cropped ground (Table 3). Among the individual fallow sites at both locations the Conventional Fallow had the highest Nitrogen level, possibly due to better organic matter breakdown as a result of tillage burying the residue. The Stubble Mulch practice had the lowest Nitrogen levels at both locations, possibly due mainly to the excessive weed growth that existed.

**Table 3. Soil Fertility Levels**

|                     | Williston     |               |               | Dickinson     |               |               |
|---------------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                     | N             | P             | K             | N             | P             | K             |
|                     | Lb/Ac<br>0-2' | Lb/Ac<br>0-6" | Lb/Ac<br>0-6" | Lb/Ac<br>0-2' | Lb/Ac<br>0-6" | Lb/Ac<br>0-6" |
| <b>May 25</b>       | 26            | 14            | 550           | 24            | 17            | 375           |
| <b>Sept. 28:</b>    |               |               |               |               |               |               |
| Conventional Fallow | 114           | 11            | 470           | 102           | 13            | 380           |
| Stubble Mulch       | 65            | 11            | 520           | 56            | 10            | 235           |
| Reduced Tillage     | 110           | 8             | 570           | 78            | 11            | 270           |
| Chemical Fallow     | 83            | 15            | 710           | 72            | 12            | 340           |
| Cropped Ground      | 25            | 16            | 630           | 17            | 9             | 350           |

## SUMMARY:

This demonstration shows several positive aspects for using Conventional Fallowing Systems – less cost, higher Nitrogen levels, and similar moisture retention to other fallowing practices. However, the thing it doesn't show is the high erosion hazard that exists with this method.

At the Dickinson location where a small amount of residue existed at the start, the land was rolling and several hard rainstorms hit the area throughout the summer, there's no doubt that Chemical Fallow was worth the expense. A high level of soil loss was observed on all treatments using tillage, especially the Conventional Fallow.

At the Williston location, it was a different story. A large amount of residue existed at the start, the land was only gently rolling, and the rainfall through the summer was moderate. Here, even after three passes with a chisel plow (Conventional Fallow), enough residue was retained on the surface to provide adequate erosion protection. No erosion was observed on any of the practices. Consequently, the best fallow system to use in this situation was the one that had the lowest cost.

For a farmer, the decision on which fallowing system to use depends mostly on the residue level available at the start and the erodibility of the land being fallowed.



## FUEL AND LABOR REQUIREMENTS:

As stated earlier, fields were four acres in size so that full sized equipment could be used enabling records to be kept on fuel and labor requirements. The Experiment Station's equipment normally used for field work was used here except for the Chemical Fallow at Dickinson. Here, an 18' sprayer normally used for plot work was used. Fuel and labor requirements for the entire fallow season are shown in Table 4.

**Table 4. Fuel and Labor Requirements for Fallowing Practices**

|                     | Williston      |                 | Dickinson      |                 |
|---------------------|----------------|-----------------|----------------|-----------------|
|                     | Fuel<br>Gal/Ac | Labor<br>Hrs/Ac | Fuel<br>Gal/Ac | Labor<br>Hrs/Ac |
| Conventional Fallow | 2.37           | .38             | 1.77           | .40             |
| Stubble Mulch       | 3.49           | .53             | 2.17           | .55             |
| Reduced Tillage     | 1.78           | .31             | 1.88           | .50             |
| Chemical Fallow     | .52            | .15             | .48            | .50             |

The tillage equipment used at both locations was basically the same size. However, a larger tractor was used at Williston allowing for faster field yields and lower labor requirements. The sprayer used at Dickinson was smaller than that normally used for field work, which is the reason for the higher than expected labor requirements for the Chemical Fallow. The Fuel and Labor Requirements for the Stubble Mulch practice at Williston were considerably higher than the others mainly due to the additional treatment required. The Conventional Fallow at Dickinson had lower requirements than the Reduced Tillage Fallow because a field cultivator was used for all three treatments compared to using a chisel plow twice on the Reduced Tillage.

## SOIL MOISTURE:

Soil samples were taken to the four foot depth the week of May 23, before treatment, and again the week of September 26, after all treatments. Cropped ground adjacent to the fallow site was also sampled in the fall. The results are shown in Figure 1.

**PLEASE INSERT SOIL MOISTURE LEVEL ON FALLOW PRACTICES FIGURE.**

Soil bulk density readings were not taken so results are shown in percent soil moisture on a dry basis. The soil type within each field was extremely variable, especially at the Dickinson site. This allowed making general comparisons only between practices at each location. At both locations, the fallowed ground was considerably wetter than adjacent cropped fields. Soil moisture on the fallow sites to the 2' depth was actually higher in the spring prior to any fallowing treatments. Moisture from 2 to 4 feet went basically unchanged. Little soil moisture difference was observed between individual fallowing practices.

## COSTS:

A cost analysis was performed between the different fallowing practices. The treatments used on each practice at both locations are shown in Table 1. Herbicide rates were adjusted slightly depending on the weed situation. For the most part, a tank mix of  $\frac{1}{3}$  to  $\frac{3}{4}$  pts. Roundup plus  $\frac{1}{4}$  to  $\frac{1}{3}$  pts. Banvel or 2-4D plus .5% surfactant was used.

Two cost analyses were generated (Figure 2). The first shows strictly the variable costs: the cost of fuel, labor, and herbicides. This obviously has the greatest affect on a farmer's cash flow.


The second analysis included both Variable and Fixed Cost. Data here was obtained from Minnesota Extension Folder 589, "Minnesota Farm Machinery Economic Cost Estimates for 1983". This information uses new machinery prices and is given for several machine sizes. Information for a wide sweep machine and a rod weeder were not given so data for a similar sized chisel plow and field cultivator were used instead.

In the Variable Cost Analysis, fallowing practices utilizing herbicide treatment are considerably higher priced. The gap is narrowed considerably when Fixed Costs are figured in however; Chemical Fallow still has the highest cost.

**Figure 2. Cost Per Acre for the Entire Season**

| Practice            | Variable Costs          | Fixed and      |
|---------------------|-------------------------|----------------|
|                     | (Fuel, Labor, Chemical) | Variable Costs |
|                     | \$/Acre                 | \$/Acre        |
| <b>Dickinson:</b>   |                         |                |
| Conventional Fallow | 4.59                    | 12.75          |
| Stubble Mulch       | 5.99                    | 18.48          |
| Reduced Tillage     | 12.55                   | 21.50          |
| Chemical Fallow     | 27.21                   | 29.79          |
| <b>Williston:</b>   |                         |                |
| Conventional Fallow | 5.28                    | 16.57          |
| Stubble Mulch       | 7.54                    | 24.64          |
| Reduced Tillage     | 8.42                    | 18.71          |
| Chemical Fallow     | 22.56                   | 27.17          |
| Fuel @ \$1.25/gal.  |                         |                |
| Labor @ \$6.00/hr.  |                         |                |

FIGURE 1: SOIL MOISTURE LEVEL ON FALLOW PRACTICES



% Soil Moisture - Avg. to 4'

|                     | <u>Williston</u> | <u>Dickinson</u> |
|---------------------|------------------|------------------|
| SPRING              | 13.5             | 21.2             |
| Conventional Fallow | 11.6             | 20.5             |
| Chemical Fallow     | 10.8             | 20.6             |
| Reduced Till Fallow | 10.9             | 18.9             |
| Stubble Mulch       | 11.1             | 20.8             |
| Cropped             | 6.0              | 16.7             |

## **FIELD EVALUATION PLANTING: TECHNICAL REPORT – 1982**

- Project 38I316K:** North Dakota State University, Dickinson Branch Experiment Station, Dickinson, North Dakota
- Project Title:** Field Evaluation of Woody Plant Materials
- Introduction:** There is a need to evaluate the performance of shrub and tree species/cultivars for windbreaks, wildlife, and recreational plantings under diverse soil and climatic conditions. To meet this need, field evaluation planting sites representative of the major land resource areas were located in the three states served by the center. These sites provide planting locations for assemblies of trees and shrubs to be evaluated under uniform culture and management.
- Objective:** The objective is to assemble and evaluate woody plant materials for conservation use. Superior cultivars will be selected and released for increase by commercial nurseries.
- Cooperators:** The Soil Conservation Service, Plant Materials Center, Bismarck, North Dakota, in cooperation with the North Dakota State University, Dickinson Branch Experiment Station, Dickinson, North Dakota.
- Location:** This project is located one mile west of Dickinson, North Dakota, on the NDSU Dickinson Branch Experiment Station.
- Legal description: NE ¼ 5, T139N., R96W., Stark County, North Dakota
- Major Land Resource Area:** The site is located in Major Land Resource Area 054, Rolling Soft Shale Plain. This moderately dissected rolling plain is underlain by calcareous shales and sandstones. Strongly dissected areas of sharp local relief or badland topography border major streams and valleys in some areas. Elevation is 1,800 to 3,100 feet. Sixty percent of the area is rangeland.
- Soils:** The soil type is a Parshall fine sandy loam. The Parshall series consists of deep, well drained soils formed in fine sandy loam alluvium on terraces and outwash plains and in upland swales. The surface layer and subsoil is dark grayish-brown fine sandy loam. The underlying material is dark grayish-brown fine sandy loam and loamy fine sand. Permeability is moderately rapid. The available water capacity is moderate. Organic matter is high and fertility is medium.

This soil is in North Dakota Windbreak Suitability Group 5. Included in this group are nearly level to hilly soils of the Flaxton, Lihen, Livonia, Parshall and Vebar series among others. These are well-drained, loamy and sandy soils. They are suited to windbreak and other plantings, but selection of species is limited. Erosion hazard is serious. The moderate available water capacity is the main limitation.

Climate: For MLRA 054 the average annual precipitation is 13 to 19 inches; increasing from west to east for this semiarid area. Rainfall is highest from late spring to midsummer and very low during the rest of the year. Winter precipitation is snow. Average annual temperature is 40° to 45°F. Average freeze-free period is 110 to 135 days. The plant hardiness zone is 4a, with an average annual minimum temperature of -30 to -20°F.

### **Methods and Materials**

Assembly: Refer to Table 31 for a list of woody species planted from 1978 through 1982.

Planting Plan: The planting site is approximately 500 feet long and 200 feet wide. The area is divided into four blocks. Each block consists of single row, non-replicated plots. Each plot contains a minimum of 5 plants. Row length is 100 feet and spacing between rows is 20 feet. Block 1 contains conifers spaced 5 feet within row. Block 2 contains shrubs and small trees spaced 5 feet within row. Block 3 contains medium sized trees, spaced 10 feet within row. Block 4 contains tall trees spaced 10 feet within row. All rows run from west to east.

Plot Preparation: A clean, firm planting site is prepared annually by disking and harrowing.

Planting Method: All tree and shrubs were hand planted using approved forestry methods.

Planting Date: Refer to Table 31 for a list of woody species planted from 1978 through 1982.

Fertilization: No fertilizer has been applied to planting area.

Weed Control: No herbicide has been applied to any plot during year of establishment or in succeeding years. Weeds were controlled by clean cultivating between rows, within row, and in fallow areas. Six to seven tillage operations were performed each year in the months of May through August. Hand hoeing was done as needed to control weeds in rows.

Biological Control: Previous years: No animal repellent or insecticide was applied in 1978. In the fall of 1979, an animal repellent, Arasan 50, was sprayed on fruit trees to discourage rodent damage.

1980 – 1981: On November 6, 1980 and October 29, 1981, Arasan 50 was applied to the trunks and lower limbs of fruit trees to deter rodents from damaging bark and cambium. Conifers also received this spray treatment to discourage animal browse. No insecticides were applied.

1982: No rodent repellent was applied.

Irrigation: Each year, newly planted materials were watered with a portable tank. No water was added following year of establishment.

Crop Residue Management: No cover crop has been established.

Silvicultural Practices: A minimum of pruning was done in 1979 to reshape trees damaged by animals. Dead trees and broken branches were cut and removed each year for sanitation. Replacements were used when available.

Evaluations and Measurements: Previous years: Records of planting date, survival, vigor, canopy width, height, cold hardiness, animal damage, and unusual or outstanding features have been maintained since 1978.

1982: Climatic data recorded at Dickinson Branch Experiment Station, Dickinson, North Dakota is shown in Table 30.

Plant performance data was reported on SCS-ECS-58 Woody Plant Initial Evaluation sheets. Survival, vigor, canopy cover and height, and special remarks were recorded for all hardwoods and remaining conifers on May 5 and September 16, 1982.

## Results

Plant Performance: Mean data for individual accessions of trees and shrubs is shown in Table 31. The following accessions exhibit potential for further evaluation:

| Accession Number             | Genus/Species Origin/Source  | Plot Location           | Remarks |
|------------------------------|--|-------------------------|---------|
| ND-1765<br>5980T             | Siberian larch<br><u>Larix sibirica</u><br>USDA, FS, Shelterbelt Lab.,<br>Bottineau, ND                                  | 1/03/1-10               |         |
| ND-313                       | Red tatarian honeysuckle<br><u>Lonicera tatarica sibirica</u><br>USDA, ARS, Cheyenne, WY<br>USDA, SCS, PMC, Bismarck, ND | 2/01/1-10               |         |
| ND-1730<br>5994T             | Red tatarian honeysuckle<br><u>Lonicera tatarica sibirica</u><br>Lincoln-Oakes Nursery,<br>Bismarck, ND                  | 2/01/11-20              |         |
| ND-628<br>5887T              | Silverberry<br><u>Elaeagnus commutata</u><br>Wells, Co., ND  | 2/02/1-10               |         |
| ‘Midwest’                    | Manchurian crabapple<br><u>Malus baccata mandshurica</u><br>Echo Manchuria<br>Res. Sta. Morden, Manitoba, Canada         | 3/01/1-5                |         |
| WY-843<br>‘Bighorn’<br>4646T | Skunkbush sumac<br><u>Rhus trilobata</u><br>Basin, WY<br>USDA, SCS, PMC, Bismarck, ND                                    | 2/02/11-20<br>2/04/1-10 |         |
| ‘Red Splendor’<br>6004T      | Flowering crabapple<br><u>Malus sp. x</u><br>Lee Nursery, Fertile, MN  | 3/01/6-10               |         |



| Accession Number               | Genus/Species<br>Origin/Source  | Plot Location | Remarks |
|--------------------------------|---|---------------|---------|
| ND-14                          | Harbin pear<br><u>Pyrus ussuriensis</u><br>Res. Sta. Morden, MB, Canada<br>SCS, PMC, Bismarck, ND | 3/02/6-10     |         |
| SD-134<br>6066T                | Apricot<br><u>Prunus armeniaca</u><br>Brookings Co., Brookings, SD                                | 3/04/1-5      |         |
| ND-416<br>6067T                | Apricot<br><u>Prunus armeniaca</u><br>Burleigh Co., Bismarck, ND                                  | 3/05/6-10     |         |
| ND-1336<br>6088T               | Chokecherry<br><u>Prunus virginiana</u><br>Mercer Co., Stanton, ND                                | 3/06/6-10     |         |
| ND-1873<br>5648T               | Amur maple<br><u>Acer ginnala</u><br>Lincoln-Oakes Nursery,<br>Bismarck, ND                       | 3/09/1-5      |         |
| SD-156<br>5890T                | Green ash<br><u>Fraxinus pennsylvanica</u><br>Deuel Co., Clear Lake, SD                           | 4/01/1-5      |         |
| 'Cardan'<br>MDN-12002<br>5895T | Green ash<br><u>Fraxinus pennsylvanica</u><br>USDA, ARS, Mandan, ND<br>Carlyle, MT                | 4/02/1-5      |         |
| ND-1759<br>5893T               | Green ash<br>SD-156 x MDN-12002<br><u>Fraxinus pennsylvanica</u><br>USDA, SCS, PMC, Bismarck, ND  | 4/02/6-10     |         |
| ND-364<br>5867T                | Russian olive<br><u>Elaeagnus angustifolia</u><br>Burleigh Co., Menoken, ND                       | 4/06/1-5      |         |

The following accessions failed to survive:

| <b>Accession Number</b> | <b>Genus/Species Origin/Source</b>  | <b>Plot Location</b> | <b>Remarks</b>         |
|-------------------------|---|----------------------|------------------------|
| ND-1717<br>6045T        | Scotch pine<br><u>Pinus sylvestris</u><br>USDA, FS, For. Sci. Lab.,<br>Lincoln, NE<br>Pieria, Greece              | 1/10/1-5             | Failed to<br>Establish |
| ND-1760<br>6035T        | Engelman spruce<br><u>Picea engelmannii</u><br>Coeur D' alene, ID<br>USDA, FS, Shelterbelt Lab.,<br>Bottineau, ND | 1/05/1-5             | Failed to<br>Establish |
| ND-1719<br>6047T        | Scotch pine<br><u>Pinus sylvestris</u><br>USDA, FS, For. Sci. Lab.,<br>Lincoln, NE<br>Prague, Czechoslovakia      | 1/05/6-10            | Failed to<br>Establish |
| ND-1710<br>4364T        | <u>Pinus nigra</u><br>USDA, FS, For. Sci. Lab.<br>Lincoln, NE<br>Turkey   | 1/06/1-5             | Failed to<br>Establish |
| ND-1712<br>6040T        | <u>Pinus nigra</u><br>USDA, FS, For. Sci. Lab.<br>Lincoln, NE<br>Turkey   | 1/06/6-10            | Failed to<br>Establish |
| ND-1714<br>6039T        | <u>Pinus nigra</u><br>USDA, FS, For. Sci. Lab.<br>Lincoln, NE<br>Turkey   | 1/07/1-5             | Failed to<br>Establish |
| ND-1716<br>6041T        | <u>Pinus nigra</u><br>USDA, FS, For. Sci. Lab.<br>Lincoln, NE<br>Kellog Forest, Michigan                          | 1/08/1-5             | Failed to<br>Establish |

## Accessions which failed to survive (Continued):

| Accession Number | Genus/Species<br>Origin/Source  | Plot Location | Remarks             |
|------------------|---|---------------|---------------------|
| ND-1720<br>6037T | <u>Pinus densiflora</u><br>USDA, FS, For. Sci. Lab.<br>Lincoln, NE<br>Tono, Japan                                   | 1/08/6-10     | Failed to Establish |
| ND-1722<br>6093T | <u>Pseudotsuga menzeisii</u><br>USDA, For. Sci. Lab.<br>Lincoln, NE<br>Douglas Co., Colorado                        | 1/09/6-10     | Failed to Establish |
| ND-1723<br>6232T | Northern white cedar<br><u>Thuja occidentalis</u><br>USDA, FS, For. Sci. Lab.<br>Lincoln, NE<br>Quebec, Canada      |               | Failed to Establish |
| ND-3805          | Amur corktree<br><u>Phellodendron amurense</u><br>UM, Dept. of Hort. Arboretum<br>Chaska, MN<br>SDSU, Brookings, SD |               | Failed to Establish |
| ND-81<br>6078T   | Sloe<br><u>Prunus spinosa</u><br>Res. Sta. Morden, MB, Canada<br>USDA, SCS, PMC, Bismarck, ND                       | 3/08/1-5      | Winter Killed       |
| ND-573<br>5967T  | Cathay walnut<br><u>Juglans cinerea</u><br>Res. Sta. Morden, MB, Canada   | 4/04/1-5      | Failed to Establish |

Rabbit damage was severe the winter of 1981-1982. Nearly all species showed some signs of rabbit damage. The following accessions of Apricot were completely girdled and were pruned back to the ground May 12, 1982, to stimulate regrowth.

| <b>Accession Number</b> | <b>Genus/Species Origin/Source</b>   | <b>Plot Location</b> |
|-------------------------|--|----------------------|
| SD-132<br>6064T         | Apricot<br><u>Prunus armeniaca</u><br>Brookings Co., Brookings, SD<br>USDA, SCS, PMC, Bismarck, ND | 3/03/1-5             |
| SD-133<br>6065T         | Apricot<br><u>Prunus armeniaca</u><br>Brookings Co., Brookings, SD<br>USDA, SCS, PMC, Bismarck, ND | 3/03/6-10            |
| SD-134<br>6066T         | Apricot<br><u>Prunus armeniaca</u><br>Brookings Co., Brookings, SD<br>USDA, SCS, PMC, Bismarck, ND | 3/04/1-5             |
| Mantoy<br>6069T         | Apricot<br><u>Prunus armeniaca</u><br>USDA, ARS, Mandan, ND<br>USDA, SCS, PMC, Bismarck, ND        | 3/04/6-10            |
| ND-1178<br>6070T        | Apricot<br><u>Prunus armeniaca</u><br>Walsh Co., Park River, ND<br>USDA, SCS, PMC, Bismarck, ND    | 3/05/1-5             |
| ND-416<br>6067T         | Apricot<br><u>Prunus armeniaca</u><br>Burleigh Co., Bismarck, ND<br>USDA, SCS, PMC, Bismarck, ND   | 3/05/6-10            |
| ND-423<br>6068T         | Apricot<br><u>Prunus armeniaca</u><br>Stark Co., Dickinson, ND<br>USDA, SCS, PMC, Bismarck, ND     | 3/06/1-5             |

Project No: 38I316K

**Table 30. 1982 Weather Summary – Official Station, North Dakota State University  
Dickinson Branch Experiment Station, Dickinson, North Dakota**

| <b>Month</b> | <b>Temp.<br/>(Mean)</b> | <b>Normal<br/>Temp.<br/>(Mean)</b> | <b>Deviation<br/>From Norm.</b> | <b>Total<br/>Precip.</b> | <b>Normal<br/>Precip.</b> | <b>Deviation<br/>From Norm.</b> |
|--------------|-------------------------|------------------------------------|---------------------------------|--------------------------|---------------------------|---------------------------------|
| January      |                         | 10.6°F                             |                                 | 0.85 in.                 | 0.43 in.                  | +0.42 in.                       |
| February     |                         | 15.5                               |                                 | 0.40                     | 0.41                      | -0.01                           |
| March        |                         | 24.3                               |                                 | 1.68                     | 0.72                      | +0.96                           |
| April        |                         | 40.5                               |                                 | 1.85                     | 1.42                      | +0.43                           |
| May          |                         | 52.2                               |                                 | 4.32                     | 2.36                      | +1.96                           |
| June         |                         | 61.3                               |                                 | 3.48                     | 3.56                      | -0.13                           |
| July         |                         | 68.4                               |                                 | 2.02                     | 2.15                      | -0.13                           |
| August       |                         | 67.5                               |                                 | 2.63                     | 1.78                      | +0.85                           |
| September    |                         | 55.8                               |                                 | 1.77                     | 1.32                      | +0.45                           |
| October      |                         | 45.2                               |                                 | 6.51                     | 0.91                      | +5.60                           |
| November     |                         | 28.4                               |                                 | 0.63                     | 0.52                      | +0.11                           |
| December     |                         | <u>15.6</u>                        |                                 | <u>0.49</u>              | <u>0.41</u>               | <u>+0.08</u>                    |
| Annual       |                         | 40.4                               |                                 | 26.58                    | 15.99                     | +10.59                          |

|      | <b>Last Killing Frost</b> | <b>First Killing Frost</b> | <b>Frost Free Days</b> |
|------|---------------------------|----------------------------|------------------------|
| 1982 | May 7                     | September 15               | 131                    |

Total Seasonal Precipitation (Average): 11.27 inches

Total Seasonal Precipitation (April-August) 1982: 14.25 inches

USDA, SCS, PMC, Bismarck, North Dakota

210 – Project No: 38I316K

Project Title: Field Evaluation of Woody Plant Materials (FEP)

Location: North Dakota State University, Dickinson Branch Experiment Station, Dickinson, North Dakota

Major Land Resource Area: 054

202 – Soil Series Texture: Parshall fine sandy loam

201 – Year of Record: 1982

---

|                               |                            |                |                                   |
|-------------------------------|----------------------------|----------------|-----------------------------------|
| 211-PLOT-LOCATION             |                            | 304-MATL-PLTD  | (Establishment, material planted) |
| 23,1-ACC-NO                   | (Prime-PMC-control number, | 306-AGE        | (Age of stock)                    |
| 2-PLANT SYMBOL                | PI number)                 | 305-NO-PLTS    | (Number of plants)                |
| 12-COMMON-NAME                |                            | 310-NO-PLT-SRV | (Number of plants surviving)      |
| 4-GENUS-NAME                  |                            | 363-PCT-SRV    | (Percent survival)                |
| 5-SPECIES-NAME                |                            | 337-VI         | (Vigor, plant)                    |
| 29,30-COLL-SITE-STATE, COUNTY | (Origin/Source)            | 347-R-CO       | (Resistance to cold)              |
| 212-YR-PLT                    | (Year planted)             | 359-CAN-COV    | (Canopy cover, cm)                |
| 209-TRANS-DATE                | (Transplant date)          | 360-PLNT-HT    | (Plant height, cm)                |

---

Table 31. 38I316K Field Evaluation of Woody Plant Materials – Dickinson, ND – 1982

| Plot Location | Accession Number                        | Plant Symbol | Genus/Species Origin/Source  | Yr. Plt. | Trans. Date | Matl. Pltd. | Age | No. Plts. | No. Plt. Srv. | Pct. Srv. | V I | C O | Can. Cov. (cm) | Plnt. Ht. (cm) | Remarks       |
|---------------|---|--------------|--|----------|-------------|-------------|-----|-----------|---------------|-----------|-----|-----|----------------|----------------|---------------|
| 1/01/1-10     | ND-1729<br>5979T                        | LASI*        | Siberian larch<br><u>Larix sibirica</u><br>NDFS State Nursery<br>Towner, ND  | 78       | 05/16       | PLBR        | 1-0 | 10        | 8             | 80        | 8   |     | 29             | 46             |               |
| 1/02/1-10     | SL-383-T<br>Pallet No.<br>2392<br>5976T | LASI*        | Siberian larch<br><u>Larix sibirica</u><br>USDA, FS, Shelterbelt Lab.,<br>Bottineau, ND<br>Denbigh Ex. Forest                        | 78       | 05/16       | PLBR        | 1-0 | 10        | 9             | 90        | 6   |     | 47             | 69             |               |
| 1/03/1-10     | ND-1765<br>5980T                        | LASI*        | Siberian larch<br><u>Larix sibirica</u><br>USDA, FS, Shelterbelt Lab.,<br>Bottineau, ND  | 78       | 05/17       | PLBR        | 2-0 | 10        | 10            | 100       | 5   |     | 63             | 122            |               |
| 1/04/1-5      | ND-1763<br>6043T                        | PIPO*        | Ponderosa pine<br><u>Pinus ponderosa</u> var. <u>ponderosa</u><br>USDA, FS, Shelterbelt Lab.,<br>Bottineau, ND<br>757-5 Todd Co., SD | 78       | 05/16       | CONT        | 1-1 | 5         | 4             | 80        | 7   |     | 74             | 134            | Winter Injury |
| 1/04/6-10     | ND-1565<br>6036T                        | PIAR         | Bristle cone pine<br><u>Pinus aristata</u><br>USDA, FS, Shelterbelt Lab.,<br>Bottineau, ND   | 78       | 05/16       | CONT        | 1-1 | 5         | 1             | 20        |     |     | 65             | 90             |               |
| 1/06/1-10     | ND-1863<br>5909T                        | GLTR         | Honey locust<br><u>Gleditsia triacanthos</u><br>Brown Co., SD<br>USDA, SCS, PMC,<br>Bismarck, ND                                     | 82       | 05/12       | PLBR        | 2-0 | 10        | 9             | 90        | 5   |     | 33             | 46             |               |

| Plot Location | Accession Number             | Plant Symbol | Genus/Species<br>Origin/Source   | Yr. Plt. | Trans. Date | Matl. Pltd. | Age | No. Plts. | No. Plt. Srv. | Pct. Srv. | V I | C O | Can. Cov. (cm) | Plnt. Ht. (cm) | Remarks                                |
|---------------|------------------------------|--------------|--|----------|-------------|-------------|-----|-----------|---------------|-----------|-----|-----|----------------|----------------|--|
| 1/07/1-10     | ND-3804                      | ROPS         | Black locust<br><u>Robinia pseudoacacia</u><br>Darby, MT<br>ND Forest Service<br>State Nursery, Towner, ND               | 82       | 05/12       | CONT        | 1-0 | 10        | 7             | 70        | 5   |     | 53             | 76             |  |
| 2/01/1-10     | ND-313<br>5996T              | LOTAS*       | Red tatarian honeysuckle<br><u>Lonicera tatarica sibirica</u><br>USDA, ARS, Cheyenne, WY<br>USDA, SCS, PMC, Bismarck, ND | 78       | 05/17       | PLBR        | 2-0 | 10        | 10            | 100       | 4   |     | 162            | 136            | Moderate grasshopper damage, leaf spot |
| 2/01/11-20    | ND-1730<br>5994T             | LOTAS*       | Red tatarian honeysuckle<br><u>Lonicera tatarica sibirica</u><br>Lincoln-Oakes Nursery,<br>Bismarck, ND                  | 78       | 05/17       | PLBR        | 2-0 | 10        | 10            | 100       | 4   |     | 181            | 160            | Leaf spot, and mildew                  |
| 2/02/1-10     | ND-628<br>5877T              | ELCO*        | Silverberry<br><u>Elaeagnus commutata</u><br>Wells Co., ND   | 78       | 05/17       | PLBR        | 2-0 | 10        | 10            | 100       | 5   |     | 151            | 145            |  |
| 2/02/11-20    | WY-843<br>'Bighorn'<br>4646T | RHTR         | Skunkbush sumac<br><u>Rhus trilobata</u><br>USDA, SCS, PMC, Bismarck, ND<br>Bighorn Co., WY                              | 78       | 05/17       | PLBR        | 2-0 | 10        | 10            | 100       | 3   |     | 232            | 153            |  |
| 2/03/1-10     | ND-26<br>11852T              | *            | Honeysuckle<br><u>Lonicera</u> sp.<br>USDA, ARS<br>Mandan, ND  | 79       | 05/2        | PLBR        | 2-0 | 10        | 10            | 100       | 6   |     | 104            | 118            | Many grasshoppers                      |
| 2/03/11-15    | ND-452<br>19978T             | LOXYM*       | Honeysuckle<br><u>Lonicera xylosteum mollis</u><br>USDA, ARS, Cheyenne, WY<br>USDA, SCS, PMC, Bismarck, ND               | 79       | 05/2        | PLBR        | 2-0 | 5         | 5             | 100       | 4   |     | 133            | 137            | Leaf mildew, grasshoppers              |



| Plot Location | Accession Number             | Plant Symbol | Genus/Species Origin/Source  | Yr. Plt. | Trans. Date | Matl. Pltd.   | Age | No. Plts. | No. Plt. Srv. | Pct. Srv. | V I | C O | Can. Cov. (cm) | Plnt. Ht. (cm) | Remarks                       |
|---------------|------------------------------|--------------|--|----------|-------------|---------------|-----|-----------|---------------|-----------|-----|-----|----------------|----------------|-------------------------------|
| 2/04/1-10     | WY-843<br>'Bighorn'<br>4646T | RHTR         | Skunkbush sumac<br><u>Rhus trilobata</u><br>USDA, SCS, PMC, Bismarck, ND<br>Bighorn Co., WY  | 79       | 05/2        | PLBR          | 2-0 | 10        | 10            | 100       | 5   |     | 126            | 93             | Leaf spots                    |
| 2/04/11-20    | PM-ND-283<br>6079T           | PRTE*        | Russian almond<br><u>Prunus tenella</u><br>ND Fish & Game Dept.<br>USDA, SCS, PMC, Bismarck, ND  | 80       | 05/08       | PLBR          | 2-0 | 10        | 10            | 100       | 4   |     | 54             | 60             | Leaf spots                    |
| 2/05/1-10     | ND-11<br>5993T               | LOMA6        | Amur honeysuckle<br><u>Lonicera maackii</u><br>Res. Sta. Morden, MB, Canada  | 81       | 05/07       | CONT          | 0-1 | 10        | 10            | 100       | 4   |     | 42             | 44             |                               |
| 2/06/1-5      | ND-995<br>PI-303584          | SAHU         | Prairie willow<br><u>Salix humilis</u><br>USDA, PI Sta.,<br>Ames, IA   | 82       | 05/12       | PLBR-<br>CONT | 1-2 | 5         | 4             | 80        | 4   |     | 58             | 66             |                               |
| 2/06/6-10     | PI-370126                    | SALIX        | Willow<br><u>Salix sp.</u>   | 82       | 05/12       | PLBR-<br>CONT | 0-1 | 5         | 5             | 100       | 4   |     | 33             | 48             |                               |
| 2/07/1-10     | ND-624<br>6094T              | PTTR         | Common hop tree<br><u>Ptelea trifoliata</u><br>Ramsey Co., ND<br>USDA, SCS, PMC,<br>Bismarck, ND   | 82       | 05/12       | PLBR          | 2-0 | 10        | 9             | 90        | 5   |     | 24             | 33             |                               |
| 3/01/1-5      | 'Midwest'<br>6003T           | MABAM*       | Manchurian crabapple<br><u>Malus baccata mandshurica</u><br>Echo Manchuria/Res. Sta.<br>Morden, MB, Canada<br>USDA, SCS, PMC, Bismarck, ND | 78       | 05/17       | PLBR          | 2-0 | 5         | 5             | 100       | 3   |     | 144            | 169            | Deer browse,<br>rodent damage |

| Plot Location | Accession Number        | Plant Symbol | Genus/Species Origin/Source  | Yr. Plt. | Trans. Date | Matl. Pltd. | Age | No. Plts. | No. Plt. Srv. | Pct. Srv. | V I | C O | Can. Cov. (cm) | Plnt. Ht. (cm) | Remarks        |
|---------------|-------------------------|--------------|--|----------|-------------|-------------|-----|-----------|---------------|-----------|-----|-----|----------------|----------------|----------------|
| 3/01/6-10     | 'Red Splendor'<br>6004T | *            | Flowering crabapple<br><u>Malus sp.</u> x<br>Lee Nursery, Fertile, MN  | 78       | 05/17       | PLBR        | 2-0 | 5         | 5             | 100       | 3   |     | 181            | 256            |                |
| 3/02/1-5      | ND-1731<br>6001T        | MABA*        | Siberian crabapple<br><u>Malus baccata</u><br>Lincoln-Oakes Nursery,<br>Bismarck, ND   | 78       | 05/17       | PLBR        | 2-0 | 5         | 5             | 100       | 3   |     | 178            | 249            | 2 replacements |
| 3/02/6-10     | ND-14<br>1095T          | PYUS*        | Harbin pear<br><u>Pyrus ussuriensis</u><br>Harbin, Manchuria/Res. Sta.<br>Morden, MB, Canada<br>USDA, SCS, PMC, Bismarck, ND | 78       | 05/17       | PLBR        | 2-0 | 5         | 5             | 100       | 3   |     | 195            | 272            | Good growth    |
| 3/03/1-5      | SD-132<br>6064T         | PRAR*        | Apricot<br><u>Prunus armeniaca</u><br>USDA, SCS, PMC, Bismarck, ND<br>Brookings Co., SD                                      | 78       | 05/17       | PLBR        | 2-0 | 5         | 5             | 100       | 5   |     | 159            | 183            | 2 replacements |
| 3/03/6-10     | SD-133<br>6065T         | PRAR*        | Apricot<br><u>Prunus armeniaca</u><br>USDA, SCS, PMC, Bismarck, ND<br>Brookings Co., SD                                      | 78       | 05/17       | PLBR        | 2-0 | 5         | 3             | 60        | 6   |     | 185            | 185            |                |
| 3/04/1-5      | SD-134<br>6066T         | PRAR*        | Apricot<br><u>Prunus armeniaca</u><br>USDA, SCS, PMC, Bismarck, ND<br>Brookings Co., SD                                      | 78       | 05/17       | PLBR        | 2-0 | 5         | 5             | 100       | 7   |     | 124            | 146            |                |
| 3/04/6-10     | 'Mantoy'<br>6069T       | PRAR*        | Apricot<br><u>Prunus armeniaca</u><br>USDA, SCS, PMC, Bismarck, ND<br>USDA, ARS, Mandan, ND                                  | 78       | 05/17       | PLBR        | 2-0 | 5         | 5             | 100       | 6   |     | 195            | 212            |                |

| Plot Location | Accession Number     | Plant Symbol | Genus/Species<br>Origin/Source   | Yr. Plt. | Trans. Date | Matl. Pltd. | Age | No. Plts. | No. Plt. Srv. | Pct. Srv. | V I | C O | Can. Cov. (cm) | Plnt. Ht. (cm) | Remarks                                 |
|---------------|----------------------|--------------|--|----------|-------------|-------------|-----|-----------|---------------|-----------|-----|-----|----------------|----------------|---|
| 3/05/1-5      | ND-1178<br>6070T     | PRAR*        | Apricot<br><u>Prunus armeniaca</u><br>USDA, SCS, PMC, Bismarck, ND<br>Walsh Co., ND              | 78       | 05/17       | PLBR        | 2-0 | 5         | 4             | 80        | 7   |     | 168            | 161            |   |
| 3/05/6-10     | ND-416<br>6067T      | PRAR*        | Apricot<br><u>Prunus armeniaca</u><br>USDA, SCS, PMC, Bismarck, ND<br>Burleigh Co., ND           | 78       | 05/17       | PLBR        | 2-0 | 5         | 5             | 100       | 7   |     | 142            | 137            |   |
| 3/06/1-5      | ND-423<br>6068T      | PRAR*        | Apricot<br><u>Prunus armeniaca</u><br>USDA, SCS, PMC, Bismarck, ND<br>Stark Co., ND              | 78       | 05/17       | PLBR        | 2-0 | 5         | 5             | 100       | 7   |     | 155            | 190            |   |
| 3/06/6-10     | ND-1336<br>6088T     | PRVI         | Chokecherry<br><u>Prunus virginiana</u><br>Mercer Co., ND  | 78       | 05/17       | PLBR        | 2-0 | 5         | 5             | 100       | 3   |     | 259            | 313            | Slight leaf spot                        |
| 3/07/1-5      | ND-1732<br>6090T     | PRVI         | Chokecherry<br><u>Prunus virginiana</u><br>Lincoln-Oakes Nursery,<br>Bismarck, ND                | 78       | 05/17       | PLBR        | 2-0 | 5         | 5             | 100       | 4   |     | 247            | 293            | Moderate<br>mildew, slight<br>leaf spot |
| 3/07/6-10     | 'Schubert'<br>12608T | PRVI         | Chokecherry<br><u>Prunus virginiana</u><br>USDA, ARS, Mandan, ND<br>USDA, SCS, PMC, Bismarck, ND | 78       | 05/17       | PLBR        | 2-0 | 5         | 5             | 100       | 6   |     | 116            | 169            | Mildew                                  |
| 3/08/1-5      | ND-81<br>6078T       | PRSP         | Sloe<br><u>Prunus spinosa</u><br>Res. Sta. Morden, MB, Canada<br>USDA, SCS, PMC, Bismarck, ND    | 78       | 05/17       | PLBR        | 2-0 | 5         | 0             | 0         |     |     |                |                |   |

| Plot Location | Accession Number  | Plant Symbol | Genus/Species Origin/Source  | Yr. Plt. | Trans. Date | Matl. Pltd. | Age | No. Plts. | No. Plt. Srv. | Pct. Srv. | V I | C O | Can. Cov. (cm) | Plnt. Ht. (cm) | Remarks        |
|---------------|-------------------|--------------|--|----------|-------------|-------------|-----|-----------|---------------|-----------|-----|-----|----------------|----------------|----------------|
| 3/08/6-10     | ND-629<br>5645T   | ACGI         | Amur maple<br><u>Acer ginnala</u><br>Res. Sta. Morden, MB, Canada                        | 79       | 05/2        | PLBR        | 2-0 | 5         | 4             | 80        | 5   |     | 121            | 138            | 3 replacements |
| 3/09/1-5      | ND-1873<br>5648T  | ACGI         | Amur maple<br><u>Acer ginnala</u><br>Lincoln-Oakes Nursery,<br>Bismarck, ND              | 79       | 05/2        | PLBR        | 2-0 | 5         | 5             | 100       | 5   |     | 166            | 193            | 2, 4-D damage  |
| 3/09/6-10     | ND-686<br>6225T   | SYAMJ*       | Japanese tree lilac<br><u>Syringa amurensis japonica</u><br>ND Game & Fish Dept.         | 79       | 05/2        | PLBR        | 2-0 | 5         | 3             | 60        | 4   |     | 62             | 77             |                |
| 3/10/1-5      | ND-3773<br>21576T |              | Willow<br><u>Salix sp.</u><br>Norman Co., MN<br>USDA, SCS, PMC, Bismarck,<br>ND          | 82       | 05/12       | PLBR        | 0-1 | 5         | 3             | 60        | 7   |     | 11             | 22             |                |
| 3/10/6-10     | Mich-433          | SAPE         | Laurel willow<br><u>Salix pentandea</u><br>USDA, SCS, Rose Lake<br>PMC, East Lansing, MI | 82       | 05/12       | PLBR        | 0-1 | 5         | 5             | 100       | 5   |     | 13             | 38             |                |
| 4/01/1-5      | SD-156<br>5890T   | FRPE         | Green ash<br><u>Fraxinus pennsylvanica</u><br>Deuel Co., SD                              | 78       | 05/17       | PLBR        | 2-0 | 5         | 5             | 100       | 3   |     | 171            | 232            | 3 replacements |
| 4/01/6-10     | ND-1734<br>5891T  | FRPE         | Green ash<br><u>Fraxinus pennsylvanica</u><br>Lincoln-Oakes Nursery,<br>Bismarck, ND     | 78       | 05/17       | PLBR        | 2-0 | 5         | 5             | 100       | 4   |     | 143            | 222            | 2 replacements |

| Plot Location | Accession Number   | Plant Symbol | Genus/Species Origin/Source   | Yr. Plt. | Trans. Date | Matl. Pltd.   | Age | No. Plts. | No. Plt. Srv. | Pct. Srv. | V I | C O | Can. Cov. (cm) | Plnt. Ht. (cm) | Remarks                   |
|---------------|--------------------|--------------|---|----------|-------------|---------------|-----|-----------|---------------|-----------|-----|-----|----------------|----------------|---------------------------|
| 4/02/1-5      | MDN-12002<br>5895T | FRPE         | Green ash<br><u>Fraxinus pennsylvanica</u><br>USDA, ARS, Mandan, ND<br>Wibaux Co., MT               | 78       | 05/17       | PLBR          | 2-0 | 5         | 5             | 100       | 3   |     | 228            | 308            | 5 replacements            |
| 4/02/6-10     | ND-1759<br>5893T   | FRPE         | Green ash<br><u>Fraxinus pennsylvanica</u><br>SD-156 x MDN-12002<br>USDA, SCS, PMC,<br>Bismarck, ND | 78       | 05/17       | PLBR          | 2-0 | 5         | 5             | 100       | 4   |     | 176            | 246            | 5 replacements            |
| 4/03/1-5      | ND-647<br>5887T    | FRNI         | Black ash<br><u>Fraxinus nigra</u><br>Res. Sta. Morden, MB, Canada                                  | 78       | 05/17       | PLBR          | 2-0 | 5         | 5             | 100       | 4   |     | 126            | 243            | Good growth               |
| 4/03/6-10     | ND-1432<br>5658T   | AEGL         | Ohio buckeye<br><u>Aesculus glabra</u><br>Res. Sta. Morden, MB, Canada                              | 78       | 05/17       | PLBR          | 2-0 | 5         | 1             | 20        | 6   |     | 45             | 65             |                           |
| 4/04/1-5      | ND-1879<br>11850T  | GLSI         | Chinese honey locust<br><u>Gleditsia sinensis</u><br>Woodward, OK<br>USDA, ARS, Mandan, ND          | 80       | 05/08       | PLBR-<br>CONT | 2-1 | 5         | 5             | 100       | 4   |     | 43             | 68             |                           |
| 4/04/6-10     | ND-548<br>5969T    | JUMA*        | Manchurian walnut<br><u>Juglans mandshurica</u><br>Res. Sta. Morden, MB, Canada                     | 78       | 05/17       | PLBR          | 2-0 | 5         | 3             | 60        | 6   |     | 168            | 110            | Winter injury             |
| 4/05/1-5      | ND-1170<br>6009T   | MOAL         | Mulberry<br><u>Morus alba</u><br>Burleigh Co., ND   | 78       | 05/17       | PLBR          | 2-0 | 5         | 5             | 100       | 3   |     | 420            | 325            | Moderate<br>Winter injury |
| 4/05/6-10     | ND-363<br>5866T    | ELAN         | Russian olive<br><u>Elaeagnus angustifolia</u><br>Burleigh Co., ND                                  | 78       | 05/17       | PLBR          | 2-0 | 5         | 5             | 100       | 4   |     | 240            | 296            |                           |

| Plot Location | Accession Number     | Plant Symbol | Genus/Species Origin/Source  | Yr. Plt. | Trans. Date | Matl. Pltd. | Age | No. Plts. | No. Plt. Srv. | Pct. Srv. | V I | C O | Can. Cov. (cm) | Plnt. Ht. (cm) | Remarks |
|---------------|----------------------|--------------|--|----------|-------------|-------------|-----|-----------|---------------|-----------|-----|-----|----------------|----------------|---------|
| 4/06/1-5      | ND-364<br>5867T      | ELAN         | Russian olive<br><u>Elaeagnus angustifolia</u><br>Burleigh Co., ND                       | 78       | 05/17       | PLBR        | 2-0 | 5         | 5             | 100       | 2   |     | 375            | 400            |         |
| 4/06/6-10     | ND-1735<br>5874T     | ELAN         | Russian olive<br><u>Elaeagnus angustifolia</u><br>Lincoln-Oakes Nursery,<br>Bismarck, ND | 78       | 05/17       | PLBR        | 2-0 | 5         | 5             | 100       | 4   |     | 335            | 441            |         |
| 4/07/1-5      | ND-541<br>5868T      | ELAN         | Russian olive<br><u>Elaeagnus angustifolia</u><br>Haakon Co., SD                         | 78       | 05/17       | PLBR        | 2-0 | 5         | 5             | 100       | 3   |     | 390            | 360            |         |
| 04/07/6-10    | PM-ND-1843<br>11840T | ELAN         | Russian olive<br><u>Elaeagnus angustifolia</u><br>Res. Sta. Morden, MB, Canada           | 80       | 05/08       | PLBR        | 2-0 | 5         | 5             | 100       | 4   |     | 144            | 153            |         |
| 4/09/1-10     | MDN-12003<br>T05725  | CEOC         | Hackberry<br><u>Celtis occidentalis</u><br>USDA, ARS, Mandan, ND                         | 80       | 05/08       | PLBR        | 2-0 | 10        | 8             | 80        | 6   |     | 40             | 48             |         |
| 4/10/1-10     | PM-SD-75<br>5713T    | CEOC         | Hackberry<br><u>Celtis occidentalis</u><br>Potter Co., SD                                | 81       | 05/07       | PLBR        | 2-0 | 10        | 7             | 70        | 6   |     | 28             | 44             |         |