

Barley Disease by Location, Year and Variety																			
Location	Foliar Necrosis - % of Flag						DON - ppm								FHB Field Severity - %				
	4 Site	L	Lnl	Llo	L	P	7 Site	LV	LI	L	T	R	P	L	4 Site	LV	L	Lnl	Llo
Year	Ave.	06	04	04	03	03	Ave.	07	07	05	05	05	05	04	Ave.	07	05	04	04
Variety:																			
Six-Rowed																			
Robust	40	15	50	70	23	5	4.7	1	7	5	9	6	5	1	3	2	1	1	7
Drummond	19	3	25	35	14	3	4.9	1	8	4	9	8	4	1	--	--	1	0	1
Lacey	34	14	33	65	24	8	4.0	1	4	7	5	6	5	1	--	--	2	0	3
Legacy	34	17	40	60	20	5	3.9	1	6	5	7	4	2	2	2	1	2	1	5
Tradition	26	2	30	55	16	3	3.3	1	5	3	7	6	1	1	1	1	1	0	2
Stellar-ND	43	6	55	85	27	--	4.5	1	7	7	13	3	2	1	1	2	1	0	0
Two-Rowed																			
Pinnacle	--	--	--	--	--	--	--	1	3	--	--	--	--	--	--	--	--	--	--
Bowman	75	8	100	100	91	--	--	1	--	2	--	--	--	1	--	--	1	1	2
Conlon	51	18	70	40	75	--	--	0	2	3	--	--	--	0	1	2	2	0	1
CDC Copeland	--	--	--	--	--	--	--	0	--	--	--	--	--	--	--	--	--	--	--
Scarlett	--	--	--	--	--	--	--	0	--	--	--	--	--	--	--	--	--	--	--
Rawson	26	13	10	60	22	--	--	1	--	4	--	--	--	0	--	--	1	0	1
AC Metcalfe	--	4	20	70	--	--	--	0	--	2	--	--	--	1	--	--	0	0	3

L=Langdon, Lnl=Langdon not lodged, Llo=Langdon badly lodged, LI=Langdon Irrigated, P=Pembina, W=Walsh, T=Towner, R=Ramsey, N=Nelson
 Field Severity = (Incidence x Head Severity) 2007 ratings on selected varieties.