

North Central Research Extension Center—Minot
Camelina Variety Trial (spring)

Variety	Early Vigor	Days	Days	Days	Plant Height	Seed Weight g/1000	Test Weight lb/bu	Oil ¹ %	Seed Yield		
		10% Flower	90% Flower	Maturity DAP ³					2007	2008	2009
Blaine Creek	2.5	44	65	94	32	1.1	51.5	35.3	585	1528	2110
Calena	3.0	45	63	94	33	1.5	51.5	36.3	--	1602	2541
Celine	2.5	46	63	95	33	1.2	51.0	35.1	469	1299	1975
Galena	2.8	44	61	93	33	1.5	52.1	36.0	593	2270	2613
Ligena	3.0	45	62	95	33	1.6	50.4	35.8	546	2043	2427
Robinson	3.0	44	62	93	32	1.3	51.6	35.3	204	1816	2266
Suneson	2.8	44	61	94	30	1.3	52.1	36.0	604	1820	1962
LSD 5%	NS	1	1	1	2	0.2	0.5	0.6	135	NS	326
C.V.%	24.0	1.2	1.4	0.9	5.4	11.1	0.7	1.2	21.8	20.9	10.1
Mean	2.7	44	62	94	32	1.4	51.3	35.7	434	1742	2215

Trial was planted on May 17 with a seeding rate of 7 million PLS and harvested on August 21.

¹ Oil adjusted to 8.5% moisture.

² Early Vigor: Vigor of plants 4 weeks after planting (1 = poorest, 5 = excellent).

³ DAP = Days after planting.