

# Field evaluation of fungicides for management of *Ascochyta* blight of chickpeas

Hofflund / Nesson Valley, ND –  
25 miles east of Williston (2012)

Tyler Tjelde, Irrigation Agronomist, NDSU Williston Res. Extension Ctr., Nesson Valley Irrigation Research Site  
Michael Wunsch, Plant Pathologist, NDSU Carrington Research Extension

## KEY FINDINGS:

- Proline (5.7 fl oz/ac) and Priaxor (4 or 6 fl oz/ac) showed excellent efficacy against *Ascochyta* blight on chickpeas.
- Priaxor performed similarly when applied at 4 fl oz/ac and 6 fl oz/ac.
- Rotational strategies with Proline and Priaxor performed equivalently as sequential applications of Proline.
- Omega (fluazinam), Bravo Top (difenoconazole + chlorothalonil), and Vertisan (penthiopyrad) showed efficacy against *Ascochyta* blight on chickpeas. Additional testing is needed to optimize the use of these products in a fungicide resistance management program. For Bravo Top and Vertisan, additional testing is needed confirm that their efficacy is equivalent to Proline and Priaxor, respectively, which they would replace in a fungicide resistance management program.

## DETAILED RESULTS:

The fungicides BRAVO TOP, OMEGA, and QUASH are currently not registered for use on chickpeas and should not be used. Future registration of these fungicides is anticipated, and results for these products are provided for reference only.

Treatment (fungicide application timing) <sup>w</sup>	Ascochyta Blight		Yield <sup>†</sup>	Test weight	Seed diameter <sup>y</sup>				
	Severity <sup>z‡</sup>	Severity <sup>z</sup>			> 10.3 mm	9.5-10.3 mm	8.7-9.5 mm	< 8.7 mm	
	July 7 <sup>x</sup>	July 19 <sup>x</sup>	lbs/ac	lbs/bu	percent	percent	percent	percent	
1 Non-treated check (water; A,B,C,D)	3 abc*	64 bcd *	2065 bc *	60.9 a *	0.0	5 bc *	24 d *	72 d *	
2 Priaxor 500SC 4 fl oz/ac (A,B,C,D)	1 ab	25 a	2930 ab	62.0 a	0.0	15 ab	49 abc	36 abc	
3 Priaxor 500SC 6 fl oz/ac (A,B,C,D)	1 ab	31 a	2932 ab	62.6 a	0.0	14 ab	47 abc	39 abc	
4 Proline 480SC 5.7 fl oz/ac + NIS 0.125% v/v (A,B,C,D)	2 ab	27 a	3437 a	62.1 a	0.0	14 ab	46 abc	40 abc	
5 Omega 500F 13.6 fl oz/ac (A,B,C,D)	2 ab	34 a	2921 ab	61.7 a	0.1	12 abc	41 bc	47 abc	
6 Omega 500F 8 fl oz/ac (A) / Priaxor 500SC 4 fl oz/ac (B,D) / Proline 480SC 5.7 fl oz/ac + NIS 0.125% v/v (C)	4 abc	41 abc	2773 abc	62.4 a	0.1	15 ab	47 abc	38 abc	
7 Omega 500F 13.6 fl oz/ac (A) / Priaxor 500SC 4 fl oz/ac (B,D) / Proline 480SC 5.7 fl oz/ac + NIS 0.125% v/v (C)	3 abc	40 abc	2877 ab	62.6 a	0.1	14 ab	42 abc	44 abc	
8 Omega 500F 16 fl oz/ac (A) / Priaxor 500SC 4 fl oz/ac (B,D) / Proline 480SC 5.7 fl oz/ac + NIS 0.125% v/v (C)	4 abc	39 abc	2468 abc	62.7 a	0.2	15 a	47 abc	37 abc	
9 Bravo WS 1.5 pt/ac (A) / Priaxor 500SC 4 fl oz/ac (B,D) / Proline 480SC 5.7 fl oz/ac + NIS 0.125% v/v (C)	3 abc	31 a	3137 a	63.4 a	0.1	10 abc	42 abc	47 abc	
10 Non-treated check water (A) / Priaxor 500SC 4 fl oz/ac (B,D) / Proline 480SC 5.7 fl oz/ac + NIS 0.125% v/v (C)	9 c	66 cd	2670 abc	62.1 a	0.0	11 abc	39 bc	50 bc	
11 Omega 500F 13.6 fl oz/ac (A,B) / Priaxor 500SC 4 fl oz/ac (C) / Proline 480SC 5.7 fl oz/ac + NIS 0.125% v/v (D)	5 bc	40 abc	2749 abc	61.4 a	0.0	13 abc	43 abc	44 abc	
12 Non-treated (A,B) / Priaxor 500SC 4 fl oz/ac (C) / Proline 480SC 5.7 fl oz/ac + NIS 0.125% v/v (D)	6 bc	75 d	1914 c	62.5 a	0.0	2 c	19 d	79 d	
13 Proline 480SC 5.7 fl oz/ac + NIS 0.125% v/v (A,C) / Priaxor 500SC 4 fl oz/ac (B,D)	3 abc	25 a	3378 a	61.4 a	0.1	17 a	48 abc	35 abc	
14 Proline 480SC 5.7 fl oz/ac + NIS 0.125% v/v (A,C) / Priaxor 500SC 6 fl oz/ac (B,D)	2 abc	30 a	3175 a	61.9 a	0.2	19 a	49 ab	32 ab	
15 Priaxor 500SC 4 fl oz/ac (A,C) / Proline 480SC 5.7 fl oz/ac + NIS 0.125% v/v (B,D)	3 abc	29 a	3095 ab	61.8 a	0.0	15 a	45 abc	40 abc	
16 Priaxor 500SC 6 fl oz/ac (A,C) / Proline 480SC 5.7 fl oz/ac + NIS 0.125% v/v (B,D)	4 bc	33 a	3209 a	62.3 a	0.1	11 abc	47 abc	41 abc	
17 Priaxor 500SC 4 fl oz/ac (A,C) / Bravo Top 4.59SC 2 pt/ac (B,D)	2 ab	32 a	3046 ab	63.0 a	0.1	13 abc	47 abc	40 abc	
18 Bravo Top 4.59SC 2 pt/ac (A,C) / Priaxor 500SC 4 fl oz/ac (B,D)	1 ab	38 ab	2708 abc	62.2 a	0.0	14 ab	42 abc	43 abc	
19 Proline 480SC 5.7 fl oz/ac + NIS 0.125% v/v (A,C) / Vertisan 1.67EC 20 fl oz/ac + NIS 0.25% v/v (B,D)	3 abc	32 a	3065 ab	61.8 a	0.1	15 ab	42 abc	43 abc	
20 Vertisan 1.67EC 20 fl oz/ac + NIS 0.25% v/v (A,C) / Proline 480SC 5.7 fl oz/ac + NIS 0.125% v/v (B,D)	6 bc	45 abc	2716 abc	62.2 a	0.1	11 abc	38 c	51 c	
21 Proline 480SC 5.7 fl oz/ac + Bravo WS 1.5 pt/ac + NIS 0.125% v/v (A,C) / Priaxor 500SC 4 fl oz/ac (B,D)	0 a	35 a	3071 ab	61.9 a	0.0	18 a	53 a	29 a	
22 Quash 50WDG 3 oz/ac + NIS 0.25% v/v (A,C) / Priaxor 500SC 4 fl oz/ac (B,D)	2 abc	41 abc	2640 abc	62.5 a	0.2	14 ab	43 abc	42 abc	
	F:	3.95	6.53	3.37	1.67	1.16	3.88	13.34	11.29
	P > F:	< 0.0001	< 0.0001	0.0003	0.0762	0.3320	< 0.0001	< 0.0001	< 0.0001
	CV:	37.7	26.9	1.91	1.38	205.9	29.7	10.0	15.4

<sup>z</sup> *Ascochyta* disease severity: Percent of the canopy exhibiting symptoms of *Ascochyta*.

<sup>y</sup> Seed size: Seed size was determined by assessing the percent (by weight) of a 200-gram seed sample that passed through sieves with round 26/64, 24/64, and 22/64-inch diameter holes.

<sup>x</sup> The chickpeas were in full bloom on July 7 and July 19.

<sup>w</sup> Fungicide application timing:

**Application A:** June 27, 2012 at 9:30-11:00 am. Chickpeas at bloom initiation (less than 10% of plants with at least one open blossom); canopy was closed; trace levels of *Ascochyta* present. Wind = 12-16 mph, temperature = 66-69°F, relative humidity = 40-45%.

**Application B:** July 10, 2012 at 2:00-4:00 pm. Wind = 9 to 10 mph, temperature = 84 to 91°F, relative humidity = 30 to 40%.

**Application C:** July 20, 2012 at 7:00-10:00 am. Wind = 6.5 to 9 mph, relative humidity = 60-88%, temperature = 69 to 75°F.

**Application D:** August 2, 2012 at 11:30 am - 1:00 pm. Wind = 7 to 8 mph, relative humidity = 41-47%, temperature = 78 to 82°F.

\* Within-column means followed by different letters are significantly different ( $P < 0.05$ ; Tukey multiple comparison procedure).

‡ In order to meet meet model assumptions of normality and homoskedasticity, analysis of variance was conducted on the natural-log transformation of disease severity and yield [ $\ln(x + 1)$  for data sets including values < 1.0; otherwise,  $\ln(x)$ ]. For ease of interpretation, treatment means are reported as the untransformed yield (lbs/ac) and disease severity (percent).

# Field evaluation of fungicides for management of *Ascochyta* blight of chickpeas

Hofflund / Nesson Valley, ND – 25 miles east of Williston (2012)

Tyler Tjelde, Irrigation Agronomist, NDSU Williston Research Extension Center, Nesson Valley Irrigation Research Site

Michael Wunsch, Plant Pathologist, NDSU Carrington Research Extension Center

701-652-2951 / michael.wunsch@ndsu.edu

## METHODS:

- **Location of trial:** NDSU Williston Research Extension Center, Nesson Valley Irrigation Research Site, Williston, ND.
- **GPS coordinates of research trial location:** 48.167,-103.104
- **Soil type:** Lihen - fine sandy loam
- **Soil preparation:** conventional tillage (fall - disked once, ripped twice, and mulched once; spring - harrowed once with a noble spring tooth harrow)
- **Seeding equipment:** double-disc seeder (plot cone seeder)
- **Variety:** CDC 'Frontier'
- **Experimental design:** randomized complete block      **Replicates:** 4
- **Seeded plot size:** 5 feet wide (center-to-center) x 18 feet long      **Harvested plot size:** 5 feet wide (center-to-center) x 14 feet long
- **Row spacing:** 7 inches      **Rows per plot:** 6      **Non-treated buffer plots were established between treatment plots.**
- **Previous crop:** durum wheat
- **Planting date:** May 2, 2012      **Seeding rate:** targeted plant population = 4.5 plants/square foot. Stand count assessments taken on May 29 showed an average stand count of 4.3 plants/square foot.
- **Seed treatment:** Cruiser 5FS 1.28 fl oz/cwt + ApronMaxxRTA 5.0 fl oz/cwt + Mertect 340F 2.04 fl oz/cwt
- **Fungicide application A:** June 27, 2012 at 9:30-11:00 am. Chickpeas at bloom initiation (less than 10% of plants with at least one open blossom); canopy was closed; trace levels of *Ascochyta* present. Wind = 12-16 mph, temperature = 66-69°F, relative humidity = 40-45%.
- **Fungicide application B:** July 10, 2012 at 2:00-4:00 pm. Wind = 9 to 10 mph, temperature = 84 to 91°F, relative humidity = 30 to 40%.
- **Fungicide application C:** July 20, 2012 at 7:00-10:00 am. Wind = 6.5 to 9 mph, relative humidity = 60-88%, temperature = 69 to 75°F.
- **Fungicide application D:** August 2, 2012 at 11:30 am - 1:00 pm. Wind = 7 to 8 mph, relative humidity = 41-47%, temperature = 78 to 82°F.
- **Fungicide application details:** Fungicides were applied with a 56-in hand boom with four equally spaced flat-fan Spraying Systems TeeJet 8002VS nozzles. Applications were made at 40 psi in 20 gal/ac water.
- ***Ascochyta* inoculation details:** To promote disease development, guard and buffer plots were inoculated with laboratory-grown pycnidiospores of *Ascochyta rabiei* at 11:15 to 11:45 pm on July 2. Spores of *A. rabiei* were grown on potato dextrose agar, suspended in water, and applied to the guard plots at a spore concentration of  $1.25 \times 10^6$  spores/ml and an application rate of 50 ml per plot. The spores were applied by tossing the spore solution over the center of each guard and buffer plot. To facilitate disease establishment, the chickpeas were irrigated with 0.5 inches of water earlier in the evening; the canopy was wet at the time of spore application.
- **Disease assessments:** *Ascochyta* severity was assessed as the percent of the canopy exhibiting *Ascochyta* disease symptoms. Severity was evaluated at four locations per plot.
- **Desiccation:** This trial was desiccated with paraquat (Gramoxone at 2 pts/ac + NIS at 2.5 oz/ac in 15 gallons water/ac) on August 21, 2012.
- **Harvest date:** September 7, 2012.
- **Seed size:** Seed diameter was determined by assessing the percent (by weight) of a 200-gram seed sample that passed through sieves with round 26/64, 24/64, and 22/64-inch diameter holes.
- **Statistical analysis:** Data were evaluated with analysis of variance. The assumption of constant variance was assessed by plotting residuals against predicted values, and the assumption of normality was assessed with a normal probability plot. To meet model assumptions of homoskedasticity, a systematic natural-log transformation  $[\text{LN}(x+1)]$  was applied to the July 5 disease severity data and a systematic square-root transformation was applied to the yield data. All other data met model assumptions. Single-degree-of-freedom contrasts were performed for all pairwise comparisons of isolates; to control the Type I error rate at the level of the experiment, the Tukey multiple comparison procedure was employed. Analyses were conducted with replicate and treatment as main factor effects, and they were implemented in PROC GLM of SAS (version 9.2; SAS Institute, Cary, NC).

## WE GRATEFULLY ACKNOWLEDGE:

This project was made possible with grants from the **Northern Pulse Growers Association** and the **North Dakota Department of Agriculture Crop Protection Product Harmonization Board and Registration Board**. Supplementary financial support was provided by the **BASF Corporation** and **ISK BioSciences**.

We gratefully acknowledge **Syngenta Crop Protection** for donating the seed treatment products Cruiser and Mertect used in this trial and **JM Grain** for helping us obtain seed of CDC Frontier chickpeas for use in this trial.

## IMPORTANT NOTICE:

- Fungicide performance can differ in response to which diseases are present, levels of disease when products are applied, environmental conditions, plant architecture and the susceptibility to disease of the chickpea variety planted, crop growth stage at the time of fungicide application, and other factors.
- This report summarizes fungicide performance as tested at the NDSU Williston Research Extension Center's Nesson Valley Irrigation Research Site in 2012 under the conditions partially summarized in the methods section (above).
- Fungicide efficacy may differ under other conditions; when choosing fungicides, always evaluate results from multiple trials.
- This report is shared for educational purposes and is not an endorsement of any specific products.