



Avian Nest Survival in a Patch-burn Grazing System

Cameron Duquette and Torre Hovick

North Dakota State University, School of Natural Resource Sciences, Fargo, N.D.

C.A. Duquette, NDSU

We are evaluating the effect of a patch-burn grazing management strategy on avian nest success. Our treatment structure consists of four replicates of the following: (1) season-long grazing, (2) season-long grazing with dormant-season patch burning (one-quarter of pasture) at a four-year return interval and (3) season-long grazing with dormant-season (one-eighth of pasture) and growing-season (one-eighth of pasture) patch burning at a four-year return interval. Here we present preliminary results following two years of study.

Introduction

Common range management practices focus on even utilization of forage by grazers. This grazing strategy produces a homogeneous vegetation structure and composition centered on the middle of the disturbance gradient (Fuhlendorf and Engle, 2001). In contrast, grassland species have evolved with a shifting mosaic of disturbance through the interaction of fire and grazing (Fuhlendorf and Engle, 2004).

In intact disturbance regimes, grazers preferentially select for high-quality forage in patches regenerating after fire (Vermiere et al., 2003). Selection for newly burned areas by grazers releases unburned patches from grazing pressure, resulting in biomass accumulation. This, in turn, increases the propensity of unburned patches to carry fire and perpetuate the fire cycle (Fuhlendorf and Engle, 2004).

In fire-adapted rangeland systems, an intact natural disturbance regime creates heterogeneous vegetation structure across the landscape. This diversity in habitat conditions maintains or promotes biodiversity in plants, arthropods, small mammals and birds (Doxon et al., 2011; Fuhlendorf et al., 2006; Fuhlendorf et al. 2010).

Patch-burn grazing also increases the temporal stability of grassland avian communities (Hovick et al., 2015). Through a shifting mosaic of vegetation structure, the application of fire and grazing (hereafter, patch-burn grazing) can provide habitat for species relying on diverse aspects of the disturbance gradient to complete their life histories (Fuhlendorf et al., 2009).

Traditional range management can be especially limiting to avian species that rely on vegetation structure characteristic of the far ends of the grazer utilization spectrum as part of their nesting strategy. Some examples include mountain plovers, which rely on sparse ground cover, and Le Conte's sparrows, which use areas with thick litter as part of their nesting strategy (Graul, 1975; Hovick et al., 2014).

When using a traditional management strategy, managers often achieve uniform grazing pressure through fencing and rapid rotation of grazers (Briske et al., 2011). This increased intensity of use by grazers for short time periods increases the risk of nest trampling (Bleho et al., 2014; Churchwell et al., 2008).

Woody encroachment also threatens rangeland systems subject to an inactive disturbance regime. Woody species can increase the incidence of predation and cowbird parasitism and reduce nesting cues for grassland species (Archer et al., 2017; Klug et al., 2010;

With, 1994).

In grassland avian species, woody encroachment has been shown to impact landscape-level species diversity and nesting success (Bakker, 2003; Coppedge et al., 2001; Sirami et al., 2009). Increases in grassland shrub cover also result in decreases in arthropod richness and abundance, which may impact the initiation timing and success of nesting attempts (van Hengstum et al., 2013).

We will study the use of experimental pastures by nesting birds during a time-since-fire gradient by monitoring nest success and density, as well as associated vegetation characteristics. Increases in within-patch homogeneity with accompanying heterogeneity between patches may create spatially explicit nesting habitat for a higher diversity of species, in turn creating more source habitat for grassland birds (Davis et al., 2016).

In addition, imposed heterogeneity should allow species to select for vegetation structure that maximizes nest success. Results will increase conservation of grassland bird species of conservation concern, such as the grasshopper sparrow (*Ammodramus* *savannarum*), Sprague's pipit (*Anthus spragueii*) and upland sandpiper (*Bartramia longicauda*).

Procedures

Study Area

The Central Grasslands Research Extension Center (CGREC) is in Kidder and Stutsman counties, N.D., (46° 42' 56" N, 99° 27' 08" W) in the Missouri Coteau ecoregion of the northern mixed-grass prairie. The herbaceous community is dominated by native cool-season grasses such as green needlegrass (*Nassella viridula*), western wheatgrass (*Pascopyrum smithii*) and needle-and-thread grass (*Heterostipa comata*).

Common invasive grasses on site include Kentucky bluegrass (*Poa pratensis*) and smooth brome (*Bromus inermis*) (Patton et al., 2007). Western snowberry (*Symphoricarpos occidentalis*) is the dominant woody species at the CGREC, although silverberry (*Eleagnus commutata*) and wild rose (*Rosa arkansana*) are present.

The forb community is diverse and dominated by western ragweed (*Ambrosia psilostachya*), prairie coneflower (*Ratibida columnifera*), goldenrod (*Solidago spp.*), yarrow (*Achillea millefolium*) and Flodman's thistle (*Cirsium flodmanii*) (Rogers et al., 2005). The climate is characterized as temperate and experiences an average yearly rainfall of 40.28 centimeters (cm) (15.9 inches) and average annual temperatures of 4.94 C (40.9 F) (1991-2016, North Dakota Agricultural Weather Network).

Treatment Structure

Our treatment structure consists of four replicates, each consisting of a 160-acre pasture divided into eight subpatches. The treatments are: (1) season-long grazing (SLG), (2) season-long grazing with dormant-season patch burning (one-fourth of pasture) at a four-year return interval (PBG40) and (3) season-long grazing with dormant-season (one-eighth of pasture) and

growing-season (one-eighth of pasture) patch burning at a four-year return interval (PBG20).

Annual burn plots in treatment 3 will be two adjacent 20-acre subpatches. Growing-season burns are incorporated to increase forage quality for livestock in the middle of the season (Scasta et al., 2016). Fire return intervals are designed to mimic the historical disturbance regime of mixed-grass prairie.

Cow-calf pairs will graze freely within pastures from May 1 to Oct. 1 each year at a moderate stocking rate designed to achieve 30 percent forage utilization. Soil type and vegetation communities are similar among replicates, as defined by the Natural Resources Conservation Service’s ecological site descriptions and equivalent land use histories.

Nest Searching

We designated a 4-hectare nest-searching plot in each subpatch (one-eighth of pasture) for a total of 96 plots. We searched each plot four times from May 19 to July 15. We searched for nests via hand-dragging a 30-meter (m) long rope with aluminum can bundles attached every 2.5 m.

Upon flushing a bird, we searched the immediate area for a nest. If the bird displayed a nesting behavior, such as chipping, a broken-wing display or a refusal to leave the immediate area, we marked the location and searched the area again within three days (Hovick et al., 2012). We recorded the coordinates of each nest, and flagged vegetation 5 m north and south of the nest to avoid the association between markings and nests by visual predators (Winter et al., 2003).

We candled two representative eggs from each nest to determine nest age (Lokemoen et al., 1996). We also assessed parasitism rates by brown-headed cowbirds (*Molothrus ater*) because cowbird parasitism may lower nest success in grassland species (Shaffer et al., 2003).

We monitored active nests every two to four days until depredation, completion or abandonment. We considered nests successful if at least one conspecific individual fledged.

Vegetation Monitoring

We standardized the collection date of all nest vegetation data to the actual or expected fledge date of each nest (McConnell et al., 2017). At each nest and at 5 m in each cardinal direction, we assessed the cover of vegetation functional groups using a Daubenmire frame and Daubenmire cover classes, as well as assessed visual obstruction and litter depth (Daubenmire, 1959; Dieni and Jones, 2003).

Statistics

We analyzed nest survival in the RMark interface (Laake, 2013). Daily nest survival was modeled using a logit function in a generalized linear model (Rotella et al., 2004).

For each species, we constructed a continuous model for daily survival, as well as a scale-based hierarchical model detailing the effects of vegetation and management (Dinsmore and Dinsmore, 2007; Hovick et al., 2012; Winter et al., 2003). The first model step evaluates the effects of cowbird parasitism, time-since-fire and incubation stage (laying, incubating or brooding). The second step considers the effects of local (5 m) vegetation. The final modeling step includes nest-site vegetation measurements.

We used nonmetric dimension scaling to evaluate the divergence of avian nesting communities along a time-since-fire gradient using the VEGAN package in R (Oksanen, 2009). We used the

anosim function to test for differences between time-since-fire groupings.

Results

In 2018, we monitored 7,385 nests from 27 species (Table 1).

Nesting	
Nest Total	385
Total Species	27
PBG20	127
PBG40	149
SLG	109
CCSP	72
WEME	53
BWTE	42
NOPI	30
GRSP	23
BRBL	37
MALL	21
CCLO	20
Table 1. Nesting species monitored and number of nests by species and treatment for individuals monitored at the Central Grasslands Research Extension Center, May-August 2018.	

Daily Survival Rate

We were able to run nest survival metrics on every species with 20 or more nests per year (six species total; Table 2).

Blue-winged teal (*Anas discors*) had a constant daily survival rate of 0.96. This corresponds to a total survival rate of 0.38. Greater cover of woody vegetation at the nest site decreased overall survival.

Northern pintails (*Anas acuta*) also had a constant daily survival rate of 0.96, corresponding with a total survival rate of 0.39. Shrub cover enhanced nesting success at the microsite-scale, and was decreased by bare ground cover at the nest site.

Clay-colored sparrows (*Spizella pallida*) had a daily nest survival rate of 0.94, corresponding with a total survival rate of 0.29. Their nest success was decreased by brown-headed cowbird parasitism and positively correlated with visual obstruction at the nest site.

Western meadowlark (*Sturnella neglecta*) daily nest survival was 0.95, with a total survival rate of 0.20. Western meadowlark survival was higher in the nestling stage, as well as in areas with a greater cover of smooth brome at the nest site and bluegrass at the microsite level. Nesting success decreased with increasing visual obstruction.

Brewer’s blackbird (*Euphagus cyanocephalus*) daily survival probability was 0.95, corresponding to a total survival

rate of 0.20. Their survival decreased during the course of the nesting season, and with brown-headed cowbird parasitism and nest-site cool-season grass cover. Nest survival increased with greater vegetation height.

Species (N ≥ 20)	Daily Survival Probability	Model Coefficients
Blue-winged teal	0.96	Nest Shrub -
Northern pintail	0.96	5m shrub+, Nest Bare-
Clay-colored sparrow	0.94	BHCO Parasitism-, nest visual obstruction+
Grasshopper sparrow	0.94	Stage+, nest vegetation height-
Western meadowlark	0.95	Stage+, 5m C3 invasive grasses+, 5m bluegrass+, nest visual obstruction-
Brewer's blackbird	0.95	Time2-, BHCO parasitism-, 5m vegetation height+, nest C3 grass-

Table 2. Daily nest survival rates and final hierarchical model coefficients and directionality for grassland bird species at the Central Grasslands Research Extension Center near Streeter, N.D., in 2018. BHCO = brown-headed cowbird; C3 = cool-season.

Discussion

After two years of data collection, early results highlight the differences in preferred vegetation structure among grassland species. We discovered that new burns create habitat for blackbirds and is reflected in blackbird density and blackbird selection for unburned areas for nesting.

In upcoming years, additional times-since-fire will allow for bird species to exhibit selection for vegetation characteristics at an experimental patch level. We will test to see if patch contrast creates more niches for nesting and breeding birds, and enhances abundance and diversity of birds, compared with traditional range management.

Literature Cited

- Archer, S.R., Andersen, E.M., Predick, K.I., Schwinning, S., Steidl, R.J., Woods, S.R. 2017. Woody plant encroachment, causes and consequences. In: Briske D. (eds) *Rangeland Systems*. Springer Series on Environmental Management. Springer, Cham.
- Bakker, K.K., 2003. The effect of woody vegetation on grassland nesting birds: An annotated bibliography. *Proceedings of the South Dakota Academy of Science* 82, 119-141.
- Bleho, B.I. Koper, N., Machtans, C.S. 2014. Direct effects of cattle on grassland birds in Canada. *Conservation Biology* 28, 724-734.
- Briske, D.D., Sayre, N.F., Huntsinger, L., Fernandez-Gimenez, M., Budd, B., Derner, J.D. 2011. Origin, persistence, and resolution of the rotational grazing debate: Integrating human dimensions into rangeland research. *Rangeland Ecology and Management* 64, 325-334.
- Churchwell, R.T., Davis, C.A., Fuhlendorf, S.D., Engle, D.M. 2008. Effects of patch-burn management on dickcissel nest success in a tallgrass prairie. *Journal of Wildlife Management* 72, 1596-1604.
- Conkling, T.J., Belant, J.L., DeVault, T.L., Martin, J.A. 2017. Effects of crop type and harvest on nest survival and productivity in semi-natural grasslands. *Agriculture, ecosystems, and environment* 240, 224-232.
- Coppedge, B.R., Engle, D.M., Masters, R.E., Gregory, M.S. 2001. Avian response to landscape change in fragmented southern great plains grasslands. *Ecological Applications* 11, 47-59.
- Daubenmire, R.F. 1959. A canopy coverage method of vegetational analysis. *Northwest Science* 33, 43-64.
- Davis, C.A., Churchwell, R.T., Fuhlendorf, S.D., Engle, D.M., Hovick, T.J. 2016. Effect of pyric herbivory on source-sink dynamics in grassland birds. *Journal of Applied Ecology* 53, 1004-1012.
- Dieni, J.S., Jones, S.L. 2003. Grassland songbird nest site selection patterns in northcentral Montana. *Wilson Ornithological Society* 115, 388-396.
- Dinsmore, S.J., Dinsmore, J.J. 2007. Modeling avian nest survival in program MARK. *Studies in Avian Biology* 34, 73-83.
- Doxon, E.D., Davis, C.A., Fuhlendorf, S.D., Winter, S.L. 2011. Aboveground macroinvertebrate diversity and abundance in sand sagebrush prairie managed with the use of pyric herbivory. *Rangeland Ecology and Management* 64, 394-403.
- Fuhlendorf, S.D., Engle, D.M. 2001. Restoring heterogeneity on rangelands: Ecosystem management based on evolutionary grazing patterns. *BioScience*, 51(8), 625-632.
- Fuhlendorf, S.D., Engle, D.M. 2004. Application of the fire-grazing interaction to restore a shifting mosaic on tallgrass prairie. *Journal of Applied Ecology*, 41(4), 604-614.
- Fuhlendorf, S.D., Engle, D.M., Kerby, J., Hamilton, R. 2009. Pyric herbivory: rewilding landscapes through the recoupling of fire and grazing. *Conservation Biology* 23, 588-598.
- Fuhlendorf, S.D., Harrell, W.C., Engle, D.M., Hamilton, R.G., Davis, C.A., Leslie, D.M. Jr. 2006. Should heterogeneity be the basis for conservation? Grassland bird response to fire and grazing. *Ecological Applications* 41, 604-614.
- Fuhlendorf, S.D., Townsend, D.E. Jr., Elmore, R.D., Engle, D.M. 2010. Pyric-herbivory to promote rangeland heterogeneity: Evidence from small mammal communities. *Rangeland Ecology and Management* 63, 670-678.
- Graul, W.D. 1975. Breeding biology of the mountain plover. *The Wilson Bulletin* 87, 6-31.
- Hovick, T.J., Elmore, R.D., Fuhlendorf, S.D. 2014. Structural heterogeneity increases diversity of non-breeding grassland birds. *Ecosphere* 5, 1-13.
- Hovick, T.J., Elmore, R.D., Fuhlendorf, S.D., Engle, D.M., Hamilton, R.G. 2015. Spatial heterogeneity increases diversity and stability in grassland bird communities. *Ecological Applications* 25, 662-672.

- Hovick, T.J., Miller, J.R., Dinsmore, S.J., Engle, D.M., Debinski, D.M., Fuhlendorf, S.D. 2012. Effects of fire and grazing on grasshopper sparrow nest survival. *Journal of Wildlife Management* 76, 19-27.
- Klug, P.E., Jackrel, S.L., With, K.A. 2010. Linking snake habitat use to nest predation risk in grassland birds: The dangers of shrub cover. *Oecologia* 162, 803-813.
- Laake, J.L. 2013. RMark: an R Interface for analysis of capture-recapture data with MARK. AFSC Processed Rep 2013-01, 25 p. Alaska Fisheries Science Center, NOAA, National Marine Fisheries Service, Seattle, Washington..
- Lokemoen, J.T., Koford, R.R. 1996. Using candlers to determine egg incubation stage of passerine eggs. *Journal of Field Ornithology* 67, 660-668.
- Madden, E.M., Murphy, R.K., Hansen, A.J., Murray, L. 2000. Models for guiding management of prairie bird habitat in northwestern North Dakota. *American Midland Naturalist* 144, 377-392.
- McConnell, M.D., Monroe, A.P., Wes Burger Jr., L., Martin, J.A. 2017. Timing of nest vegetation measurement may obscure adaptive significance of nest-site characteristics: A simulation study. *Ecology and Evolution* 7, 1259-1270.
- Oksanen, J. 2009. Multivariate analysis of ecological communities in R: VEGAN Tutorial. <http://cran.r-project.org>.
- Patton, B.D., Dong, X., Nyren, P.E., Nyren, A. 2007. Effects of grazing intensity, precipitation, and temperature on forage production. *Range Ecology and Management* 60:656-665.
- Rogers, W.M., Kirby, D.R., Nyren, P.E., Patton, B.D., Dekeyser, E.S. 2005. Grazing intensity effects on northern plains mixed-grass prairie. *Prairie Naturalist* 37:73-83.
- Rotella, J.J., Dinsmore, S.J., Shaffer, T.L. 2004. Modeling nest-survival data: a comparison of recently developed methods that can be implemented in MARK and SAS. *Animal Biodiversity and Conservation* 27, 187-205.
- Scasta, J.D., Thacker, E.T., Hovick, T.J., Engle, D.M., Allred, B.W., Fuhlendorf, S.D., Weir, J.R. 2016. Patch-burn grazing (PBG) as a livestock management alternative for fire-prone ecosystems of North America. *Renewable Agriculture and Food Ecosystems* 31,550-567.
- Shaffer, J.A., Goldade, C.M., Dinkins, M.F., Johnson, D.H., Igl, L.D. 2003. Brown-headed cowbirds in grasslands: Their habitats, hosts, and response to management. *Prairie Naturalist* 35: 145-186.
- Sirami, C., Seymour, C., Midgley, G. Barnard, P., The impact of shrub encroachment on savanna bird diversity from local to regional scale. *Diversity and Distributions* 2009, 948-957.



C.A. Duquette, NDSU

- Van Hengstum, T., Hooftman, D.A., Oostermeijer, J.G.B., van Tienderen, P.H. 2013. Impact of plant invasions on local arthropod communities: a meta-analysis. *Journal of Ecology* 102, 4-11.
- Vermeire, L.T., Mitchell, R.B., Fuhlendorf, S.D., Gillen, R.L. 2003. Patch burning effects on grazing distribution. *Journal of Range Management* 57, 248-252.
- With, K.A. 1994. The hazards of nesting near shrubs for a grassland bird, the McCown's longspur. *The Condor* 96, 1009-1019.
- Winter, W., Hawks, S.E., Shaffer, J.A., Johnson, D.H. 2003. Guidelines for finding nests of passerine birds in tallgrass prairie. *Prairie Naturalist* 35: 197-211.