

**NDSU Carrington Research Extension Center  
2005 Variety Trial Data**

<b>Oat - Direct Seeded</b>			<b>Tri-County - Wishek (Carrington REC)</b>			
Variety	Days to Head	Plant Lodge	Plant Height	Grain Protein	Test Weight	Grain Yield
		0-9	inch	%	lb/bu	bu/ac
AC Assiniboia	76.8	4.0	37.1	9.9	31.5	73.7
Beach	72.5	3.0	38.4	9.8	32.5	92.8
Buff	68.8	2.3	37.3	12.9	39.1	54.6
Ebeltoft	78.0	3.5	34.9	9.7	29.2	89.5
HiFi	75.8	4.5	39.2	10.2	29.2	73.6
Jerry	72.5	4.0	39.4	10.6	33.2	88.2
Killdeer	71.3	4.0	34.8	9.4	30.9	91.7
Maida	76.8	2.5	39.2	10.8	31.4	86.3
Morton	74.5	3.5	42.9	9.8	29.1	91.7
Paul	77.5	4.0	40.4	11.5	32.2	58.0
Stark	78.0	3.3	39.2	11.6	34.1	37.0
MEAN	74.9	3.5	38.3	10.5	32.0	76.3
C.V.%	1.2	35.9	4.4	4.1	5.4	11.8
LSD.05	1.3	NS	2.4	0.6	2.5	12.9

**Planting Date = April 20 ; Harvest Date = August 2 ; Previous Crop = Spring Wheat**

\* This trial was direct seeded into a non-tilled spring wheat residue.