

**NDSU Carrington Research Extension Center
2008 Variety Trial Data**

Oat - Direct Seeded						Tri-County - Wishek	
Variety	Days to Head	Plant Lodge 0-9	Plant Height inch	Grain Protein %	Test Weight lb/bu	----- Grain Yield -----	
						2008	3-yr. Avg. bu/ac
AC Pinnacle	65.0	1.8	29.7	13.4	36.9	100.1	--
Beach	61.8	1.8	30.5	15.2	39.3	100.2	83.4
HiFi	63.8	2.3	29.5	14.8	37.5	91.8	79.1
Jerry	60.0	4.0	27.4	15.5	39.1	79.1	75.3
Killdeer	61.0	2.0	26.0	14.4	37.7	97.4	73.7
Maida	62.3	4.0	29.1	15.9	37.2	89.8	76.8
Morton	63.3	1.8	30.1	15.1	38.3	90.0	78.0
Paul	65.5	1.8	27.8	20.2	45.5	57.5	57.5
Souris	62.5	2.5	25.4	14.0	37.5	84.6	74.0
Stallion	63.0	2.8	28.5	16.1	39.4	96.9	--
Stark	66.3	2.0	27.8	17.7	42.2	51.7	50.8
Youngs	62.8	2.5	28.7	15.6	37.0	90.2	--
MEAN	63.1	2.3	28.3	15.6	39.0	85.9	--
C.V.%	1.2	50.9	7.8	3.5	1.6	15.4	--
LSD.05	1.1	NS	NS	0.8	0.9	19.0	--

Planting Date = April 29 ; Harvest Date = August 6 ; Previous Crop = Annual Forages

This trial was direct seeded into a non-tilled grass forage residue.