

**2021 Durum Varietal Response to Foliar Fungicide at Mohall**

Cooperators: Dean Schoenberg and the Renville/Bottineau Ag Improvement Association

Financial Contributor: North Dakota Wheat Commission

Variety	Test Weight		Grain Protein		Yield	
	Untrt	Trt*	Untrt	Trt*	Untrt	Trt*
	lbs/bu		%		bu/A	
ND Stanley	59.8	59.2	15.1	15.7	38.6	32.4
AAC Stronghold	59.1	58.7	16.0	16.9	30.6	25.2
AAC Spitfire	57.5	57.1	16.4	16.8	32.7	28.5
ND Riveland	58.7	58.5	15.4	16.1	37.0	27.7
Joppa	59.3	59.2	14.4	15.3	37.5	28.8
ND Grano	59.5	59.2	15.1	15.8	37.1	27.7
Divide	58.6	58.2	15.3	16.3	36.5	26.8
Grenora	58.4	58.7	15.3	16.0	35.5	26.4
Lebsock	59.5	58.9	15.3	16.0	31.5	24.4
Alkabo	59.4	59.4	14.5	14.7	35.8	23.2
Carpio	57.9	57.7	15.2	16.0	38.1	27.7
Mountrail	58.5	58.5	14.4	15.2	32.4	26.7
Tioga	59.2	58.9	16.1	16.7	35.7	28.8
VT Peak	60.0	59.4	15.4	16.4	34.1	29.0
Trial Mean	59.0	58.7	15.3	16.0	35.2	27.4
C.V.%		0.9		3.5		16.2
LSD 0.05		0.9		0.9		8.3

\*8 oz/A Prosaro applied at flowering.

*Planted on May 6 with a seeding rate of 1.5 million PLS/A and harvested on August 16.**Previous Crop: canola**Tillage: minimum till**Soil Type: Barnes loam*

Note: The trial sustained moderate drought. View data with caution.