

WINTER WHEAT VARIETY TRIALS

Table 6 - Off-station winter wheat variety trials			
Variety	Yield in bushels per acre		
	Beach	Hettinger	Avg. 2-station
Froid	35	19	27
Hume	34	19	27
Winoka	44	21	33
Trapper	39	23	31
Lancer	41	21	31
Centurk	44	22	33
Bronze	42	20	31
Standard error of a treatment mean	2.1	1.0	
L.s.d. @ 5%	6.6	2.9	4.8
The c.v. = P.C.	9.3	9.5	

Table 7 - Off station winter wheat variety trial			
Variety	Test weight per bushel		
	Beach	Hettinger	Avg. 2-station
Froid	60.5	56.5	58.5
Hume	59.5	58.0	58.8
Winoka	60.5	58.5	59.5
Trapper	60.0	60.5	60.3
Lancer	58.5	57.0	57.8
Centurk	60.0	58.5	59.3
Bronze	61.5	59.0	60.3

Table 8 - Off station winter wheat variety trials			
Variety	Protein percent		
	Beach	Hettinger	Avg. 2-station
Froid	12.2	13.5	12.9
Hume	11.3	14.3	12.8
Winoka	12.1	13.5	12.8

Trapper	11.0	12.7	11.9
Lancer	11.1	13.1	12.1
Centurk	11.5	12.9	12.2
Bronze	11.0	12.7	11.9
Wheat protein at 14.0% moisture.			

[Back to 1975 Research Reports Table of Contents](#)

[Back to Research Reports](#)

[Back to Dickinson Research Extension Center \(http://www.ag.ndsu.nodak.edu/dickinso/\)](http://www.ag.ndsu.nodak.edu/dickinso/)

[Email: drec@ndsuent.nodak.edu](mailto:drec@ndsuent.nodak.edu)
