

Table 1. 1990 Dickinson Hard Red Spring Wheat Variety Trial

Variety	Bu/A Avg.	Test Wt lbs.	Heading date	Height in.
Amidon	47.9	62.0	26-Jun	35
Bergen*	39.1	60.5	26-Jun	28
Butte 86	40.2	62.5	23-Jun	32
Cutless*	32.7	62.0	27-Jun	32.5
Fjeld*	45.4	62.0	27-Jun	31.5
Grandin*	49.5	61.5	26-Jun	32.5
Gus*	44.3	60.0	27-Jun	31.5
Laura	32.2	61.5	26-Jun	32
Len*	43.7	62.0	29-Jun	31.5
Marshall*	41.0	59.5	29-Jun	30
Minnpro*	40.4	60.5	27-Jun	31
N 86-0370*	46.0	62.0	29-Jun	31.5
2369*	57.5	62.5	28-Jun	31.5
2370*	44.3	61.5	26-Jun	32
2375*	52.5	60.0	27-Jun	34
2385*	43.7	60.0	25-Jun	32.5
Stoa	45.9	62.0	28-Jun	34.5

Vance*	49.5	61.5	30-Jun	30.5
MT 8402*	43.7	59.5	27-Jun	31
ND 655*	45.4	63.5	27-Jun	34
ND 656	42.9	61.5	26-Jun	35
ND 657*	48.1	62.5	27-Jun	34
ND 659*	50.9	59.5	27-Jun	33.5
ND 660	48.1	62.0	27-Jun	34
ND 661	41.8	63.0	25-Jun	34
ND 662*	58.0	62.0	27-Jun	33
ND 663	37.7	63.0	25-Jun	32
ND 664*	49.0	59.5	25-Jun	30.5
ND 665*	45.4	61.5	26-Jun	28.5
SD 2980	46.5	62.0	24-Jun	28.5
W 2501*	53.9	60.0	25-Jun	26
W 2502*	48.7	58.5	25-Jun	28.5
XW 371*	44.8	59.5	25-Jun	28.5
Golden 86*	41.0	63.0	29-Jun	24.5
Stoa+alga Min	39.9	60.0	28-Jun	28

*Semidwarf

Seeding Rate: 1,000,000 live seeds per acre (approx 1 Bu/A); Seeding Date: April 23; Harvest Date: August 3; Fertilizer Applied: 50 lbs/A 18-46-

0 drill application; Herbicide Applied: Hoelon/Buctril tank mix; L.S.D. 5% =
6.72 Bu/A; C.V. % = 9.00