

**NDSU Carrington Research Extension Center  
2020 Variety Trial Data**

<b>Hemp</b>								<b>Carrington</b>
Treatment	Emergence % 2 weeks after planting	Emergence % 3 weeks after planting	Plant Height inch	Days to Flower	Oil %	Test Weight lb/bu	KWT g/1000	Yield lb/ac
CFX-2	53.9	75.8	53.3	37.8	32.8	40.4	16.75	1174
CRS-1	56.1	68.7	57.1	38.0	32.6	40.0	17.01	1358
Katani	47.7	75.6	42.6	37.0	32.6	40.5	14.90	1255
Grandi	39.1	66.6	40.6	37.0	32.4	40.5	15.97	1128
Canda	58.9	67.2	61.1	40.5	30.4	39.4	18.60	1113
Joey	54.6	68.2	58.5	39.5	30.4	39.7	17.95	1265
X-59	26.6	43.2	53.3	38.0	29.2	40.0	16.73	1382
Altair	39.2	65.6	65.2	53.8	30.1	37.1	15.97	1026
Vega	45.1	67.6	62.7	52.0	30.6	38.7	17.08	1208
Henola	39.6	64.1	56.0	53.5	29.7	39.6	13.79	1720
Bilobrzeskie	53.1	94.6	65.7	63.5	31.6	38.4	12.88	963
CFX-1	40.8	58.7	55.1	37.8	32.8	40.3	17.09	1146
Mean	44.9	66.0	58.8	45.3	31.3	39.2	15.64	1155
C.V. (%)	35.3	20.4	16.8	1.9	2.2	1.2	4.3	13.3
LSD 0.05	22.7	19.2	14.2	1.3	1.0	0.7	0.97	220
LSD 0.10	18.9	16.0	11.8	1.1	0.8	0.6	0.81	183

**Planting Date = May 25; Harvest Date = September 8; Previous Crop = Spring Wheat**