North Dakota hard red spring wheat variety descriptions, agronomic traits, 2010.

| | • | | | | | Reaction to Disease ⁴ | | | |
|----------------------------|---------------------------------|------------------|--------|--------------------------------|------------------------------|----------------------------------|--------------|---------------------------|-------------------|
| Variety | Agent or Origin ¹ | Year Released | Height | Straw Strength ² | Days to Head ³ | Stem Rust | Leaf Rust | Leaf Spot ⁵ | Head (Scab) |
| Agawam ⁶ | WestBred | 2008 | 30 | 7 | 58 | NA | S | NA | MS ¹⁰ |
| Albany | Trigen Seed | 2008 | 32 | 5 | 62 | NA | NA | NA | \mathbf{M}^{10} |
| Alpine ⁷ | AgriPro | 2008 | 34 | 6 | 62 | NA | NA | MS | MS |
| Alsen | ND | 2000 | 34 | 3 | 63 | R | MR/MS | S | MR |
| Barlow | ND | 2009 | 35 | 6 | 62 | R | R | MR | M |
| Blade | WestBred/Sabre | 2007 | 33 | 4 | 64 | R | MR | MS | \mathbf{M}^{10} |
| Breaker | WestBred | 2007 | 34 | 3 | 64 | R | MR | MS | M^{10} |
| Brennan | AgriPro | 2009 | 30 | 4 | 62 | R | MR | M | MS |
| Brick | SD | 2009 | 35 | 5 | 60 | R | R | NA | MR |
| Briggs | SD | 2002 | 35 | 7 | 61 | R/MR | R | MS | S |
| Brogan | WestBred | 2009 | 32 | 3 | 64 | MR | MR | MS | S |
| Carberry | Can. | 2009 | 32 | 4 | 57 | NA | NA | NA | NA |
| Choteau ⁶ | MT | 2004 | 32 | 7 | 66 | NA | NA | NA | NA |
| Cromwell | Thunder Seed | 2007 | 33 | 5 | 67 | NA | MR | MR | S |
| Dapps | ND | 2003 | 39 | 5 | 59 | R | R | M | S |
| Edge | WestBred/Sabre | 2008 | 33 | 4 | 62 | NA | NA | NA | MS^{10} |
| Faller | ND | 2007 | 35 | 5 | 65 | R | R | MR | M |
| Freyr | AgriPro | 2004 | 34 | 6 | 64 | R | MR/MS | MS | MR |
| Glenn ⁹ | ND | 2005 | 37 | 4 | 61 | R | R | M | MR |
| Granite ⁹ | WestBred | 2002 | 29 | 2 | 66 | R/MR | MR | S | MS |
| Hat Trick ⁹ | Trigen Seed | 2007 | 34 | 5 | 61 | R | MR | M | MS |
| Howard | ND | 2006 | 36 | 7 | 63 | R | R | M | M |
| Jenna | AgriPro | 2009 | 32 | 4 | 66 | R | MR | M | M |
| Kelby ⁹ | AgriPro | 2006 | 30 | 4 | 62 | MR | R | M | M |
| Knudson | AgriPro | 2001 | 32 | 5 | 64 | MR | MR | MR | M |
| Kuntz ⁹ | AgriPro | 2007 | 31 | 4 | 65 | R | MR | MS | \mathbf{M}^{10} |
| Mott ⁶ | ND | 2009 | 36 | 3 | 66 | MR | MS | MS | MS |
| Muchmore | Can. | 2009 | 32 | 4 | 57 | NA | NA | NA | NA |
| ND 901CL PLUS ⁸ | ND | 2010 | 36 | 4 | 60 | R/MR | MR/R | NA | \mathbf{M}^{10} |
| Pivot | WestBred | 2010 | 27 | 3 | 67 | MS | MR | MR | \mathbf{S} |
| RB07 ⁹ | MN | 2007 | 32 | 6 | 62 | R | R | MS | MR |
| Reeder | ND | 1999 | 35 | 3 | 63 | R | MS | S | S |
| Sabin | MN | 2009 | 33 | 6 | 65 | R | MR | NA | M |
| Samson | WestBred | 2007 | 31 | 2 | 63 | R | MR/MS | MR/MS | S |
| Select ⁹ | SD | 2010 | 35 | 6 | 60 | R/MR | R/MR | R/MR | MR |
| Steele-ND | ND | 2004 | 35 | 7 | 63 | R | R | MS | M |
| SY605 CL ⁸ | AgriPro | 2009 | 34 | 7 | 62 | R/MR | S | MS | S |
| Tom | MN | 2008 | 34 | 6 | 64 | R | R | NA | M |
| Traverse | SD | 2006 | 37 | 6 | 60 | R | MR | NA | M |
| Vantage | WestBred | 2007 | 32 | 2 | 67 | R | MR/MS | MS | MS ¹⁰ |
| WB Digger | WestBred | 2009 | 34 | 6 | 63 | MR | MR/MS | NA | MS |

¹Refers to agent or developer: MN = University of Minnesota; MT = Montana State University; ND = North Dakota State University; SD = South Dakota State University. **Bold** varieties are those recently released so data is limited and rating values may change. NA indicates insufficient information is available to make an accurate assessment.

²Straw Strength = 1-9 scale with 1 the strongest and 9 the weakest. These values are based on recent data and may change as more data become available.

³Days to Head = the number of days from planting to head emergence from the boot averaged over several locations in 2010.

 $^{^4}$ R = resistant; MR = moderately resistant; M = intermediate; MS = moderately susceptible; S = susceptible; VS = very susceptible.

⁵Leaf spot refers to the leaf fungal diseases such as Tan Spot and Septoria. It does not include Bacterial leaf streak.

⁶Solid stemmed or semisolid stem, imparting resistance to sawfly.

⁷Hard white wheat.

⁸CL = refers to a Clearfield variety, with tolerance to Beyond™ family of herbicide.

⁹Indicates varieties that showed Bacterial Leaf Streak symptoms based on very limited data.

¹⁰Based on one year's data.