## Hard Red Spring Wheat Variety Descriptions

						Reaction to Disease <sup>4</sup>			
	Agent or	Year	Height	Straw	Days to	Stem	Leaf	Leaf	Head
Variety	Origin <sup>1</sup>	Rlsd	(in)	Strgth <sup>2</sup>	Head <sup>3</sup>	Rust	Rust	Spot⁵	Scab
Advance	SD	2012	32	6	64	MR	MR	NA	MS
Agawam <sup>6</sup>	WestBred	2008	30	7	58	NA	MR/MS	NA	MS
Alpine <sup>7</sup>	AgriPro	2008	34	6	62	NA	S	MS	MS
Alsen	ND	2000	34	3	63	R	MR/MS	S	MR
Barlow	ND	2009	35	6	62	R	MR/MS <sup>9</sup>	MR	М
Blade	WestBred/Sabre	2007	33	4	64	R	MS <sup>9</sup>	MS	М
Breaker	WestBred	2007	34	3	64	R	MR/MS <sup>9</sup>	MS	М
Brennan	AgriPro	2009	30	4	62	R	MR	М	MS
Brick	SD	2009	35	5	60	R/MR	MR/MS	MS/S	MR
Briggs	SD	2002	35	7	61	R/MR	MR/MS	MS	S
Brogan	WestBred	2009	32	3	64	MR	MR/MS	MS	S
Choteau <sup>6</sup>	MT	2004	32	7	66	NA	MR/MS	NA	NA
Edge	WestBred/Sabre	2008	33	5	62	NA	S	MS	MS
Elgin	ND	2012	36	5	65	R	M <sup>9</sup>	NA	М
Faller	ND	2007	35	5	65	R	S <sup>9</sup>	MR	М
Forefront	SD	2012	37	5	61	MR	MR	NA	MR
Freyr	AgriPro	2004	34	6	64	R	S	MS	MR
Glenn	ND .	2005	37	4	61	R	MR/MS <sup>9</sup>	М	MR
LDS Hat Trick	Limagrain	2007	34	5	61	R	MR/MS	М	MS
Howard	ND	2006	36	7	63	R	MS <sup>9</sup>	М	М
Jenna	AgriPro	2009	32	4	66	R	MR/MS	М	М
Kelby	AgriPro	2006	30	4	62	MR	MR/MS <sup>9</sup>	М	М
Kuntz	AgriPro	2007	31	4	65	R	MS <sup>9</sup>	MS	М
LCS Albany	Limagrain	2008	32	5	67	MR	S	MS	М
LCS Breakaway	Limagrain	2011	32	5	63 05	NA	R	MS	M
LCS Powerplay	Limagrain	2011	33	5	65	NA	MR S <sup>9</sup>	MS	M
Mott <sup>6</sup>	ND	2009	36	3	66	MR		MS	MS
ND 901CL PLUS <sup>8</sup>	ND	2010	36	4	60	R/MR	MR	NA	М
Norden	MN	2012	32	3	6	R	R/MR S <sup>9</sup>	M	M
Pivot	WestBred	2010	27	3	67	MS		MR	S
Prosper	NDSU	2011	35	5	65	R	S <sup>9</sup>	М	М
RB07	MN	2007	32	5	62	R	R/MR <sup>9</sup>	MS	MR
Rollag	MN	2011	32	3	63	R	MS	MR	MR
Sabin	MN	2009	33	6	65	R	MR	MS	M
Samson	WestBred SD	2007	31	2	63	R	MR/MS	MS	S
Select SY Soren	-	2010	35 30	6 4	60 62	R/MR	R/MR MR	R/MR M	MR M
SY Tyra <sup>6</sup>	Sygenta/AgriPro Sygenta/AgriPro	2011		4 5	63 62	R R	MR	NA	S
SY 191a <sup>®</sup> SY605 CL <sup>8</sup>		2011	31 34	5 7	62 62	R/MR		MS	S
	AgriPro	2009			-		MR/MS	-	
Tom	MN WestBred	2008 2007	34 32	6 2	64 67	R MR	MR/MS	NA MS	M MS
Vantage				4			MR/MS		
Velva	NDSU WestBred	2011	35 34	4	63 63	R MR	MR/MS <sup>9</sup>	M NA	MS MS
WB Digger		2009 2011	34 30	6 4	63 63	R	MR/MS MR/MS	MA MS	INIS S
WB Mayville	Monsanto/WB	2011	30	4	03	ľ,	IVIT/IVIO	IVIO	3

<sup>1</sup> Refers to agent or developer: MN = University of Minnesota, MT = Montana State University, ND = North Dakota State University, SD = South Dakota State University. **Bold** varieties are those recently released so data is limited and rating values may change.

<sup>2</sup> Straw Strength = 1 to 9 scale, with 1 the strongest and 9 the weakest. These values are based on recent data and may change as more data become available.

 $^{3}$  Days to Head = the number of days from planting to head emergence from the boot averaged over several locations in 2010 and 2011.

 $^{4}$  R = resistant, MR = moderately resistant, M = intermediate, MS = moderately susceptible, S = susceptible. NA indicates insufficient information is available to make an accurate assessment.

<sup>5</sup> Leaf spot refers to the leaf fungal diseases such as tan spot and septoria. It does not include bacterial leaf streak.

<sup>6</sup> Solid stemmed or semisolid stem, imparting resistance to sawfly. <sup>7</sup> Hard white wheat.

<sup>8</sup> CL = refers to a Clearfield variety, with tolerance to Beyond <sup>TM</sup> family of herbicides.

<sup>9</sup> These lines were resistant to moderately resistant to races prevalent prior to 2011, and show some level of susceptibility to a new race of the pathogen that was detected at low frequency in 2012.

Note: Published in NDSU publication A-574 (revised).