2020 Public Lentil Variety Trial at Minot

							1000				
	Market	Days to	Days to	Vine	Pod	Seed	Seed	Seeds/	Test	Seed	Significant
Variety	Class	Flower	Maturity	Length	Height	Protein	Weight	Pound	Weight	Yield	Difference ⁴
		DAP ¹	DAP ¹	cm	cm	% ²	g	#	lb/bu	bu/A ³	
CDC Richlea	MG	45	73	32	19	23.9	47	10199	62.4	18.4	Α
CDC Greenland	LG	45	76	39	23	24.5	51	9494	60.1	18.2	Α
CDC Imvincible-C	SG	46	73	34	19	27.2	39	11830	63.9	18.1	Α
CDC Kermit	SG	46	73	35	21	25.1	37	12619	64.7	18.1	Α
CDD Impala-CL	ESR	46	74	31	16	25.4	45	11459	65.0	18.1	Α
CDC Greenstar	LG	45	73	36	25	23.3	61	7908	60.3	17.9	Α
CDC Maxim-CL	SR	45	73	36	21	23.7	44	10482	64.5	17.9	Α
ND Eagle	SG	46	74	26	13	25.6	45	10432	64.1	17.9	Α
Avondale	MG	44	71	36	19	23.7	49	9547	62.3	17.8	Α
Trial Mean		45	73	34	19	24.7	47	10441	63.0	18.0	_
C.V. %		1.3	1.7	11.3	17.8	5.07	15.2	13.5	2.94	0.99	
LSD 5%		1.2	8.0	2.6	2.6	0.45	16.8	3504.6	0.60	NS	

¹ Days after planting

<u>Note</u>: No significant differences were observed for lodging. All plots maintained upright habit throughout the season.

2020 Public Chickpea Variety Trial at Minot

•	-			1000				
	Days to	Days to	Vine	Seed	Seeds/	Test	Seed	Significant
Variety	Flower	Maturity	Length	Weight	Pound	Weight	Yield	Difference ³
	DAP ¹	DAP ¹	cm	g	#	lb/bu	bu/A ²	
CDC Frontier	44	99	45	334	1306	63.3	26.9	Α
ND Crown	44	98	44	374	1148	62.7	23.7	Α
CDC Palmer	44	94	44	395	1222	62.6	21.3	Α
CDC Leader	44	99	43	364	1270	61.8	15.9	В
CDC Orion	42	105	43	342	1311	60.6	9.1	C*
Sawyer	43	104	43	432	1078	60.8	8.6	C*
Royal	46	101	44	478	956	61.2	8.1	C*
Sierra	43	103	43	422	1119	61.3	6.9	C*
Trial Mean	44	101	44	393	1176	61.8	15.0	
C.V. %	2.8	3.5	1.5	12.5	10.6	1.59	53.18	
LSD 5%	2.2	3.2	6.8	60.3	178.7	1.53	7.84	

^{*}Yields were severely impacted by Ascochyta Blight

Planting date: 5/11/2020

Planting date: 5/11/2020

Harvest date: 8/17/2020

Tillage: Minimal Tillage

Previous Crop: Flax

Harvest date: 9/4/2020 Previous Crop: Flax

Tillage: Minimal Tillage

<u>Note</u>: No significant differences were observed for lodging or seed size. All varieties met the RMA requirement for large kabuli types that at least 51% stayed above a 22/64" round hole screen.

² Protein adjusted to 0% moisture

³ Yield adjusted to 13.5% moisture

⁴ Seed yields within a market class with different letters are significantly (p<0.05) different

¹ Days after planting

² Yield adjusted to 13.5% moisture

³ Seed yields within a market class with different letters are significantly (p<0.05) different