

EFFECTS OF DORMANT SEASON GRAZING ON HERBAGE PRODUCTION AND PLANT GROWTH

Kristy-Layne Nelson¹, Kevin Sedivec¹, Jackie Olson², Mitch Faulkner³, Tim Faller⁴,
Christopher Schauer⁴.

¹Graduate Student, Extension State Rangeland Specialist, Animal and Range Sciences
Department, NDSU, Fargo

²Assistant Professor of Agriculture, Dickinson State University, Dickinson, ND

³Range Specialist, Bureau of Land Management, Lander WY

⁴Director, and Post-Doc Research Associate, Hettinger Research Extension Center, NDSU,
Hettinger, ND

Impact Statement

Preliminary data indicated that brief early summer use of dormant-season pastures and winter stocking rates intended to achieve 50% utilization of standing aboveground biomass is the preferred management grazing option, relative to grazing treatments of 30 or 50% winter utilization with no summer use.

Introduction

Winter, or dormant-season grazing, is practiced by many North and South Dakota livestock producers in an effort to lower winter feed costs. Dormant-season grazing, while not an exclusive winter-grazing period, is defined as grazing during that time period between plant quiescence in late fall and green up in early spring. Although adequate information exists regarding nutritional management of winter grazing cattle, little is known about the ecological effects of these practices on range or pasture land in the upper Midwest and northern Great Plains. Furthermore, research emphasizing inferences for specific winter-grazing management is lacking. Various aspects of dormant-season grazing have been examined in a variety of ecosystem types, and conventional wisdom dictates that defoliation during winter months while plants are dormant has little to no effect on plant vigor (Riesterer et al. 2000).

Vallentine (2001) stated that determining when to harvest standing forage is dependent on plant factors, physical site factors, animal factors, and economic and management factors. Determining the effectiveness and practicality of dormant-season grazing requires assessing management practices to address these four points. Research is abundant regarding the effects of grazing and clipping on rangeland condition and productivity, animal production, and economics during the growing season, yet insufficient information exists regarding the effects of such practices during the dormant season. Furthermore, with the implementation of new grazing management techniques that include season-of-use changes, differences in economic return, plant growth and vigor, and subsequent plant-species composition can be expected.

Winter grazing is an appealing management option to many ranchers. Producing hay or purchasing winter feeds is labor and capital intensive, while winter grazing offers the potential for flexibility in making management decisions. Furthermore, this practice allows for a more efficient utilization of range resources. The objectives of this study were to determine the impacts of winter grazing on herbage production, growth rate of dominant grass species (short-

term), and changes in plant species composition using various levels and combinations of winter and summer use (long-term subsequent research).

Study Area

This study was located in Adams County, North Dakota and Perkins County, South Dakota. The Adams County study site was approximately 62 ha and located 8 km southwest of Hettinger, North Dakota (El. 817m) on sections 16, T129N, R96W and 25, R97W, T129N. The Perkins County study site was approximately 58 ha and located 25 km south of Lodgepole, South Dakota (El. 803m) on sections 13, T19N, R12E, and 18, T19N, R13E.

Climate

Western North and South Dakota are characterized by a continental, cool temperate, semi-arid climate with warm summers and cold winters. Average annual precipitation is about 41 cm/yr near Hettinger, North Dakota and Lodgepole, South Dakota (Owenby and Enzell 1992). Approximately 70% of the precipitation at both locations occurs during the growing season occurring from April through early October. The study locations receive 120-130 frost-free d/yr (Omodt et al. 1968) with the warmest month, July, averaging 22° Celsius. January is the coldest month with an average temperature of -8° Celsius (Owenby and Enzell 1992).

Growing-season precipitation was 28.6 cm in 2000, which was 7.6 cm below the annual average, with all months except May and July below average during the growing season. The 2001 growing season was characterized as a dry spring and wet July, with average precipitation 4.0 cm below the 30-year average. The fall and winter of 2000-01 received above average precipitation; however the fall and winter of 2001-02 received considerably less precipitation, particularly in November and December. In 2002, growing-season precipitation was 13.74 cm, which was 20.42 cm below the annual average. All growing season months were below average. Less-than-average precipitation was received in November and December of 2002. Growing-season precipitation received in 2003 was 6.68 cm below the 30-year average, with only the months of May and September receiving above average precipitation during the growing season. During the 2004 growing season, precipitation was 11.6 cm below the 30-year average. Only the months of July and October were above average precipitation.

Monthly average temperatures were generally above the 30-year average in 2000, with the exception of June, November, and December. Warmer-than-average temperatures characterized the winter of 2001-2002, as November and December 2001 and January and February 2002 were substantially warmer than the 30-year average. The winter of 2002-2003 also had warmer-than-average temperatures compared to the 30-year average. Spring and summer temperatures were near average for 2000-2003. The winter of 2003-2004 was characterized by warmer than average temperatures. The year 2004 fell below the 30-year average in nearly every month except March, September and November. Temperatures in 2005 were similar with numbers either near or below the 30-year average. The only months above average were April and September.

Vegetation

The study areas were found in the northern mixed-grass prairie and described as the Missouri Slope Vegetation Zone (SCS 1984). The plant communities are described as a wheatgrass-

needlegrass vegetation type (Barker and Whitman 1994). Dominant midgrass species were western wheatgrass and needle-and-thread (*Stipa comata* Trin. & Rupr.), and dominant short graminoid species were thread-leaf sedge (*Carex filifolia* Nutt.) and blue grama (*Bouteloua gracilis*; Barker and Whitman 1994; Shiflet 1994). Plant names were referenced from McGregor et al. (1986) and USDA-USFS (2002).

Uplands of the study areas were represented by shallow ecological sites. The soils were moderately coarse to moderately fine textured with restricting layers of shales and sandstone occurring at depths of 25 to 50 cm below the soil surface. This site is comprised of upland sedges (*Carex spp.*), little bluestem, blue grama, prairie sandreed (*Calamovilfa longifolia* (Hook.) Scribn.), western wheatgrass, prairie junegrass, fringed sagewort, and numerous upland forb species. Average total primary production on shallow ecological sites in the Dakotas varies from 1,350 to 2,000 kg/ha (Sedivec et al. 1991).

Midland sites on the North and South Dakota study areas were comprised of a loamy ecological site with well-drained, moderately deep and medium to fine textured soils. Plant communities were comprised of western wheatgrass, needle-and-thread, blue grama, Kentucky bluegrass (*Poa pratensis* L.), prairie junegrass, cheatgrass (*Bromus tectorum* L.), upland sedges (*Carex spp.*), green needlegrass (*Stipa viridula* Trin.) and numerous forb species. Average total primary production on these sites range from 1,700 to 3,150 kg/ha (Sedivec et al. 1991).

Materials and Methods

Treatments: Two study areas (blocks) were selected in North and South Dakota based on similar range condition and native plant species composition. Each study area was blocked and divided into four paddocks with one of four treatments 1) 50% summer-utilization (season-long control; **SL**), 2) 25% summer use for 2 weeks in early and mid June and 50% dormant-season utilization [flash grazing (Hart 2001); **FL**], 3) 30% dormant-season utilization (**DS 30**), and 4) 50% dormant-season utilization (**DS 50**) assigned randomly to a paddock within block. The SL treatment was a 32 ha paddock and the dormant-season-use treatments were each 9 ha paddocks in North Dakota. The South Dakota DS 30 and SL treatment paddocks each 12 ha; the FL treatment was 15 ha; and the DS 50 treatment was 19 ha.

Stocking Rates: Stocking rates for the summer-use treatments were determined using the United States Department of Agriculture (USDA) Natural Resource Conservation Service (NRCS) Technical Guide (2001) for the Missouri Slope Vegetation Zone. Summer use paddocks were surveyed for ecological site composition using the USDA SCS soil surveys for Adams County, North Dakota (Ulmer 1987) and Perkins County, South Dakota. (Wiesner 1980). The stocking rate for the season-long grazing treatment was calculated for a 4-month grazing period beginning June 1 and ending October 1. The North Dakota site was stocked at 0.76 ha/AUM with ten 544 kg cows and their calves. The South Dakota site was stocked at 0.63 ha/AUM with seven 280 kg spayed heifers (Table 3). Summer use of the flash grazing paddocks was targeted for 25% utilization.

The summer-use FL treatment grazing carrying capacities were calculated by stocking for 50% use of the total available AUMs in June while considering that 50% of the total annual production occurred by mid June, thus achieving a 25% utilization of total annual biomass. The

North and South Dakota sites were stocked with ten and sixteen 1200 lb. cows and calves or 1.78 ha/AUM and 1.69 ha/AUM, respectively, for two weeks.

Stocking rates for the winter grazing treatments were calculated after determining dry-standing plant biomass on 15 Nov. 2000. Ten randomly placed 0.25m² frames were clipped for each ecological site (n=2) existing within a given replicate (n=20). The USDA SCS (Wiesner 1980, Ulmer 1987) soil survey maps and technical guides were used to estimate ecological site composition within each paddock to calculate total standing biomass. Final stocking rates for each treatment were computed by calculating 25% grazing-use efficiency with 50% disappearance (Laycock et al. 1972, Pearson 1975) and a dry matter intake for an 1150 lb non-lactating cow (NRC 1996).

The North Dakota DS 50 and FL grazing treatments were each stocked with four 1200 lb. cows, or 1.23 and 1.22 ha/AUM, respectively, and the DS 30 treatment was stocked with three 1200 lb. cows, or 1.64 ha/AUM. The DS 50 treatment was stocked with 11 cows or 1.02 ha/AUM, the FL treatment stocked with 8 cows or 0.95 ha/AUM, and the DS 30 treatment stocked with 6 cows or 1.02 ha/AUM at the South Dakota site. All South Dakota treatments were stocked with cows weighing an average of 1150 lb.

Winter grazing cattle were allowed ad libitum access to white salt and trace minerals and were supplemented with 3 lb/day on an as-fed basis of 30% crude protein all-natural cake. During the winter grazing period of 2000-2001, cattle grazed as snow cover allowed for 53 days beginning November 15 on both the North and South Dakota study sites. During the dormant-season grazing period of 2001-2002, cattle grazed on the North Dakota site for 53 days beginning November 15. The cattle on the South Dakota site grazed for 35 days and animal numbers were increased to meet set stocking rate guidelines, as turn out was delayed until January 12 due to mechanical failures affecting the watering system. Cattle were allowed to graze at both the North Dakota and South Dakota sites for 53 days beginning on November 15 during the 2002-2003 and 2003-2004 winter grazing period. During the 2004-2005 winter grazing period, cattle grazed on the North Dakota sites for 50 days beginning November 22, while the cattle on the South Dakota site grazed for 56 days beginning November 15. Cattle grazed the North Dakota site for 51 days during the 2005-2006 winter grazing season beginning November 15. Cattle grazed for 36 days on the South Dakota site beginning November 29 and were pulled off early due to crusted snow.

Table 1 reports ha/AUM comparisons of treatments and percent reduction in carrying capacities compared to the SL treatment (control). From a perspective of utilized AUMs, the dormant season only grazing treatments had reduced carrying capacities relative to season-long use, however, the FL treatment numerically increased carrying capacities slightly relative to season-long use (3.2 to 5.3%).

Table 1. Stocking rate comparisons among grazing treatments in North and South Dakota.

	SL	FL	DS 30	DS 50
N.D.				
Ha/AUM	0.8	0.7	1.6	1.2
% Difference from SL	0.0	+5.3	-115.8	-61.8
S.D.				
Ha/AUM	0.6	0.6	1.0	0.9
% Difference from SL	0.0	+3.2	-61.9	-38.1

SL = season-long summer grazing, FL = 25% summer use for 2 weeks in early and mid June and 50% dormant season utilization, DS30 = 30% dormant season utilization, DS50 = 50% dormant season utilization

Herbage Disappearance: The degree of disappearance of graminoids and forbs for each treatment was determined using a paired-plot clipping technique (Milner and Hughs 1968). Twenty cages were distributed in each pasture during the treatment period. One plot within and outside each cage was clipped using a 0.25m² quadrat. Clipped herbage was separated into grasses and forbs, dry matter weights recorded, and kg/ha plant biomass and standard error of the mean calculated for each ecological site. Percent utilization and standard errors of the mean were calculated for graminoids and forbs combined and graminoids alone.

In the summer through winter periods of 2000-2001, 5 cages were systematically placed on each of the 2 shallow ecological sites and 2 loamy ecological sites before grazing began on each treatment (n=20), with the exception of the South Dakota 30% treatment where only 5 cages were placed on a shallow ecological site since this site made up only 10% of the study area on the treatment. During the winter of 2001 on the North Dakota sites, 5 plots were clipped for both the loamy and shallow ecological sites on the 30% treatment, 5 shallow plots were clipped on the 50% treatment, and no plots were clipped on the FL treatment due to ice and snow cover. On the summer treatments of 2001, the 20 sites within each pasture selected for the tiller study were used to determine utilization. In 2001-2002, 2002-2003, 2003-2004 and 2004-2005 all plots from the winter grazing treatments were clipped since ice and snow cover did not prevent clipping as it had in 2000-2001.

In May 2004, 30 cages were distributed evenly across the entire pasture, regardless of ecological differences. This created a more representative sampling of the pastures.

Leaf Heights: A study examining leaf heights throughout the growing season was initiated in May of 2001 to determine the growth patterns of western wheatgrass, needle-and-thread, thread-leaf sedge, and blue grama within each treatment. The species were selected as they were described as the predominant forage base of the study region (Barker and Whitman 1994, Shiflet 1994). Furthermore, these species are described as commonly existing together in various successional stages of rangeland in western North Dakota (Hansen and Hoffman 1988). Goetz (1963) monitored the growth and development of native range plants in western North Dakota and used leaf height as a main indicator of plant growth. Furthermore, researchers have

correlated leaf and plant height with plant vigor, forage yield, competition, range condition and trend, and defoliation levels (Short and Woolfolk 1956, Buwai and Trlica 1977).

Twenty locations indicative of the dominant forage base were selected randomly within each treatment in May 2001. On each location, a 0.25 m² quadrat was selected containing at least 10 western wheatgrass tillers, 5 needle-and-thread tillers, 10 thread-leaf sedge tillers, and 10 blue grama tillers. Cool-season tillers were marked with uniquely-colored rings upon the selection of each site in mid May and each tiller measured monthly until senescence was observed for each species. Western wheatgrass and needle-and-thread tillers were measured mid monthly for leaf height (height of tallest leaf) from May to August. Thread-leaf sedge was measured mid monthly for leaf height from May to July. Blue grama was the only warm-season grass investigated for growth, thus leaf heights were measured mid monthly during its growth period as described by Goetz (1964), from June to September. Needle-and-thread and thread-leaf sedge were not measured in 2002 and 2003 to improve random distribution of the plots throughout the four treatments.

Thirty locations were selected in May 2004 within each treatment. On each location, a 0.25 m² quadrat was selected containing 10 western wheatgrass tillers (cool season) and 10 blue grama tillers (warm season). Tillers were marked with uniquely colored rings and measured monthly until senescence. Western wheatgrass was measured from May until August and blue grama was measured from June until September.

Herbage production was analyzed as a randomized complete design with the GLM procedure of SAS (SAS Inst., Inc., Cary, NY) using frame as the experimental unit. The model included treatment, year, and treatment X year using the residual error term. When differences were found ($P < 0.05$), a mean separation was conducted using a Tukey's Honest Significant Difference Test (Steel and Torrie, 1980).

Results and Discussion

Herbage Production: No differences in herbage production were found between locations ($P=0.30$). Following one year of treatment, peak primary production on the winter-only and flash treatments were not different ($P>0.05$) than the SL control treatment (Figure 1). In 2002, peak primary production on the SL control treatment was lower ($P<0.05$) than FL and DS 50. No differences ($P>0.05$) in peak primary production were found on SL, FL, and DS 30 following the third year (2003) of treatment; however peak herbage production was lower ($P<0.05$) on DS 50 compared to the other three treatments (Figure 1). In 2004 as well as in 2005, the SL treatment had lower ($P<0.05$) peak herbage production than the DS 50 and 30 treatments. The FL treatment had lower production ($P<0.05$) than the DS 30% in 2004, and lower production ($P<0.05$) than the DS 50% in 2005.




Figure 1. Peak grass production (mean \pm S.E.) on the season-long (SL), June flash and 50% dormant-season use (FL), 30% dormant-season use (DS 30), and 50% dormant-season use (DS 50) for 2001-2005.

^{ab}Treatments with the same letter are not significantly different ($P>0.05$).

^{fg}Treatments with the same letter are not significantly different ($P>0.05$).

^{mn}Treatments with the same letter are not significantly different ($P>0.05$).

^{stu}Treatments with the same letter are not significantly different ($P>0.05$).

^{xyz}Treatments with the same letter are not significantly different ($P>0.05$).

On average, moderate dormant-season grazing treatments did not effect herbage production, which corresponds with data reported by Coughenour (1991) who found increased nitrogen in live and dead grasses and fringed sagebrush on winter grazed areas. Increases in various aspects of rangeland productivity under moderate growing-season grazing use versus light or no grazing use were reported by Manley et al. (1995). Comparisons of grazed and ungrazed range in Wyoming revealed elevated levels of C and N in subsurface soil on grazed areas. This phenomenon is attributed to the reduction of litter and standing plant biomass as these sources immobilize a significant amount of N and C. Furthermore, increased animal traffic may enhance physical breakdown of residual plant material and soil incorporation. Likewise, Schacht et al. (1998) observed that mowing dormant range of switchgrass, little bluestem, and big bluestem resulted in a higher yield of annual growth than a non-mowed control. Engle et al. (1998) also reported which grazing strategies emphasizing defoliation during the dormant season that decreased probability of multiple defoliations during the growing season were less detrimental than those that increased the probability of multiple defoliations, such as the FL treatment in this study. Relevant research by Auen and Owensby (1988), Coughenour (1991); Engle et al. (1998), Schacht et al. (1998) and Reisterer et al. (2000) indicate dormant-season harvesting of grasses had little or no negative effect on subsequent herbage production.

Leaf Heights: No significant interaction was found for treatment by month, which means that all months react in the same way to the treatments. In 2001, the SL treatment leaf heights were higher than the FL treatment (Figure 2). The FL treatment leaf heights were lower ($P<0.05$) than the DS 50 treatment. Negative effects of grazing treatment on late growing season plant production were also observed by Trent et al. (1988). Fall grazed winter wheat plants relied on photosynthesis later in the growing season than non-grazed wheat plants, as they were unable to draw from carbohydrate reserves during grain filling. Similarly, Buwai and Trlica (1977) found

heavy quiescent defoliation of western wheatgrass reduced TNC relative to a non-defoliated control. Furthermore, moderate and heavy dormant defoliation of western wheatgrass reduced both herbage yield and plant height when compared to the control.




Figure 2. Western wheatgrass leaf heights in July (Peak) (mean \pm S.E.) on the season-long (SL), June flash and 50% dormant-season use (FL), 30% dormant-season use (DS 30), and 50% dormant-season use (DS 50) for 2001-2005.

^{abc}Treatments with the same letter are not significantly different ($P>0.05$).

^{fgh}Treatments with the same letter are not significantly different ($P>0.05$).

^{mn}Treatments with the same letter are not significantly different ($P>0.05$).

stTreatments with the same letter are not significantly different ($P>0.05$).

^{xyz}Treatments with the same letter are not significantly different ($P>0.05$).

In 2002, the FL and DS 30 treatments had lower western wheatgrass heights ($P<0.05$) than the DS 50 treatments. In 2003, no differences ($P<0.05$) in western wheatgrass leaf heights were detected among any treatments (Figure 2).

In 2004, the leaf heights in the FL and SL treatments were shorter ($P<0.05$) than the DS 30 and DS treatment leaf heights (Figure 2). Western wheatgrass leaf heights on the FL and SL treatments were also lower ($P<0.05$) than the DS 30 and DS 50 treatments in 2005. The DS 30 treatment also had lower leaf heights ($P<0.05$) than the DS 50 treatment.

Light winter use (DS 30) resulted in shorter leaf heights than heavy winter use for needle-and-thread, thread-leaf sedge, and blue grama in 2001. Light winter use also resulted in lower needle-and-thread and blue grama leaf heights than SL. These data suggest increased utilization during the dormant period result in increased herbage yield the following year. Treatment by date interactions were not detected ($P<0.05$) for blue grama in 2001, thus monthly leaf height

data were combined. However, in 2002 there was treatment by date interactions ($P < 0.05$) in July. No treatment by date interaction was detected ($P < 0.05$) in 2003.

As with the western wheatgrass leaf heights, no month by treatment interaction was detected. In 2001, the SL and FL treatments were lower ($P < 0.05$) than DS 30 treatment. The DS 30 treatment leaf heights were lower ($P < 0.05$) than the DS 50 treatment (Figure 3). In 2002, the SL treatment leaf heights were lower ($P < 0.05$) than the FL and DS 50 treatments. There were no differences found ($P < 0.05$) in blue grama leaf heights in 2003 (Figure 3). In 2004, the SL treatment blue grama leaf heights were shorter ($P < 0.05$) than the FL, DS 30, and DS 50 treatments. The leaf heights for the FL and DS 30 treatments were lower ($P < 0.05$) than the DS 50 treatment. Blue grama leaf heights for the SL treatment were lower ($P < 0.05$) than the FL and DS 30 treatments in 2005. The FL and DS 30 treatments had lower ($P < 0.05$) leaf heights than the DS 50.




Figure 3. Blue grama leaf heights (mean \pm S.E.) in August (Peak) on the season-long (SL), June flash and 50% dormant-season use (FL), 30% dormant-season use (DS 30), and 50% dormant-season use (DS 50) for 2001-2005.

^{abc}Treatments with the same letter are not significantly different ($P > 0.05$).

^{fg}Treatments with the same letter are not significantly different ($P > 0.05$).

^{mn}Treatments with the same letter are not significantly different ($P > 0.05$).

stTreatments with the same letter are not significantly different ($P > 0.05$).

^{wxyz}Treatments with the same letter are not significantly different ($P > 0.05$).

These findings are consistent with the peak herbage production observations and studies by Coughenour (1991) and Manley et al. (1995) who reported positive effects on herbage production with increased levels of herbage removal during the dormant season. If dormant-season defoliation is not detrimental to needle-and-thread, blue grama, and thread-leaf sedge, removal of standing-dead plant material and the corresponding reduction in litter on the soil surface may be important to subsequent herbage production and plant growth. Removal of standing dead plant material has been reported to elevate soil temperatures, thus accelerating

decomposition and mineralization in the spring. Furthermore, nutrient turnover rates are accelerated under grazed systems by returning mineral nitrogen to the soil in a readily decomposable form, thereby bypassing slower plant litter decomposition pathways (Coughenour 1991).

Summary

During the months of May through November in 2000, seasonal forage availability declined considerably between the time of peak production and the beginning of the winter grazing season. Considering these losses, stockpiling of forage throughout the growing season for use in late fall or winter results in lost herbage production potential. Furthermore, AUMs/ha for winter-only grazing areas were severely reduced relative to season-long grazing use. Incorporating a brief early-summer grazing period on winter pastures can increase land use and reduce economic losses by increasing stocking rates.

From an ecological and land-use efficiency perspective, a dormant-season grazing system that incorporates moderate early summer use combined with winter stocking rates utilizing 50% of the standing plant biomass is a preferable, and moreover, a beneficial management alternative. This method yielded greater herbage production than other treatments and resulted in greater needle-and thread and thread-leaf sedge leaf heights than the season-long or DS 30 grazing treatments. This method reduced western wheatgrass leaf heights late in the growing season in 2001, however, western wheatgrass and blue grama leaf heights were not affected in 2002 and 2003 compared to summer grazing alone. If the conclusion is made that dormant-season defoliation has little effect on these grasses, limiting litter accumulation on stockpiled pastures by ensuring at least moderate utilization (50%) of standing plant biomass may positively affect subsequent herbage production. Furthermore, SL grazing may have a more negative affect on needle-and-thread and thread-leaf sedge growth than winter use at higher (50%) utilization levels. The direct effects of dormant-season grazing on individual plant species versus conventional SL use, at present, is undistinguished in relevant literature. This research indicates western wheatgrass and blue grama were generally unaffected by dormant-season grazing.

Conclusion

Preliminary data regarding dormant-season grazing of native rangeland in the western Dakotas indicated that brief early summer use of dormant-season pastures and winter stocking rates intended to achieve 50% utilization of standing aboveground biomass is the preferred management option, relative to grazing treatments of 30 or 50% winter utilization with no summer use. This method was beneficial from both a land-use and ecological standpoint. Subsequent data are necessary; however, to evaluate the long-term ecological and economic sustainability of this management.

Literature Cited

- Auen, L.M. and C.E. Owensby. 1988.** Effects of dormant-season herbage removal on Flint hills rangeland. *J. Range Manage.* 41:481-482.
- Barker, W.T. and W.C. Whitman. 1994.** Rangeland cover types of the northern Great Plains. *In: Shiflets, T.N., Rangeland cover types of the United States. Soc. for Range Manage., Denver, CO. 154pp.*

- Buwai, M. and M.J. Trlica. 1977.** Multiple defoliation effects on herbage yield, vigor, and total nonstructural carbohydrates of five range species. *J. Range Manage.* 30:164-171.
- Coughenour, M.B. 1991.** Biomass and nitrogen response to grazing of upland steppe on Yellowstone's northern winter range. *J. Appl. Ecol.* 28:71-82.
- Engel, R.K., J.T. Nichols, J.L. Dodd, and J.E. Brummer. 1998.** Root and shoot responses of sand bluestem to defoliation. *J. Range Manage.* 51:42-46.
- Goetz, H. 1963.** Growth and development of native range plants in the mixed grass prairie of western north Dakota. M.S. Thesis. North Dakota State Univ., Fargo. 165pp.
- Hansen, P.L. and G.R. Hoffman. 1988.** The vegetation of the Grand River/Cedar River, Sioux, and Ashland Districts of the Custer National Forest: a habitat type classification. Gen. Tech. Rep. RM-157. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station. 68p.
- Hart, R.H. 2001.** Where the buffalo roamed-or did they? *Great Plains Res.* 11:83-102.
- Laycock, W.A., H. Buchanan, and W.C. Krueger. 1972.** Three methods for determining diet, utilization, and trampling damage on sheep ranges. *J. Range Manage.* 25:352-356.
- Manley, J.T., G.E. Schuman, J.D. Reeder, and R.H. Hart. 1995.** Rangeland soil carbon and nitrogen responses to grazing. *J. Soil and Water Cons.* 50:294-298.
- McGregor, R.L., T.M. Barkley, R.E. Brooks, and E.K. Schofield 1986.** Flora of the Great Plains. University Press of Kansas. Lawrence, K.S..
- Milner, C. and R.E. Hughes. 1968.** Methods of the measurement of primary production of grassland. Blackwell Sci. Pub. Oxford, England.
- National Research Council. 1996.** Nutrient requirements for beef cattle (7th ed.) Nat. Acad. of Sciences-NRC, Washington, D.C.
- Omodt, H.W., G.A. Johnsgard, D.D. Patterson, and O.P. Olson. 1968.** The major soils of North Dakota. Dept. of Soils, Ag. Exp. Sta. North Dakota State Univ., Fargo. Bull. No. 472. 60 pp.
- Owenby, J.R. and D.S. Ezell. 1992.** Monthly station normals of temperature, precipitation, and heating and cooling degree days 1961-1990, North Dakota. Climatology of the United States No. 81. NOAA, National Climatic Data Center. Asheville, NC. 60pp.
- Pearson, H.A. 1975.** Herbage disappearance and grazing capacity determinations of southern pine bluestem ranges. *J. Range Manage.* 28:71-73.
- Riesterer, J.L. M.D. Casler, D.J. Undersander, and D.K. Combs. 2000.** Seasonal yield distribution of cool-season grasses following winter defoliation. *Agron. J.* 92:974-980.
- Schacht, W.H., A.J. Smart, B.E. Anderson, L.E. Moser, and R. Rasby. 1998.** Growth responses of warm-season tallgrasses to dormant-season management. *J. Range Manage.* 51:442-446.
- Sedivec, K.K., D.L. Dodds, and D. Gault. 1991.** Range Site Identification. N.D. State Univ. Ext. Serv. Bull. R-580 (Revised). N.D. Univ. Fargo. 7 pp.
- Shiflet, T.N. 1994.** Rangeland cover types of the United States. *Soc. Range Manage.* Denver, CO. 152 pp.
- Short, L.R. and E.J. Woolfolk. 1956.** Plant vigor as a criterion for range condition. *J. Range Manage.* 9:66-69.
- SPSS. 2000.** Statistical procedures for social science. Marketing Dept. SPSS. Inc. Chicago, Ill.
- Steel, R.G.D and J.H. Torrie. 1980.** Principles and Procedures of Statistics. (2nd Edition), McGraw-Hill, New York Company. New York, NY.

- Trent, J.D., L.L. Wallace, T.J. Svejcar, and S. Christiansen. 1988.** Effects of grazing on growth, carbohydrate pools, and mycorrhizae in winter wheat. *Can. J. Plant Sci.* 68:115-120.
- Ulmer, M.G. 1987.** Soil survey of Adams County, North Dakota. U.S. Gov. Print Off. Washington, D.C. 174p.
- United States Department of Agriculture Natural Resource Conservation Service. 2001.** Range Technical Guidelines. Bismarck, ND.
- United States Department of Agriculture Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory 2002.** Fire Effects Information System, [Online]. Available: <http://www.fs.fed.us/database/feis/> [2002, February].
- Vallentine, J.F. 2001.** Grazing Management. Academic Press. San Diego, CA..
- Whitman, W.C., D.W. Bolin, E.W. Klosterman, K.D. Ford, L. Moomaw, D.G. Hoag, and M.L. Buchanan. 1951.** Carotene, protein, and phosphorus in grasses of western North Dakota. *N.D. Agr. Exp. Sta. Bull. No. 370.* Fargo. 55p.
- Wiesner, H.C. 1980.** Soil survey, Perkins County, South Dakota. U.S. Gov. Print Off. Washington, D.C. 195p.