

# 1988 NORTH DAKOTA WEED CONTROL RESEARCH





## SUMMARY OF 1988 WEED CONTROL EXPERIMENTS

Crop and Weed Sciences Department North Dakota State University Fargo, North Dakota

R. C.	G.	Ahrens Lym Messersmith Nalewaja	T. J. Jerzy	Manthey Peters Palazynski Woznieca
0.	υ.	Marewaja	Zenon	Woznieca

D. E. Peterson - Crop and Weed Sciences Department - Extension A. G. Dexter - Crop and Weed Sciences Department - Extension North Dakota State University - University of Minnesota

## **Technicians**

## Graduate Research Assistants

K. M. Christianson M. G. Ciernia J. L. Luecke R. F. Roach				Swenson Thorsness
--	--	--	--	----------------------

Experiments conducted in cooperation with: Curt Thompson, Minot Experiment Station Neil Riveland, Williston Experiment Station John Lukach, Langdon Experiment Station Blaine Schatz, Carrington Experiment Station Tom Conlon, Dickinson Experiment Station

Reference to commercial products or trade names is made with no intended endorsement and failure to mention products or trade names is done with no intended discrimination by North Dakota State University. Experiments with pesticides on non-labelled crops or target species does not imply endorsement of non-labelled uses of pesticides by North Dakota State University.

#### SUMMARY OF 1988 WEED CONTROL EXPERIMENTS

Crop and Weed Sciences Department Worth Dakota State University Fargo, Worth Dakota

M. H. Ahrens
R. G. Lym
R. G. Messersmith
G. Messersmith
Jerzy Palanynsk
J. D. Malewaja
Jennehoznikoa

D. E. Peterson - Crop and Waed Sciences Department - Extension A. G. Dexter - Crop and Weed Sciences Department - Extension North Daketa State University - University of Minhesota

## Technicians

## Graduate Research Assistants

K. M. Christianson M. W. Bredehoeft
M. G. Clernia D. W. Flake O. R. S.
J. L. Luecke J. A. Hemenway K. B. T.
R. F. Roach P. M. McMullan

Experiments conducted in cooperation with:
Curt Thempson, Minot Experiment Station
Neil Riveland, Williston Experiment Station
John Lukach, Langdon Experiment Station
Blaine Schafz, Carrington Experiment Station
Tom Conlon, Dickinson Experiment Station

Reference to commercial products or trade names is made with no intended endorsement and failure to mention products or trade names is done with no intended discrimination by North Dakoka State University. Experiments with posticides on non-labelled crops or target species does not imply endorsement of non-labelled uses of pesticides by North Dakota State University.

## Table of Contents

White Section: Experiment titles, climatic, edaphic and general informat	ion
Sugarheet weed control titles	
Sugarbeet weed control in crops titles	8 ]
General weed control in crops, titles  Perennial weed control, titles	I
Weed control in reduced tillage systems, titles	ΙΪ
Climatic data, Amenia/Prosper	7
Climatic data, Carrington	VI
Climatic data, Casselton	VII
Climatic data, Crookston	
Climatic data, Fargo	IX
Climatic data, Langdon	X
Climatic data, Minot	XI
Climatic data, Pillsbury/Page	VIII
Climatic data, Williston	XIV
Key to abbreviations	XV
List of herbicides tested	XVI
Soil characteristics	XXI
	AAI
Gold Section: Sugarbeet Weed Control	
Fall and sping applied herbicides, Amenia	
Fall and sping applied herbicides, Crookston	1
Soil applied herbicides, Bathgate	2
Soil applied herbicides, Clara City	3
Soil applied herbicides, Crookston	
Soil applied herbicides, Hillsboro	5
Soil applied herbicides, Mooreton	6
Soil applied herbicides, Raymond	7 8
Soil applied herbicides, Robbin	9
Soil applied plus postemergence herbicides, Bathgate	10
Soll applied plus postemergence herbicides, Clara City	12
Soll applied plus postemergence herbicides. Crookston	14
Soil applied plus postemergence herbicides, Hillshoro	16
soil applied plus postemergence herbicides, Mooreton	18
soil applied plus postemergence herbicides, Raymond	20
soil applied plus postemergence herbicides, Robbin	22
Common cocklebur control, Raymond	24
Effect of herbicides on sugarbeet root rot, Hector	25
Trifluralin application at different growth stages. Amenia	26
Simulated herbicide drift on sugarbeets, Amenia	28
Simulated herbicide drift on sugarbeets, Amenia	30
Herbicides on weed free sugarbeets, Fargo	31
Herbicides on weed free sugarbeets, St. Thomas	32
Time of application of sethoxydim and clopyralid plus additives. Amenia	34
Multispecies screening of PPI, Pre, and Postemergence herbicides. Fargo	36
Injury to sugarbeets and other crops from herbicide carryover. Fargo	38
Three species evaluation of the effect of additives on antagonism. Amenia	40
Broadleaf herbicides plus additives, Amenia	42

Green section: General Weed Control in Crops

Table of contents		
	Dago	File
Wild oats control in wheat, Fargo	1	WO2
Uild oate control in wheat Minor		WUT
Wild oats control in wheat Williston		WUS
Antagonism of wild pats control Williston		WUO
Antagonism of wild gate control Minot		WU/
Antagonism of wild nate control bargo		WUO
AC 222,293 plus bromoxynil and surfactant, Fargo Diclofop plus adjuvants, Fargo	8	W09
Diclofop plus adjuvants, Fargo	9	WO10
Wild nate and broadleaf herbicide combinations, ta	rdo 10	WULL
Antagonism of diclofon by various compounds, Fargo	11	WO12
Evaluation of harbicides for wild dats control. ta	ran 12	W013
Adjuvante with AC 222 203 Williston	13	WO14
Wild nate control evaluation, langdon	14	WH22
Wild out control in durum wheat Minot 1988	13	Wh26
Wild oats herbicides with adjuvants, Prosper	16	W015
Wild bass herbierass with angulary		
Hard Red Spring wheat cultivar response to herbici	des, Minot 17	VA1
Oats cultivar response to sulfonyl urea herbicides	, Minot 18	VA2
Duran wheat recorded to harbicides Minot		VA3
Wheat cultivar response to herbicides, Williston	20	VA4
Small grain response to herbicides, Fargo	21	WH1
Dwardlast wood control in whost Williston	//	WH2
Dwardlast wood control in wheat landdon	()	WH3
Proadleaf weed control in oats, Minor		WH4
Recadles weed control in wheat. Larrington	23	WH5
Rroadleaf weed control in wheat. Casselton	20	WH6
Broadleaf weed control in wheat with sulfonylureas	c, Casselton 27	WH7
Cl Cd control in wheat Eaven	7X	WH8
Preemergence fortail control in wheat. Fargo	29	WH9
postamerophie inviair control in wheat, laide	30	WH10
2 / D diothanolamino with carbonates FYD I FARGO	) 5	WH11
2 / D diothanolamine with carbonates FYD / FARGO	) 3/	WH12
		WH13
Carbonate antogonism of 2.4-1). Exp 1. Fargo	34	WH14
Carbonato antagonism of 2 4-11 typ 2 targo	33	WH15
Codium bicambonato antagonism of 2 4-11 Fargo		WHZN
DAS 514 for wood control in Wheat FYD L FARGO -	3/	WH16
DIC PIN +ON MOOD CONTROL IN MUDAL FYILL FAILIN		WH17
C 22121 +on propertions mond control in Mudal Farm	1 )7	WH18
Preemergence weed control in wheat, Williston	40	WH19.W
Small grain response to postemergence herbicides,	Fargo 41	WH19
CGA-131036 fall and spring applied on small grains	s, Williston 42	WH20
Fenoxaprop and HOF-7113 on Durum and HRS wheat. La	angdon 43	WH21
Hammaring for wood control landdon	44	WH23
Green foxtail control in wheat, Minot 1988	45	WH25
other crops from heroloide carryover, Fargo 38		

Green Section Cont.

Mond control to co			
weed control in flax. Fargo		FX1	
weed collect in flax. Carrington			
weed control in flax, williston		FX2	
Weed control in flax, Minot		FX3	
Acifluorfen for Flax designation Consolter	49	FX4	
Acifluorfen for Flax desiccation, Casselton	50	FX8	
Dessication of flax, Minot 1988	51	FX10	
weed control in flax, langdon		FX9	
Flax variety response to sulfonyl urea herbicides	53	FX11	
		LVII	
Weed control in sunflower, Carrington			
Weed control in sunflower, Minot		SF5	
Weed control in sunflower, Williston	55	SF6	
Weed control in sunflower, Williston	56	SF7	
Weed control in sunflower, Casselton	57	SF8	
Postemergence AC 222,293 in sunflower, Fargo	58	SF9	
AC 222,293 on weed free sunflower, Casselton	59	SF10	
		3110	
Weed control in soybean, Carrington	<b>CO</b>	00=	
Preemergence weed control in soybean, Casselton	60	SB5	
Postemergence weed control in soybean, Casselton		SB15	
Herbicide combinations nectors restaurant in Soybean, Casselton	62	SB16	
Herbicide combinations postemergence in soybean, Casselton	63	SB22	
beneazon with aujuvants, tasspiton	64	SP25	
IMAZECHAPYI WILLI AUTUVANTS. LASSELTON	65	SP27	
Dash volume with sethoxydim plus bentazon, Prosper	66		
	00	SP32	
Postemergence broadleaf weed control in dry bean, Casselton -	C 7	DD1	
Weed control with imazethapyr in dry bean, Casselton	67	DB1	
maze shapy in dry bean, cassellon	68	DB2	
Weed control in corn, Casselton			
Weed control in com, Cassetton	69	CO1	
Weed control in corn, Carrington	70	CO3	
Postemergence weed control in corn, Casselton	71	CO2	
		COL	
Weed control in grain sorghum, Carrington	72	GS1	
		651	
Preemergence weed control in canola, Langdon	70		
Preemergence Weed control in canola, Langdon 1988	73	CA1	
	74	CA2	
Weed control in safflower, Williston			
most control in sairlower, williston	75	SA1	
Annual forego and			
Annual forage crops response to herbicides, Minot	76	FOR1	
	, ,	TORT	
Commercial petroleum oils with sethoxydim, Exp 1, Fargo	77	CDO	
Commercial Delivienii olik With cathovidim Fun o F.		SP3	
Commercial Deciroled III Olly With Fills 71 ton Duconos	78	SP43	
Commercial Deliforenii niic With HDY Vasho I alaa	79	SP4	
Sethoxydim formulations with addition to the set of the	80	SP44	
Sethoxydim formulations with adjuvants, Exp 1, Prosper	81	SP11	
occitor Jaim Tolling at 10115 WITH an 1110 ant C Lyn 2 Easter	82	SP11Y	
oction, drill crille-or-oldy appril cation for I canad	83	SP12	
Section Julie - 01 - (dv d))) 1 (at 10n f vn 2 Drochon	84	SP13	
	85		
formation basished blass	00	SP15	

Green section cont.

Sethovydim nlus adiuvants, Fargo	88	SP19
Sethoxydim plus adjuvants, FargoFluazifop plus adjuvants, Fargo	89	SP20
Clethodim plus adjuvants, Fargo	90	SP21
Clethodim plus adjuvants, largo	91	SP22
DPX-Y6202 plus adjuvants, Exp 1, Fargo DPX-Y6202 plus adjuvants, Exp 2, Fargo	92	SP23
Fenoxaprop plus adjuvants, Prosper	93	SP33
Haloxyfop plus adjuvants, Fargo	94	SP34
Sethoxydim with NaHCO3 or Na2CO3, Fargo	95	SP26
DPX-R9674 with adjuvants, Fargo	96	SP28
2,4-D with adjuvants, Fargo 1988	97	SP29
Sethoxydim with adjuvants and antagonism, Fargo	98	SP30
Fluazifop with adjuvants and antagonism, Prosper	99	SP31
Additional with fluarifon. Fargo	100	SP39
Adjuvantes with the English Court of the Cou	101	SP40
Adjuvants with sethoxydim, FargoGlyphosate with adjuvants, Prosper	102	SP41
Paraquat with adjuvants, Prosper	103	SP42
False chamomile control on fallow and or noncrop land, Minot -	105	CAM
Common milkweed control on fallow, Minot	106	PER
weed control in soybean, Castelton		

Pink Section: Perennial Weed Control Page Fluroxypyr alone and with auxin herbicides for leafy spurge control ...... 1-2 Evaluation of sulfometuron applied mid-summer and fall for leafy spurge control ..... 3-5 Sulfometuron applied alone and with auxin herbicides as an annual treatment for leafy spurge control ..... 6-9 Leafy spurge control under trees and along waterways ...... 10-12 Leafy spurge control with various herbicides combinations Picloram and 2,4-D combinations treatments for long-term leafy spurge management ...... 17-18 Leafy spurge control with low rate annual picloram and 2,4-D Leafy spurge control with picloram plus dicamba or various 2,4-D formulations applied annually ...... 21-23 Late-season leafy spurge control with sulfometuron and picloram ..... 24-25 Leafy spurge control with resulting forage production from several herbicides treatments ...... 26-27 Leafy spurge control following a nine-year management program ...... 28-30 

Triazine-tridiphane mixtures for postemergence grass control in corn, Fargo	C020
Soybeans, Fargo	SBWT CMD1 CMD4 SB1 SB2 SB3 SB4 SB20 SF3 SF1 SF2 FA18 FA19 FA20 FA21 FA23 CMD3 FA21 FA345 FA131415 FA131415 FA131415

CLIMATIC DATA - PROSPER/AMENIA 1988

		Precipitation				Apri	<u> </u>	May		Jur		Jul		Augu	_
Date	April	May	June	July	Aug.	Max.	Min.	Max.	Min.	Max.	Min.	Max.	Min.	Max.	Min
1		Farg	580	ybea	.62			82	58	91	64	77	54	86	63
2								82	54	96	58	87	47	88	67
3								80	43	93	62	92	58	81	63
4								66	35	96	61	92	67	77	54
5								77	35	98	63	107	73	92	55
6								85	49	96	58	106	81	97	60
		.27		.23	.23	83	34	70	50	100	58	86	61	83	63
7		.61		. 23		75	39	74	42	82	58	91	56	76	51
8						46	25	73	46	82	39	90	58	90	49
					.23	56	23	71	48	90	46	76	53	97	70
10					. 23	69	24	76	54	98	60	81	46	92	65
11		.23	.19	.03	.03	69	30	82	38	79	61	88	55	90	67
12		. 23	.03		1.29	52	28	66	27	81	60	90	61	75	66
13		07	.47		102/	49	19	82	49	66	57	92	53	86	61
14		.03	.41			55	19	61	37	78	51	89	57	95	64
15		.03				82	21	68	32	79	47	91	51	103	73
16						47	26	78	40	90	58	87	56	76	66
17		07				49	13	94	61	98	66	86	61	82	63
18		.03		07		61	12	65	55	100	64	89	52	83	54
19		.03		.07		46	20	73	51	97	55	83	52	88	64
20		70	.19			51	17	58	52	94	68	90	51	82	68
21		.39				53	21	73	49	84	60	93	52	88	5
22		.03				54	28	85	43	82	61	97	63	81	53
23			.43			67	21	81	44	95	63	86	49	90	5
24						53	29	90	52	83	58	92	51	79	4
25			Farg			46	32	88	57	67	54	100	62	85	4
26		.07					27	88	53	94	61	107	71	72	4
27						63	24	91	66	86	62	98	75	70	3
28		.07	1			72		93	65	80	56	95	64	75	3
29	, lle					80	31 51	93 92	67	82	49	94	53	82	4
30	.15					77	וכ		68	02	7/	98	70	85	6
31								94	00			70	, ,	0,5	

CLIMATIC DATA - CARRINGTON 1988

		Pre	cipita	tion	-	April M			lay June			Ji	ıly	August	
Date	April	May	June	July	Aug.	Max.	Min.	Max.	Min.	Max.	Min.	Max.	Min.	Max.	Min.
1	33	.01	.50	.10	.95	58	27	71	59	90	61	75	57	68	57
2		.02			.02	61	30	69	47	88	57	84	52	82	
3			.14		.14	69	29	56	37	90	57	88	54	77	61 63
4				T		63	40	60	30	93	64	88	60	80	51
5				.30		54	31	72	28	94	61	100	65	91	56
6		T				72	28	84	47	95	60	102	65	96	65
7		.11		.18	T	82	36	76	49	97	60	99	57	91	66
8		T			T	67	44	59	40	86	64	91	53	75	66 47
9						46	26	75	35	82	51	90	60	89	47
10						53	23	75	41	93	54	82	49	99	64
11		.17				69	34	75	50	98	62	78	42	91	64
2		.20	.34	.19	.20	66	33	78	43	82	57	85	55	91	58
3					.09	52	29	67	26	82	52	85	58	88	63
4		T	.23			50	20	73	44	65	54	89	57	86	
5		.01		.08		54	15	72	44	73	46	88	62	95	55 66
6					.89	81	32	72	27	76	44	93	47	93	65
7				T		45	26	83	44	94	54	87	48	80	
8				.02		48	13	84	62	94	64	80	53	78	60
9		.09		T		60	18	71	53	93	61	80	48	82	58
0						49	21	72	52	97	55	80	46	87	52
1			.43			49	19	70	47	93	64	88	48	86	56
2						48	19	74	41	84	54	92	53	85	65
3			.63			54	17	82	42	89	58	86	65	80	49
4						54	32	84	45	95	67	85	50	88	51 49
5						53	26	90	54	86	56	95	58	86	49
6		.19				49	31	85	49	84	53	101	54	77	
7					.03	61	24	85	46	99	61	104	58		43
8		.35				71	31	89	59	85	69	102	70	70	38
9	.40					81	31	02	63	76	55	90	70 62	68	34
0						78	49	90	57	77	58	93	69	75 75	<b>3</b> 5
1								92	60		-		69	75 75	50 50

CLIMATIC DATA - CASSELTON 1988

-		Prec	ipitat	ion		Apr	il	Ma	У	Ju	ine	<u>July</u>		Aug	ust
Date	April	May	June	July	Aug.	Max.	Min.	Max.	Min.	Max.	Min.	Max.	Min.	Max.	Min.
114	45	E		89		, n	20	7/	48	93	63	85	55	98	62
1					.07	45	20	74			58	76	49	94	63
2					.47	60	38	80	60	89		85	62	89	66
3						61	34	81	42	94	62	90	60	82	56
4					.03	62	39	78	36	92	62	10,000			53
5					.01	69	26	68	36	96	61	94	70	79	
6						61	25			96	58	107	83	92	54
7		T		.22		74	32	84	41	96	58	106	64	80	49
8		.46		.16	.18	83	50	68	43	98	58	83	57	83	50
9						75	33	75	43	86	40	92	59	76	49
10						46	22	74	48	81	49	85	52	88	52
11					N8	57	26	74	48	90	53	77	46	99	66
12		.14			.01	70	36	78	54	97	64	84	56	98	66
13		.01	.42	.17	.93	49	32	83	25	78	59	88	55	94	66
14		.02	.53		.25	56	18	65	43	82	58	90	55	78	61
15			.02			51	18	82	51	73	45	93	61	87	64
16			102			55	24	66	31	74	46	90	53	98	64
17						82	24	75	35	79	54	90	58	105	64
18					.03	44	13	78	45	90	66	88	64	78	66
		.18			.05	50	13	94	57	100	63	87	54	86	55
19				.06		62	23	68	53	98	62	87	52	83	55
20		.04	4 4	.00		46	19	74	54	95	56	84	53	88	71
21		.21	.16			53	19	58	50	93	56	89	56	95	68
22		.31	20			55	27	75	47	94	58	93	60	88	54
23			.22				25	85	46	90	57	98	60	83	55
24						55			52	94	66	87	53	92	54
25						68	30	83	55	93	60	91	55	80	43
26						56	32	87					67	82	49
27		.10			T	55	29	90	49	94	62	98	79	73	40
28						68	26	87	60	93	55	106	74	71	41
29						72	31	91	63	91	56	96	74		
30	.90					80	48	90	64	80	50	94		76	42
31				.04				92	67			96	75	81	52

CLIMATIC DATA - CROOKSTON 1988

			pitati		-	Apr			ay	Jur	ne	Ju	ıly	Aug	August	
Date	April	May	June	July	Aug.	Max.	Min.	Max.	Min.	Max.	Min.	Max.	Min.	Max.	Min	
1					.89	51	32	80	62	87	60	77	48	73	63	
2			.01		T	59	33	81	42	93	65	83	57	84	65	
3						55	37	74	34	91	61	89	67	83	57	
4					.07	54	33	63	33	93	69	91	73	80	58	
5				.08		57	28	72	39	94	65	104	71	92	60	
6		.05		.12		68	34	85	59	94	65	100	66	96	74	
7		.41		.01		76	46	75	45	95	60	80	52	84	49	
8		.08				73	32	71	43	69	38	88	62	74	50	
9						42	26	70	43	79	54	86	54	85	59	
10					T	51	31	68	46	87	63	71	45	97	70	
11		.08				61	35	70	51	95	67	78	50	95	68	
12		.25	.30	.46	.21	63	30	73	28	80	59	83	62	88	66	
13			.49		.26	45	21	71	34	82	55	84	54	73	64	
4		.23	.11			41	19	78	44	74	52	85	62	85	67	
15			T		.02	51	26	45	35	74	46	85	50	94	69	
16						77	25	63	40	75	56	84	57	90	65	
17				T		40	14	74	52	86	54	84	63	76	64	
18		.07	.28	.01		45	13	89	55	89	61	84	50	81	51	
19				.05		57	21	61	56	95	53	84	57	81	59	
20		.11	.30			44	18	69	53	90	67	81	52	86	70	
21		.14			.02	46	26	56	46	88	57	87	55	83	58	
22			.03			50	28	75	43	82	60	92	67	86	58	
23			T	.32		52	26	82	43	81	61	91	62	77	65	
24						62	32	75	52	90	59	84	50	87	53	
25						44	29	86	52	81	52	86	65	75	41	
26		.05			.06	49	22	82	49	82	61	95	65	80	47	
27			T			60	24	87	62	89	66	101	71	69	47	
28						70	41	89	65	79	51	94	71	68	43	
29						77	51	91	69	80	55	89	52	68	42	
50	T			.12		77	55	89	71	80	56	89	65	78	50	
1		.14		.52	.03			91	66			91	60	83	52	

CLIMATIC DATA - FARGO 1988

		Prec	ipiat	ion		Ap	ril	Ma	ау	J	une		uly	Au	gust
Date	April			July	Aug.	Max.	Min.	Max.	Min.	Max.	Min.	Max.	Min.	Max.	Min
X		77		03		55 (	38		8 (1)		,,		50	00	
1					.43	55	24	80	60	89	66	75 05	59	88	64
2	T				.02	61	29	82	54	93	62	85	53	87	67
3	T				T	53	39	79	42	92	62	91	62	83	63
4	T				.01	65	40	68	37	93	66	92	69	78	56
5						60	34	77	36	94	69	106	78	92	56
6		T		.09		71	27	83	55	94	64	102	77	97	64
7		.28		.20	.09	80	36	71	50	97	66	86	64	86	64
8		T			T T	78	39	75	42	81	56	89	58	76	50
9		.07		T		45	27	71	45	80	42	89	63	88	52
10					T	54	26	70	49	89	55	75	52	98	71
11					.02	67	29	75	55	96	64	79	46	93	67
12		.35	.18	.15	.02	69	35	81	37	84	61	87	61	92	68
13			.03		1.36	51	26	64	24	82	60	90	63	78	66
14		.04	.69			48	20	82	51	67	58	92	57	87	62
15		.05	T			54	18	62	36	78	51	88	62	96	67
16	T				T	81	32	66	34	78	48	89	52	104	74
17	T				T	48	24	75	47	86	58	76	67		
18		T			06	47	12	92	63	97	66	86	63	82	66
19		.19		.01		59	17	68	56	98	63	88	56	82	54
20		.26		T	.03	46	24	72	54	95	56	82	55	88	65
21		.41	.06		09 T	51	20	57	52	93	69	89	54	82	71
22		.10			.16	54	19	73	51	83	57	93	62	87	61
23			.28	T		55	25	84	44	85	62	96	66	81	56
24	T			133		68	31	77	45	94	66	86	53	89	51
25	- 65					52	28	87	53	83	58	89	56	78	50
26	T III	.04			00 T	45	29	86	55	85	53	97	65	83	47
27	83	.04			10	62	27	85	54	03	65	105	71	71	46
28		T		.08		71	32	90	68	86	62	96	74	71	42
29		•		.00		79	44	90	66	80	58	94	67	74	40
30	.01	т				76	52	91	69	81	55	93	57	80	50
	,01			т	т	, 0	J.L	92	71	٥.		98	74	84	65
31		.03		1	1			76	11			70	14	04	05

ΧI

CLIMATIC DATA - LANGDON 1988

Date	A==21		cipita			Ap	oril	M	lay	J	une	.1	July Aug		iai io t
Date	April	May	June	July	Aug.	Max.	Min.	Max.	Min.	Max.		Max.		Max.	<u>ıgust</u> Min
1						40	20	77	56	92	64	79			
2			.09		T	56	30	73	55	88	56	73	58	86	56
3			.02	?		62	20	75	37	85	56		48	74	58
4						62	33	52	28	88		79	52	78	59
5				1.00		60	28	50	23	92	60	89	59	79	51
6				.23		55	28	62	34		66	91	64	78	51
7		T		.34		65	30	79		95	64	88	68	92	59
8		.44				72	38		49	93	63	89	57	99	68
9						72	26	72	44	96	63	78	53	78	40
10						38		50	34	74	45	84	57	76	47
11							23	70	36	79	54	75	50	88	56
12		.13				52	28	65	43	89	65	64	47	98	60
13		.01	.02	7/		66	34	69	48	92	59	74	50	82	59
14		.03	.02	.36		59	24	62	24	75	50	76	57	87	61
15			25	1.04		45	17	68	32	81	53	79	52	71	60
16		.10	.25	.10	T	45	18	68	37	66	48	75	57	81	61
17		.01	.03		.10	52	26	50	27	67	40	78	47	94	66
18					.08	70	18	65	36	75	54	82	47	81	58
19			.06			39	13	77	54	92	61	81	54	76	55
		. 15		T		43	15	82	51	91	59	78	48	77	
20				.27		53	13	71	50	89	53	77	50		53
21		T				43	18	68	46	91	66	79	52	80	54
22						40	23	68	40	91	60	84		82	58
23			.05		.02	40	17	78	49	79	56		58	83	51
24						49	32	75	46	86	49	89	67	79	52
25			T			50	30	78	55	92		82	50	75	51
.6						47	23	93	46	_	57	81	55	82	49
7		.02			T	46	26	78		80	57	91	56	73	44
8					63	62	29	87	45	83	62	95	57	74	40
9		.08		.05		71	39		60	93	61	97	66	65	42
0		88 -		.05			52	91	62	79	51		61	65	38
1							22	90	64	79	48		50	70	44
								86	68			87	59	77	51

CLIMATIC DATA - MINOT 1988

		Preci	pitati	on		Apr	il	Ma	у	Ju	ine		lly_		ust
Date	April		June		Aug.	Max.	Min.	Max.	Min.	Max.	Min.	Max.	Min.	Max.	Min.
1				.84		46	26	83	50	86	62	62	57	86	62
2		.83				62	36	60	45	80	64	68	56	77	59
3		.09				72	41	45	36	88	52	82	62	75	57
4						76	41	44	34	91	58	90	60	76	48
5						60	28	48	31	94	70	86	64	83	57
						54	31	66	34	96	67	88	66	97	64
6				.03	.06	71	38	80	52	96	70	83	57	100	63
7		.22		.03		81	39	72	42	100	73	81	57	74	44
8		.07			25	46	25	58	36	95	57	87	55	73	50
9			.05			46	26	75	41	87	56	79	52	92	58
10		.06				57	30	73	45	98	71	71	44	96	63
11		.04			0.5	74	37	75	47	93	58	78	51	85	60
12						62		77	32	83	49	77	57	91	61
13		T	.48	.01	.17	52		62	43	82	53	85	60	69	59
14		.02	.72	.01		49		68	44	60	51	93	62	89	66
15			.12			56	25	56	34	70	50	81	54	95	69
16		.01				77	27	75	42	78	55	90	50	91	61
17	T					42	16	82	56	97	55	85	53	78	57
18		4.6		Т	Т	49	20	64	44	95	65	74	53	78	58
19		.11		.01		59	25	71	51	90	61	76	47	88	52
20		81				43	17	74	43	100	68	80	54	88	60
21		T				49	17	74	43	93	58	88	62	82	55
22					.02	46	22	78	49	88	63	94	66	79	55
23					.02	57	28	82	51	99	66	81	49	79	51
24						44	24	84	62	96	55	85	59	86	51
25						47	27	84	50	81	56	97	59	76	45
26					•	49	24	78	52	90	71	100	57	73	39
27		.02			Т	61	33	87	60	99	67	103		68	39
28						72	33 43	89	58	91	60	101		67	37
29			_			80	43 47	97	66	79	59	86		76	41
30	.09		. 2	.2		۵υ	47	85	69	. ,	- /	92		85	55
31								65	07			, _			

## CLIMATIC DATA - PILLSBURY AND PAGE 1988

Date	April	Precip May	oitation June	July
1		4 SE 6 SE	27 T	***
2 3				
1 2 3 4 5 6 7 8 9				
6		20		
8				
10				
11 12 13				
14				
15 16			1.04	
17 18				
19		66		
20				
23				
5		00		
26 27				
28	25			
9 0 1	.25			

CLIMATIC DATA - WILLISTON 1988

XIV

		Pr	ecipit	ation		Apr	·il	Ma			ine		ıly	-	<u>ust</u>
Date	April	May	June	July	Aug.	Max.	Min.								
1		.24			.02	64	28	71	44	79	58	78	52	89	57
2		.30			٢	73	32	65	40	82	55	89	55	83	58
3				Т		71	38	61	31	92	51	90	65	79	56
4	.14			.09		65	37	65	30	98	62	89	60	85	50
5						58	31	66	33	103	60	94	67	100	56
6						71	34	80	51	104	72	94	66	106	60
7		.01		.02		72	46	79	50	105	69	81	50	105	58
8	.01	.24				71	34	72	44	104	71	88	53	77	49
9	т.					48	26	76	40	96	61	88	56	92	54
10						58	26	79	43	105	65	81	50	93	63
11		Т				78	31	81	48	105	68	80	55	92	64
12				.07	.16	77	35	76	45	89	57	85	65	91	60
13			T		.03	59	27	75	37	81	55	84	59	90	63
14		.26	.76			58	21	75	48	72	50	94	62	95	59
15		020	.07		.05	67	26	67	49	73	48	96	62	99	58
16						77	35	83	40	83	54	86	63	99	74
17				.03		67	30	88	55	102	60	82	53	90	55
18		.03		.12		56	22	89	53	102	70	76	53	84	58
19					.04	61	22	74	42	95	63	77	53	85	59
20			.18			59	30	75	49	105	66	83	51	92	56
21			.03			53	21	77	44	95	69	91	57	92	57
22					Т	54	24	80	43	95	65	102	62	78	51
23					•	60	25	82	46	101	69	102	62	83	51
24						59	30	88	55	100	79	90	55	90	54
25						48	23	88	59	89	52	96	60	88	51
26		.01			T	50	26	83	65	99	68	99	60	79	50
27		.01			•	63	23	91	65	99	68	108	66	73	43
28		.32		.02		75	33	91	58	100	66	104	70	73	40
29						82	40	98	48	95	66	97	63	82	41
30			1.98			81	47	95	64	80	55	99	57	85	55
31			,.,0			-		88	58			100	62	84	48

## KEY TO ABBREVIATIONS AND EVALUATIONS

Crop injury, crop stand and weed control ratings are based on a visual estimate using a scale of 0 to 100 with 0 = no effect and 100 = complete kill. All preplant incorporated or preemergence treatments were applied in 17 gpa water at 35 psi through 8002 nozzle tips and all postemergence treatments were applied in 8.5 gpa water at 35 psi through 8001 nozzle tips except where

All treatments were applied with a bicycle wheel-type plot sprayer unless otherwise stated. Preplant incorporation was by field cultivator + harrow or as stated in table and preemergence incorporation was by harrowing twice.

Treatments with a + indicate tank mixtures, with an & indicate
formulation mixtures and with a / indicate a separate application.

<u>Species</u> Abww = Absinth wormwood Barl (Bar) = Barley Bdlf = Broadleaf Bygr = Barnyardgrass Cath = Canada thistle Cocb = Common cocklebur
Colq = Common lambsquarters Copu = Common purslane Cosf = Volunteer sunflower Dobr = Downy brome
Fach = False chamomile
Fibw = Field bindweed Fipc = Field pennycress
Flwe (Flix) = Flixweed
Foba - Foxtail barley
Fxtl = Foxtail species
Grft = Green foxtail
Gfpw = Greenflower pepperweed Howe = Horseweed KOCZ = Kochia Latu = Ladysthumb Lent = Lentils Lesp = Leafy spurge Mael = Marshelder Mesa = Meadow salsify Mil (Ftmi) = Foxtail millet Methods PPI = Preplant incorporated PEI = Preemergence incorporated Miscellaneous DF = Dry flowable F = FallFL = F = FlowableS = Spring L = Liquid G = Granules or gallon/A Inc = I = Incorporation %ir = inju = Percent injury rating
%sr = %std, strd = Percent stand reduction
HT = Plant height

Nabe = Navy beans Nfcf = Nightflowering catchfly Pest = Perennial sowthistle Pesw = Pennsylvania smartweed Powe = Pondweed Prle = Prickly lettuce Prpw = Prostrate pigweed Qugr = Quackgrass Rrpw = Redroot pigweed Ruth = Russian thistle Soyb (Sobe) = Soybean Soyb (Sobe) = Soybean
Spkw = Spotted knapweed
Sugb = (Sgbt) = Sugarbeet
Sunfl (Sufl, Cosf) = Sunflower
Tamu = Tansy mustard
Taoa = Tame oats
Tumu = Tumble mustard
Tymu = Tame yellow mustard
Vowh = Volunteer wheat
Wesa = Western salsify Wesa = Western salsify Wht = Wheat Wibw = Wild buckwheat Wimu = Wild mustard Wioa = Wild oats Yeft = Yellow foxtail

alk = alkanolamine dma = Dimethylamin bee = Butoxyethyl ester

PRE, PE = Preemergence P, PO, POST = Postemergence

dea = diethanolamine MS = modified seed oil PO, OC = Petroleum oil concentrate (17% emulsifier) SPK = Spike stage SURF = S = Surfactant Tswt = TW = Test weight WP = Wettable powder WK = Surfactant by DuPont X-77 = Surfactant by Ortho Yld = Yield

XVI
LIST OF HERBICIDES TESTED IN 1988

Common Name	Abbre-	100 with 0 a go a	Formulation	Trade Name
or Code Name	viation	Company	romutation	Name
		2 nozzle tips and	0.8 lb/gal	None
A-1237	A1237	Uniroyal	U.O LD/ gal	
		American Cyanamid	2.5 lb/gal	Assert
AC-222,293	AC-293	American cyanamia	200 (10, 3-1	
Imazamethabenz	dmmI			
a a 2/7 /00	Imep	American Cyanamid	2 lb/gal AS	Pursuit
AC-263,499 Imazethapyr	Incp	/ indicate a sep		
Illiazethapyi				
Acifluorfen	Acif	BASF	2 lb/gal E,S	Blazer
		Rhone Poulenc	2 lb/gal S	Tackle
			558	Barnyardgr
Alachlor	Alac	Monsanto	4 lb/gal E	Lasso
			4 lb/gal MT, 15%	me I gamma
	ng = man	al	2 lb/gal S	Amitrole T
Amitrole	Amit	Rhone-Poulenc	c injadi s	
DBSWCIO JOCID	A 4 m a	Various	80% WP, 90% DF,	Numerous
Atrazine	Atra	vai rous	4 lb/gal F	
			0991	
BAS-090		BASF	Surfactant	None
BY2-010				
BAS-51400H	BAS514	BASF		Facet
			- Hi	Name of the second
Benazolin	Bena	Nor-Am	50% SC	None
		7.05	4 lb/gal S	Basagran
Bentazon	Bent	BASF	4 (D/gat 3	565451 4
acadu cond	Brox	Rhone-Poulenc	2 lb/gal E	Buctril
Bromoxynil	ыих	Kilong I datelle	9	
Butylate + Safene	er Buty	ICI	6.7 lb/gal L 10	% G Sutan+
	191 - 110			
c-4243		Uniroyal	0.8 lb/gal	None
		9	75% ND	Amber
CGA-131036	CGA131	Ciba Geigy	75% WP	Allipet
و المعالم والمعالم	20147/	Ciba Geigy	75% DF	Beacon
CGA-136872	CGA136	Ciba deigy	10.00	
Chlorembor	Clam	Rhone-Poulenc	75% SP	Amiben
Chloramben	Ctall			
Chlorsulfuron	Clsu	DuPont	75% DF	Glean
3,110, 341, 4, 4,1				- incorpor
Cinmethylin	Cinm	DuPont	7 lb/gal EC	Cinch
act and had luce			Percent stand	- DYGZ , DJ
Clethodim	Clet	Valent	2 lb/gal	Select

or Code Name	Abbre- a viation			Trade
S. South Walle	VIACION	Company	Formulation	Name
Clomazone	Clom	FMC	4 lb/gal	Command
Clopyralid	Clpy	Dow	3 lb/gal S	Stinger
Clopyralid+2,4-	Clpy&2,4-D	Dow	0.38 + 2 lb/gal	S Curtail
Cyanazine	Cyan	DuPont	80% WP, 90% DF	Bladex
			4 lb/gal F	
Cycloate	Cycl	ICI	6 lb/gal E	Ro-Neet
Dalapon	Dala	Dow	75% SP	Dowpon
Desmedipham	Desm	Nor-Am	1.3 lb/gal E	Betanex
Desmedipham +				
Phenmedipham	Des & Phen	Nor-Am	0.65+0.65 lb/gal	E Betamix
Diallate	Dial	Monsanto	4 lb/gal E	Avadex
Dicamba	Dica	Sandoz	4 lb/gal S	Banvel
Dichlorprop		Rhone-Poulenc	4 lb/gal EC	Weedone 2,4-D
Diclofop	Difp	Hoechst-Roussel	3 lb/gal E	Hoelon
iethatyl	Diet	Nor-Am	4 lb/gal E	Antor
i fenzoquat	Dife	American Cyanamid	2 lb/gal S	Avenge
iquat	Diqu	Valent	2 lb/gal S	Diquat
PX-A7881	DPX-A7	DuPont	75% DF	Muster
PX-L5300	DPX-L5	DuPont	75% DF	Express
PX-M6316	DPX-M6	DuPont	75% DF	Harmony
PX-R9674 (DPX- 5300 + DPX-M6316)	DPX-R9	DuPont	75% DF	Harmony Extra
51 X 1105 107	E 189/41-8			
PX-Y6202(-44)	Qufp	DuPont		
Quizalofop)	lagyer a		0.75 lb/gal EC	Assure
N-1/07/0	165403.9			
2X-V9360 Scent	DPX-V9	DuPont	72% DF	Accent
dothall	Endo	Pennwalt	3 lb/gal S	Herbicide 273

Common Name	Abbre-		and the latest and the second	Trade
or Code Name	viation a	Company	Formulation	Name
EPTC	EPTC	ICI	7 lb/gal E	Eptam
Ethalfluralin	Etha	Elanco	3 lb/gal E	Sonalan
E CHOCK CO.				
Ethofumesate	Etho	Nor-Am	4 lb/gal F	Nortron
			1.5 lb/gal E	
Fenoxaprop	Fenx	Hoechst-Roussel	1.5 lb/gal E	Whip option
Fluazifop-P	Flfp-P	ICI	1 lb/gal E	Fusilade 2000
	A transact A			
Fluroxypyr	Flox	Dow	1.7 lb/gal	Starane
,				
Fomesafen	Fome	ICI	2 lb/gal	Reflex
Fosamine	Fosa	DuPont	4 lb/gal S	Krenite
, 554				
Glyphosate	Glyt	Monsanto	3 lb/gal S	Roundup
gt yphood to	,			
Haloxyfop	Halx	Dow	2 lb/gal	Verdict
na coxy rop				
HOE-00661		Hoechst-Roussel	1.68 lb/gal	Ignite
HOE GOOG!				
HOE-7113		Hoechst-Roussel	0.5 lb/gal	Puma
1106 1110				
HOE-7125 (fenox	caprop+MCPA	Hoechst-Roussel	0.75 lb/gal	Tiller
+ 2,4-D)	cop. op			
. 64 57				
HOE-7121 (feno)	caprop+MCPA	Hoechst-Roussel	0.5 lb/gal	
+ 2,4-D)				
. 2,40,				
Imazaquin	Imqn	American Cyanamid	1.5 lb/gal	Scepter
a managar				
Isoxaben	Isox	Elanco	75%	Vary 10
100/10011	8 7-57/27 3			
Imazethapyr	Imep	American Cyanamid	2.5 lb/gal	Pursuit
indre map).	- 40			
Lactofen	Lact	PPG	2 lb/gal S	Cobra
Lactorion				
Linuron	Linu	DuPont	50% WP, 4 lb/gal	F Lorox
Emaron	- 113			
MCPA	MCPA	Rhone-Poulenc	4 lb/gal E, S	Several
HOLK	11260 2501			
Metolachlor	Meto	Ciba-Geigy	8 lb/gal E	Dual
Hototadiitai		5mPGUE		
Metribuzin	Metr	Mobay	4 lb/gal F, 75% D	F Sencor
,1001 1000111	(0.00-03.5)	DuPont	4 lb/gal F, 75% D	F Lexone
		510900		
Metsul furon	Mets	DuPont	60% DF	Ally/Escort
metsutturon	Meta	<b></b>	**************************************	

Common Name	Abbre-			
or Code Name	viation <sup>a</sup>	Company	Formulation	Trade
			Tormatation	<u>Name</u>
Oryzalin	Oryz	Elanco	4 lb/gal F	Surflan
Paraquat	200		5941	out I tail
r ur aqua t	Para	ICI	1.5 lb/gal S	Gramoxone Super
			2 lb/gal S	Cyclone
Pendimethalin	Pend	American Cyanamid	4 lb/gal E	Prowl
Picloram	Picl	Dow	2 lb/gal S	Tordon 22K
Propachlor	Prcl	Monsanto	4 lb/gal F	Domasid
Droponil . None	on a second of section		12,341	Ramrod
Propanil + MCPA	Prnl + MCPA	Rohm & Haas	3.0 + 1.4 lb E	Stampede CM
Pyrazon	Pyra	BASF	( 2 1) ( ) -	
			4.2 lb/gal F	Pyramin
Pyridate	Pyri	Gilmore	3.75 lb/gal E	Tough
R-25788, Dichlormid	Demd	ICI	6 lb/gal E	None
R-33865, Dietholate	Ext	ICI	6 lb/gal E	None
R017-3664	None	Mobay	2 lb/gal	
RE-40885	None	Valent	80% WP	Benchmark
S-23121	V			berichiliar K
0 13121	None	Valent	0.83 lb/gal F	None
S-53482	None	Valent	0.07 167 1 -	
			0.83 lb/gal F	None
s-63596	None	Valent	0.89 lb/gal F	None
SAN-582H	None	Sandoz		,,,,,,,
		Sandoz	6 lb/gal E	None
SC-0051	None	ICI	3 lb/gal	None
SC-0098	None			None
00/0	None	ICI	1.7 lb/gal	None
Sethoxydim	Seth, Sth	BASF	1.5 lb/gal E	Poast
Gulfometuron	Sume			. 5451
	Suile	DuPont	75% DF	Oust
hiameturon		See DPX-M6316		

Common Name	Abbre- a viation	Company	Formulation	Trade Name
Triallate	Tria	Monsanto	4 lb/gal E, 10% G	Far-go
Triclopyr	Trcp	Dow	4 lb/gal	Garlon
Tridiphane	Trid	Dow	4 lb/gal E	Tandem
Trifluralin	Trif	Elanco	4 lb/gal E	Treflan
2,4-D	2,4-D	Various	Various E, S	Numerous
2,4-DB	2,4-DB	Various	2 lb/gal	Numerous

Abbreviations in the tables may consist of only the first one, two, or three listed letters when space was limited. Abbreviations of numbered compounds varies with available space, but usually was the first letters and numbers.

## SOIL TEST RESULTS AT VARIOUS WEED EXPERIMENT LOCATIONS

	Soil <u>Texture</u>	Organic matter	nН	N	lb/A	1/
Amenia, ND (Fall expt) Amenia, ND (Time of appl.) Amenia, ND (Treflan) Amenia, ND (bioassays) Amenia, ND (Drift 1) Amenia, ND (Drift 2) Bathgate, ND Carrington, ND Casselton, ND Clara City, MN Crookston, MN Dickinson Ranch HQ Fargo, ND (Sect. 22) Fargo, ND (sugarbeets) Hector, MN Hillsboro, ND Hunter, ND Langdon, ND Minot, ND Mooreton, ND New England, ND Pillsbury, ND Raymond, MN Robbin, MN St. Thomas, ND Valley City, ND (Sec 22)	Texture  Silt loam Silty clay Fine sandy load Silty clay Fine sandy loam Clay loam Silty clay Silty clay Silty clay Clay loam Silty clay Silty loam Loam Silty loam Loam Silty clay loam	matter  3.6 3.7 3.8 3.5 3.2 3.6 4.1 3.6 5.0 6.7 high - 4.4 6.0 5.1 4.0 - 7.4 4.6 2.7 3.0 5.8 5.9 am 9.7 am 4.4 4.2 9.4	pH 7.6 7.7 7.3 7.8 7.9 7.4 8.1 7.6 7.6 7.6 7.6 7.6 7.6 7.6 7.6 7.6 7.6	Applie 20 100 337 5 110 112 79 126 14 Fertili Fertili	P 67 66 64 75 66 63 1 zed 80 36 75 9 14 34 47 66 41	950 273 470 630 950 750 550 1360
Williston, ND	Silty clay Loam	3.2 2.3	7.5 6.8	137 Fertiliz	25 zed by	850 test

## SOIL TEST RESULTS AT VARIOUS WEED EXPERIMENT LOCATIONS

Fall and spring applied herbicides, Amenia, 1987-1988. Fall herbicide treatments were applied October 21, 1987 when the air temp. was 37F, soil temp. at six inches was 36F, and the wind was south at 8-12 mph. Spring herbicide treatments were applied May 5, 1988 when the air temp. was 71F, soil temp. at six inches was 62F, wind was northeast at 5-10 mph, soil was dry at water at 40 psi through 8002 nozzle tips to the center four rows of six row plots and incorporated with a rototiller set four inches deep for treatments containing EPTC or cycloate and two inches deep for metolachlor treatments. 'Mitsui Monohikari' sugarbeet was seeded 1.25 inches deep in 22 inch rows May evaluated July 1, 1988.

EPTC+Diethatyl (Spring) 2+4 34 99 97  HIGH MEAN LOW MEAN 99 97  EXP MEAN 0 87 68 C.V. % 87 4 13					
Cycloate (Fall)	110			Pigweed control	Foxtail
# OF REPS 20 7 22 4 4	Cycloate (Fall) EPTC+Cycloate (Fall) EPTC+Cycloate (Fall) Metolachlor (Fall) Metolachlor (Fall) Metolachlor (Fall) Metolachlor (Fall) EPTC+Diallate (Fall) EPTC+Diallate (Fall) EPTC (Spring) EPTC+Cycloate (Spring) Metolachlor (Spring) Metolachlor (Spring)	4 -2.5 3+2 2 3 4 6 1.5 3+2 2 5+2 2	0 6 10 0 0 5 9 26 20 20 16 0 10 34 34 0 12 87 15 20	92 88 97 97 97 99 99 99 99 99 99 99 99 99	75 83 92 68 88 95 97 73 87 88 92 81 87 97 97 68 85 13 16 22

## Summary

Fall applied EPTC at 4 lb/A caused greater sugarbeet injury than fall applied EPTC+cycloate at 2+2.5 or 3+2 lb/A. Fall EPTC at 4 lb/A gave less green foxtail control than fall EPTC+cycloate at 3+2 lb/A. Fall metolachlor at 2 or 3 lb/A gave sugarbeet injury and weed control similar to spring metolachlor at 2 or 3 lb/A. Only fall cycloate at 4 lb/A and fall EPTC+diallate at 3+2 lb/A gave less than 90% redroot pigweed control. Only fall EPTC at 4 lb/A, fall cycloate at 4 lb/A, fall metolachlor at 2 lb/A, and fall EPTC+diallate at 4+1.5 lb/A gave less than 80% green foxtail control.

Fall and spring applied herbicides, Crookston, 1987-1988. Fall herbicide treatments were applied October 22, 1987 when the air temp. was 38F, soil temp. at six inches was 35F, wind was northeast at 5-10 mph, and the soil was moist at 0-4 inches. Spring herbicide treatments were applied April 26, 1988 when the air temp. was 52F, soil temp. at six inches was 40F, wind was west at 5-10 mph, soil was dry at 0-1 inch, moist at 1-3 inches, and wet at 3-6 inches. All herbicides were applied in 17 gpa water at 40 psi through 8002 nozzle tips to the center four rows of six row plots and incorporated with a rototiller set four inches deep for treatments containing EPTC or cycloate and two inches deep for metolachlor treatments. 'Maribo Ultramono' sugarbeet was seeded 1.25 inches deep in 22 inch rows April 26, 1988. Common lambsquarters and redroot pigweed control and sugarbeet injury were evaluated June 1, 1988.

	November 1	Sugarbeet	Common Lambsquarters	Redroot Pigweed control
Treatment	Rate	injury	control	CONCION
	(1b/A)		(%)	
EPTC (Fall) Cycloate (Fall) EPTC+Cycloate (Fall) EPTC+Cycloate (Fall) Metolachlor (Fall) Metolachlor (Fall) Metolachlor (Fall) Metolachlor (Fall) EPTC+Diallate (Fall) EPTC+Diallate (Fall) EPTC (Spring) EPTC+Cycloate (Spring) Metolachlor (Spring) Metolachlor (Spring) Metolachlor (Spring) HIGH MEAN LOW MEAN EXP MEAN C.V. %	4 4 2+2.5 3+2 2 3 4 6 4+1.5 3+2 2 1.5+2 2	0 0 0 0 0 0 3 10 3 0 3 0 8 10 10 0 3 225 NS	84 93 91 92 75 77 94 98 93 94 95 96 73 89 89 98 73 89	65 83 64 84 98 98 99 99 78 58 81 94 99 99 99 93
LSD 5% LSD 1%		NS 4	16 4	19 4
# OF REPS		4	•	

## Summary

None of the treatments caused significant sugarbeet injury. Metolachlor at 2 lb/A fall applied gave weed control similar to 2 lb/A spring applied. Metolachlor at 3 lb/A fall applied gave similar redroot pigweed control but less common lambsquarters control than 3 lb/A spring applied. Metolachlor at 2 or 3 lb/A fall applied and at 2 lb/A spring applied gave or tended to give less common lambsquarters control than other treatments. All metolachlor treatments, spring EPTC+cycloate at 1.5+2 lb/A, and spring EPTC+diethatyl gave similar control of redroot pigweed which was or tended to be greater than redroot pigweed control from other treatments. Fall EPTC+cycloate at 3+2 lb/A gave better redroot pigweed control than fall EPTC+cycloate at 2+2.5 lb/A.

Soil applied herbicides, Bathgate, 1988. Preplant incorporated herbicides were applied and rototiller incorporated May 3 when the air temp. was 55F, soil temp. at six inches was 53F, wind was north at 8-12 mph, soil was dry at 0-2 inches and moist at 2-4 inches. The rototiller was operated four inches deep for treatments containing EPTC or cycloate and two inches deep for all inches deep in 22 inch rows May 3. Preemergence herbicides were applied May 3. All herbicides were applied in 17 gpa water at 40 psi through 8002 nozzle tips to the center four rows of six row plots. Redroot pigweed control and sugarbeet injury were evaluated June 25.

Treatment	Rate (1b/A)	Sugarbeet injury 	Redroot Pigweed control (%)
EPTC Metolachlor Metolachlor EPTC+Cycloate Metolachlor EPTC+Cycloate Cycloate Diethatyl Diethatyl Diethatyl (pre) EPTC+Cycloate+Diethatyl EPTC+Cycloate+Diethatyl EPTC+Cycloate+Diethatyl EPTC+Cycl/Diet (pre) Ethofumesate  HIGH MEAN LOW MEAN EXP MEAN C.V. % LSD 5% LSD 1% # OF REPS	2.5 1.5 2 1.5+1.5 3 2+2 4 4 1.5+1.5+4 1.5+1.5/4	6 3 4 8 8 13 0 4 0 18 10 0 18 10 6 94 8 11 4	71 64 79 66 90 73 49 64 0 94 69 83 94 0 67 12 12 16 4

## Summary

Only EPTC+cycloate at 2+2 lb/A and EPTC+cycloate+diethatyl at 1.5+1.5+4 lb/A caused sugarbeet injury over 10%. Only metolachlor at 3 lb/A and EPTC+cycloate+diethatyl at 1.5+1.5+4 lb/A gave 90% or greater redroot pigweed control.

Soil applied herbicides, Clara City, 1988. Preplant incorporated herbicides were applied and rototiller incorporated April 19 when the air temp. was 63F, soil temp. at six inches was 49F, wind was north at 8-12 mph, soil was dry at 0-1 inches, moist at 1-3 inches, and wet at 3-6 inches. The rototiller was operated four inches deep for treatments containing EPTC or cycloate and two inches deep for all other PPI treatments. 'KW 3265' sugarbeet was seeded in 22 inch rows April 20. Preemergence herbicides were applied April 19. All herbicides were applied in 17 gpa water at 40 psi through 8002 nozzle tips to the center four rows of six row plots. Sugarbeet injury was evaluated May 23.

		Sugarbeet
Treatment	Rate	injury
	(1b/A)	(%)
	2.5	9
EPTC	1.5	ő
Metolachlor	2	i i
Metolachlor	1.5+1.5	$\frac{1}{3}$
EPTC+Cycloate	3	0 1 3 4
Metolachlor	2+2	13
EPTC+Cycloate Cycloate	4	
Diethatyl		3 0 0
Diethatyl (pre)	4 4	0
EPTC+Cycloate+Diethatyl	1.5+1.5+4	16
EPTC+Cycloate/Diet (pre)	1.5+1.5/4	5
Ethofumesate	3	0
		16
HIGH MEAN		0
LOW MEAN		4
EXP MEAN		86
C.V. %		
LSD 5%		5 7
LSD 1% # OF REPS		4
# UT KEPS		

#### Summary

EPTC at 2.5 1b/A, EPTC+cycloate at 2+2 1b/A, and EPTC+cycloate+diethatyl at 1.5+1.5+4 1b/A caused significant sugarbeet injury. Weed populations were too erratic for evaluation of weed control.

Soil applied herbicides, Crookston, 1988. Preplant incorporated herbicides were applied and rototiller incorporated April 26 when the air temp. was 52F, soil temp. at six inches was 40F, wind was west at 5-10 mph, soil was dry at 0-1 inches, moist at 1-3 inches, and wet at 3-6 inches. The rototiller was operated four inches deep for treatments containing EPTC or cycloate and two inches deep for all other PPI treatments. 'Maribo Ultramono' sugarbeet was seeded 1.25 inches deep in 22 inch rows April 26. Preemergence herbicides were applied April 26. All herbicides were applied in 17 gpa water at 40 psi through 8002 nozzle tips to the center four rows of six row plots. Common lambsquarters and redroot pigweed control and sugarbeet injury were evaluated June 1.

			Common	Redroot
Treatment	Do.+-	Sugarbeet	Lambsquarters	Pigweed
TT CACINCTIC	Rate	injury		control
	(1b/A)		(%)	
EPTC	2	00		
Metolachlor	1.5	20	96	92
Metolachlor	1.5	5	66	94
EPTC+Cycloate		0	77	97
Metolachlor	1.5+1.5	11	97	92
EPTC+Cycloate	1 5.2	10	82	99
EPTC+Cycloate	1.5+2	9 5	97	94
EPTC+Cycloate	1+2.5		97	97
EPTC+Cycloate	2+2	26	98	98
EPTC+Cycloate+Triallate	1.5+2.5	15	98	95
Cycloate	1.5+2+2	13	98	99
EPTC+Cycl/Meto (pre)		5	93	94
Diethaty1	1.5+2/2	43	99	99
EPTC+Cycloate+Diethatyl	1 5.2.2	0	20	79
Cycloate+Diethatyl	1.5+2+3	36	97	99
	3+3	11	91	96
Diethatyl	1.5+1.5+4	28	98	99
EPTC+Cycl/Diethatyl (pre)	1 5.0/4	3	20	71
Ethofumesate (pre)	1.5+2/4	36	99	99
EPTC+Cycloate/Etho (pre)	3.5	3	87	98
Li Totoje Toute/Lino (pre)	1.5+2/3	50	99	99
HIGH MEAN				
LOW MEAN		50	99	99
EXP MEAN		0	20	71
C.V. %		16	85	94
LSD 5%		45	9	7
LSD 1%		10	11	9
# OF REPS		14	14	12
" OF INEFS		4	4	4

#### Summary

EPTC alone or EPTC+cycloate with EPTC at 2 lb/A caused or tended to cause greater sugarbeet injury than EPTC+cycloate that included EPTC at 1 or 1.5 lb/A. Addition of diethatyl, ethofumesate, or metolachlor to EPTC+cycloate gave greater sugarbeet injury than EPTC+cycloate alone. Metolachlor gave less control of common lambsquarters than EPTC, EPTC+cycloate, or cycloate. Diethatyl gave less control of common lambsquarters and redroot pigweed than other treatments.

Soil applied herbicides, Hillsboro, 1988. Preplant incorporated herbicides were applied and rototiller incorporated April 25 when the air temp. was 48F, soil temp. at six inches was 43F, wind was north at 20 mph, soil was dry at 0-1 inches, moist at 1-3 inches, and wet at 3-6 inches. The rototiller was operated four inches deep for treatments containing EPTC or cycloate and two inches deep for all other PPI treatments. 'Maribo Ultramono' sugarbeet was seeded 1.25 inches deep in 22 inch rows April 25. Preemergence herbicides were applied April 25. All herbicides were applied in 17 gpa water at 40 psi through 8002 nozzle tips to the center four rows of six row plots. Redroot pigweed and green and yellow foxtail control and sugarbeet injury were evaluated June 6.

<u>Treatment</u>	Rate (1b/A)	Sugarbeet injury	Gr.&Ye. Foxtail control (%)	Redroot Pigweed control
EPTC Metolachlor Metolachlor EPTC+Cycloate Metolachlor EPTC+Cycloate Cycloate Diethatyl Diethatyl EPTC+Cycloate+Diethatyl EPTC+Cycl/Diet (pre) Ethofumesate	2.5 1.5 2 1.5+1.5 3 2+2 4 4 1.5+1.5+4 1.5+1.5/4	31 0 10 8 9 21 6 4 0 39 21 0	98 78 90 96 90 98 92 57 19 98 94 85	90 74 85 83 93 94 80 68 8 98 94
HIGH MEAN LOW MEAN EXP MEAN C.V. % LSD 5% LSD 1% # OF REPS		39 0 12 61 11 15 4	98 19 83 11 14 18 4	98 8 80 10 12 16 4

#### Summary

EPTC at 2.5 lb/A, EPTC+cycloate at 2+2 lb/A, EPTC+cycloate+diethatyl at 1.5+1.5+4 lb/A, and EPTC+cycloate/diethatyl (pre) at 1.5+1.5/4 lb/A gave sugarbeet injury over 20%. Only metolachlor at 1.5 lb/A, diethatyl at 4 lb/A, diethatyl (pre) at 4 lb/A, and ethofumesate at 3 lb/A gave less than 90% foxtail spp. control. Only metolachlor at 1.5 lb/A, diethatyl at 4 lb/A, and diethatyl (pre) at 4 lb/A gave less than 80% redroot pigweed control.

Soil applied herbicides, Mooreton, 1988. Preplant incorporated herbicides were applied and rototiller incorporated April 21 when the air temp. was 59F, soil temp. at six inches was 49F, wind was northwest at 8-12 mph, soil was dry at 0-2 inches, and moist at 3-4 inches. The rototiller was operated four inches deep for treatments containing EPTC or cycloate and two inches deep for all other PPI treatments. 'Maribo Ultramono' sugarbeet was seeded 1.25 inches deep in 22 inch rows April 21. Preemergence herbicides were applied April 21. All herbicides were applied in 17 gpa water at 40 psi through 8002 nozzle tips to the center four rows of six row plots. Yellow foxtail control and sugarbeet injury were evaluated June 6.

		Sugarbeet	Yellow Foxtail
Treatment	Rate	injury	control
	(1b/A)	(%	
EPTC	2.5	26	98
Metolachlor	1.5	1	84
Metolachlor	2	13	90
EPTC+Cycloate	1.5+1.5	9	97
Metolachlor	3	14	97
EPTC+Cycloate	2+2	26	98
Cycloate	4	0	91
Diethatyl	4	3	63
Diethatyl (pre)	4	0	0
EPTC+Cycloate+Diethatyl		38	99
EPTC+Cycl/Diet (pre)		31	98
Ethofumesate	3	0	51
HIGH MEAN		38	99
LOW MEAN		0	0
EXP MEAN		13	80
C.V. %		33	7
LSD 5%		6	8
LSD 1%		8	11
# OF REPS		4	4

## Summary

EPTC+cycloate+diethatyl at 1.5+1.5+4 caused greater sugarbeet injury than all other treatments. EPTC at  $2.5\,$  lb/A, EPTC+cycloate at  $2+2\,$  lb/A, and EPTC+cycloate/diethatyl (pre) caused greater sugarbeet injury than all other treatments except EPTC+cycloate+diethatyl. All treatments except metolachlor at  $1.5\,$  lb/A, diethatyl at 4 lb/A, diethatyl (pre) at 4 lb/A, and ethofumesate at 3 lb/A gave 90% or greater control of yellow foxtail.

Soil applied herbicides, Raymond, 1988. Preplant incorporated herbicides were applied and rototiller incorporated April 20 when the air temp. was 59F, soil temp. at six inches was 54F, wind was northwest at 8-12 mph, soil was dry at 0-1 inch, moist at 1-3 inches, and wet at 3-6 inches. The rototiller was operated four inches deep for treatments containing EPTC or cycloate and two inches deep for all other PPI treatments. 'Maribo Ultramono' sugarbeet was seeded 1.25 inches deep in 22 inch rows April 20. Preemergence herbicides were applied April 20. All herbicides were applied in 17 gpa water at 40 psi through 8002 nozzle tips to the center four rows of six row plots. Green and yellow foxtail control and sugarbeet injury were evaluated May 23.

Treatment	Rate (1b/A)	Sugarbeet injury (%	Gr & Ye Foxtail control	
EPTC	2.5	0	76	
Metolachlor	1.5	0	30	
Metolachlor	2	0	45	
EPTC+Cycloate	1.5+1.5	0	71	
Metolachlor	3	0	70	
EPTC+Cycloate	2+2	0	76	
Cycloate	4	0	50	
Diethatyl	4	0	14	
Diethatyl (pre)	4	0	0	
EPTC+Cycloate+Diethatyl		0	83	
EPTC+Cycloate/Diet (pre)		0	89	
Ethofumesate	3	0	33	
HIGH MEAN		0	89	
LOW MEAN		0	0	
EXP MEAN		0	53	
C.V. %		0	21	
LSD 5%		NS	16	
LSD 1%		NS	22	
# OF REPS		4	4	

## Summary

None of the treatments caused sugarbeet injury. The control of foxtail spp. was less than expected from the treatments. However, the organic matter at this location was 9.7% and the weed density was high. This may explain the lack of efficacy by the herbicides.

Soil applied herbicides, Robbin, 1988. Preplant incorporated herbicides were applied and rototiller incorporated May 3 when the air temp. was 72F, soil temp. at six inches was 53F, wind was north at 10-15 mph, soil was dry at 0-2 inches and moist at 2-4 inches. The rototiller was operated four inches deep for treatments containing EPTC or cycloate and two inches deep for all other PPI treatments. 'Interstate Puressa II' sugarbeet was seeded 1.25 inches deep in 22 inch rows May 3. Preemergence herbicides were applied May 3. All herbicides were applied in 17 gpa water at 40 psi through 8002 nozzle tips to the center four rows of six row plots. Redroot pigweed and green foxtail control were evaluated June 25.

<u>Treatment</u> Rate	Redroot Pigweed control	Green Foxtail
(1b/A)	(%)	control
EPTC  Metolachlor  Metolachlor  EPTC+Cycloate  Metolachlor  EPTC+Cycloate  Metolachlor  SEPTC+Cycloate  Upercomparison of the process of the	59 74 83 59 93 71 35 79 0 95 88 78	98 80 85 96 89 99 88 71 26 97
HIGH MEAN LOW MEAN EXP MEAN C.V. % LSD 5% LSD 1% # OF REPS	95 0 68 11 10 14 4	99 26 84 9 10 14 4

## Summary

Only metolachlor at 3 lb/A and EPTC+cycloate+diethatyl at 1.5+1.5+4 lb/A gave over 90% redroot pigweed control. Only metolachlor at 1.5 and 2 lb/A, diethatyl at 4 lb/A, diethatyl (pre) at 4 lb/A, and ethofumesate at 3 lb/A gave green foxtail control less than 86%.

Soil applied plus postemergence herbicides, Bathgate, 1988. Preplant incorporated herbicides were applied and rototiller incorporated four inches deep May 3. 'Interstate Puressa II' sugarbeet was seeded 1.25 inches deep in 22 inch rows May 3. Preemergence herbicides were applied May 3. All soil applied herbicides were applied in 17 gpa water at 40 psi through 8002 nozzle tips to the center four rows of six row plots. Postemergence treatments containing desmedipham at 0.25 lb/A and desmedipham at 0.33 lb/A were applied May 31 and June 7 respectively. All treatments containing clopyralid were applied June 7. The final portion of treatments with three postemergence applications was applied June 13.

		Date o	f Application	
	May 3	May 31	June 7	June 13
Time of day		12:45 pm	11:00 am	12:30 pm
Sky		sunny	sunny	mostly sunny
Air temperature (OF)	55	92	93	83
Soil temp. (OF at 6 in)	53	71	74	70
Relative Humidity (%)		46	38	37
Wind Velocity (mph)	8-12	20	8-12	10
Wind Direction	north	south	southeast	north
Dry Soil (inches)	0-2	0-2	0-3	0-3
Moist Soil (inches)	2-4	2-4	3-5	3-5
Wet Soil (inches)	below 4	below 4	below 5	below 5
Sugarbeets		cotyl-2 lf	4-8 1f	6-8 lf
Redroot pigweed		cotyl-2 lf	2-6 1f	6 1f-3" tall

Postemergence herbicides were applied in 8.5 gpa water at 40 psi through 8001 nozzle tips to the center four rows of six row plots. Sugarbeet injury and redroot pigweed control were evaluated June 25.

Treatment*	Rate	Sugarbeet injury	Redroot Pigweed control
	(1b/A)	(%)	
Desmedipham/Desmedipham Desmed/Desmed/Desmed Desm/Desm+Sethoxy/Desm Desm/Desm+Flua-P/Desm Des/Des/Des+Seth+Dash Des/Des/Des+Flua-P+OC De/De+Clpy/De+Seth+Dash Des/Des/Sethoxydim+Dash Des/Des/Fluazifop-P+OC Desm+Endothall/Desm+Endotha	0.25/0.33 0.25/0.33/0.5 0.25/0.33+0.2/0.5 0.25/0.33+0.19/0.5 0.25/0.33/0.5+0.2+0.25G 0.25/0.33/0.5+0.19+0.25G 0.25/0.33+0.09/0.5+0.2+0.25G 0.25/0.33/0.2+0.25G 0.25/0.33/0.2+0.25G	9 18 15 21 13 15 20 11 10 6	83 98 97 94 98 97 98 83 84 58
Clopyralid+Endothall Clopyralid+Endothall Clopyralid+Endothall	0.09+0.5 0.19+0.5 0.09+0.75	0 0 0	0 0 0

(continued on next page)

Soil applied plus postemergence herbicides, Bathgate, 1988. (continued)

			Redroot
Tuestant		Sugarbeet	Pigweed
Treatment*	Rate	injury	control
	(1b/A)	(%)	
Clopyralid+Seth+Dash	0.09+0.2+0.25G		
Clopyralid+Seth+Dash		0	0
Clopyralid+Seth+Dash	0.14+0.2+0.25G	0	4
Clopyralid+Sethoxydim	0.19+0.2+0.25G	0	13
Clopyralid+Sethoxydim	0.09+0.2	0	0
Clopyralid+Sethoxydim	0.14+0.2	0	0 3
EPTC+Cycl(ppi)/Desmedipham	0.19+0.2	0	
Diethatyl(pre)/Desmedipham	1.5+1.5/0.33	16	74
FDTC+Cycl(nni)/Desmedipnam	4/0.33	0	38
EPTC+Cycl(ppi)/Desm/Desm	1.5+1.5/0.25/0.33	22	96
Diet(pre)/Desm/Desm	4/0.25/0.33	5	85
EPTC+Cyc(ppi)/Desm/Desm/Desm		31	99
Diet(pre)/Desm/Desm/Desm	4/0.25/0.33/0.5	21	98
EPTC+Cyc(ppi)/Des/Des+Clpy/De	s+Seth+Dash	38	99
1.5+1.5/0.25	/0.33+0.09/0.5+0.2+0.25G		
Diet(pre)/Des/Des+Clpy/Des+Se		20	98
4/0.25	/0.33+0.09/0.5+0.2+0.25G		
HIGH MEAN			
LOW MEAN		38	99
EXP MEAN		0	0
C.V. %		11	58
LSD 5%		35	6
LSD 1%		5	5 7
# OF REPS		7	
TO KEPS		4	4

<sup>\*</sup> OC = Agway Booster Plus E

Two applications of desmedipham over EPTC+cycloate gave greater sugarbeet injury and redroot pigweed control than desmedipham alone.

<sup>\*</sup> Dash = BASF surfactant

Soil applied plus postemergence herbicides, Clara City, 1988. Preplant incorporated herbicides were applied and rototiller incorporated four inches deep April 19. 'KW 3265' sugarbeet was seeded in 22 inch rows April 20. Preemergence herbicides were applied April 19. All soil applied herbicides were applied in 17 gpa water at 40 psi through 8002 nozzle tips to the center four rows of six row plots. Postemergence treatments containing desmedipham at 0.25 lb/A and desmedipham at 0.33 lb/A were applied May 12 and May 18 respectively. All treatments containing clopyralid were applied May 18. The final portion of treatments with three postemergence applications was applied June 2.

		Date of	Application	
	April 19		May 18	June 2
Time of day		12:15 pm	12:00 pm	11:00 am
Sky		sunny	em 100 em	
Air temperature (OF)	63	90	71	90
Soil temp. (OF at 6 in)	49	65	60 (	at 4")
Relative Humidity (%)		27	24	34
Wind Velocity (mph)	8-12	15	10-12	0
Wind Direction	north	west	southeast	
Dry Soil (inches)	0-1	0-1	ess etc ose	(C) (C) (C)
Moist Soil (inches)	1-3	1-4	em (m) em	an on m
Wet Soil (inches)	below 3	below 4	■ • •	
Sugarbeet		cotyl-2 lf	2-4 1f	6-8 lf
Green and Yellow Foxtai	]	2-3 leaf	1.5" tall	6" tall
		(1" tall)		
Redroot Pigweed	100 COS COS		2-6 leaf	4 to 8" tall
Common Lambsquarters	900 COM 800	2-41f(1" tall)	6 leaf	6 to 10" tall

Postemergence herbicides were applied in 8.5 gpa water at 40 psi through 8001 nozzle tips to the center four rows of six row plots. Redroot pigweed, common lambsquarters, and green and yellow foxtail control and sugarbeet injury were evaluated June 17.

			Grft		
		Sgbt	Yeft	Rrpw	Colq
Treatment*	Rate	inj	cnt1	cnt1	<u>cntl</u>
	(1b/A)		(	%)	
Desmedipham/Desmedipham	0.25/0.33	13	8	54	91
Desmed/Desmed	0.25/0.33/0.5	13	20	88	98
Desm/Desm+Sethoxy/Desm	0.25/0.33+0.2/0.5	26	98	78	95
Desm/Desm+Flua-P/Desm	0.25/0.33+0.19/0.5	19	91	87	97
Des/Des/Des+Seth+Dash	0.25/0.33/0.5+0.2+0.25G	25	96	89	98
Des/Des/Des+Flua-P+OC	0.25/0.33/0.5+0.19+0.25G	15	63	92	97
De/De+Clpy/De+Se+Dash C	0.25/0.33+0.09/0.5+0.2+0.25G	19	93	96	99
Des/Des/Sethoxydim+Dash	0.25/0.33/0.2+0.25G	8	98	56	96
Des/Des/Fluazifop-P+OC	0.25/0.33/0.2+0.25G	6	97	51	92
Desmed+Endo/Desmed+Endo	0.25+0.25/0.33+0.25	5	0	61	75
Clopyralid+Endothall	0.09+0.5	0	0	5	26
Clopyralid+Endothall	0.19+0.5	1	0	20	64
Clopyralid+Endothall	0.09+0.75	0	0	5	29

Soil applied plus postemergence herbicides, Clara City, 1988. (continued)

			061		
		Sah+	Grft	Dwn	Colo
Treatment*	Rate	Sgbt inj	Yeft cntl	Rrpw	Colq
11 Ca chiere		1111		<u>cntl</u>	cntl
	(1b/A)		(	%)	
Clopyralid+Sethoxydim+Dash	0.09+0.2+0.25G	4	96	15	58
Clopyralid+Sethoxydim+Dash	0.14+0.2+0.25G	5	98	18	54
Clopyralid+Sethoxydim+Dash	0.19+0.2+0.25G	8	98	34	73
Clopyralid+Sethoxydim	0.09+0.2	0	96	10	27
Clopyralid+Sethoxydim	0.14+0.2	0	82	15	45
Clopyralid+Sethoxydim	0.19+0.2	5	93	9	56
EPTC+Cyc1(ppi)/Desmedipham	1.5+1.5/0.33	16	76	59	95
Diethatyl(pre)/Desmedipham	4/0.33	14	36	85	81
<pre>EPTC+Cycl(ppi)/Des/Des</pre>	1.5+1.5/0.25/0.33	30	84	69	94
Diet(pre)/Desm/Desm	4/0.25/0.33	8	46	78	94
<pre>EP+Cy(ppi)/Desm/Desm/Desm</pre>	1.5+1.5/0.25/0.33/0.5	31	87	92	99
Diet(pre)/Desm/Desm/Desm	4/0.25/0.33/0.5	15	61	96	95
EP+Cy(ppi)/Des/Des+Clpy/Des+Se					
	.33+0.09/0.5+0.2+0.25G	34	99	96	98
Diet(pre)/Des/Des+Clpy/Des+Se	th+Dash				
4/0.25/0	.33+0.09/0.5+0.2+0.25G	28	98	97	99
HIGH MEAN		34	99	97	99
LOW MEAN		0	0	5	26
EXP MEAN		13	66	57	78
C.V. %		53	16	18	12
LSD 5%		9	15	15	13
LSD 1%		13	20	19	17
# OF REPS		4	4	4	4
			7	7	7

<sup>\*</sup> OC = BASF Booster Plus E

Desmedipham applied two or three times over EPTC+cycloate caused more sugarbeet injury than desmedipham alone. Desmedipham+endothall applied twice gave less control of common lambsquarters than desmedipham applied twice. Clopyralid+sethoxydim+Dash gave common lambsquarters control greater than clopyralid+sethoxydim.

<sup>\*</sup> Dash = BASF surfactant

Soil applied plus postemergence herbicides, Crookston, 1988. Preplant incorporated herbicides were applied and rototiller incorporated four inches deep April 26. 'Maribo Ultramono' sugarbeet was seeded 1.25 inches deep in 22 inch rows April 26. Preemergence herbicides were applied April 26. All soil applied herbicides were applied in 17 gpa water at 40 psi through 8002 nozzle tips to the center four rows of six row plots. Postemergence treatments containing desmedipham at 0.25 lb/A and desmedipham at 0.33 lb/A were applied May 23 and May 27 respectively. All single application postemergence treatments were applied May 27. The final portion of treatments with three postemergence applications was applied June 6.

		Date of	Application	
	April 26	May 23	May 27	June 6
Time of day		11:30 am	5:00 pm	12:45 pm
Sky		sunny	overcast	sunny
Air temperature (OF)	52	82	79	90
Soil temp. (OF at 6 in)	40	60	74	78
	-10	26	26	34
Relative Humidity (%)	5-10	5-10	2-4	15
Wind Speed	west	east	north	south
Wind Direction	0-1	0-1	0-2	0-2
Dry Soil (inches)	1-3	1-4	2-4	2-4
Moist Soil (inches)		below 4	below 4	below 4
Wet Soil (inches)	below 3		2-4 1f	6-10 lf
Sugarbeet	ann ann ann	cotyl-2 lf		1-6" tall
Common Lambsquarters	COD 6234 CEO	cotyl-6 lf		
Redroot pigweed		cotyl	cotyl-2 lf	i through 8001

Postemergence herbicides were applied in 8.5 gpa water at 40 psi through 8001 nozzle tips to the center four rows of six row plots. Common lambsquarters and redroot pigweed control and sugarbeet injury were evaluated June 13.

		Sgbt	Colq	Rrpw
Treatment*	Rate	inj	cntl	cnt1
Treatment	(1b/A)		(%) -	
	•			= 0
Desmedipham	0.33	0	73	70
Desmedipham/Desmedipham	0.25/0.33	8	98	94
Desmed/Desmed	0.25/0.33/0.5	9	98	98
Des&Phen/Des&Phen/Des&Phen	0.25/0.33/0.5	22	99	99
De/De+Clpy/De+Seth+Dash 0.25/0	.33+0.14/0.5+0.2+0.25G	26	99	99
Desmed/Desmed+Fluazifop-P	0.25/0.33+0.19	11	95	94
Desmed/Desmed+Sethoxydim	0.25/0.33+0.2	9	94	93
Fluazifop-P+OC	0.19+0.25G	0	0	0
Sethoxydim+Dash	0.2+0.25G	0	0	0
Clopyralid	0.09	0	0	0
Clopyralid	0.14	0	4	0
	0.19	0	10	0
Clopyralid Clopyralid+Endothall	0.09+0.5	0	0	0
Clopyralid+Endothall	0.19+0.5	0	5	0
	0.09+0.75	0	6	6
Clopyralid+Endothall	0.09+0.2+0.25G	0	1	0
Clopyralid+Sethoxydim+OC	0.14+0.2+0.25G	0	53	9
Clopyralid+Sethoxydim+OC	0.19+0.2+0.25G	Ö	54	8
Clopyralid+Sethoxydim+OC	0.13.0.2.0.230			

Soil applied plus postemergence herbicides, Crookston, 1988. (continued)

		Sgbt	Colq	Rrpw
Treatment*	Rate	inj	cntl	cntl
	(1b/A)		(%) -	
Desmed/Desmed+Clopyralid	0.25/0.33+0.09	5	97	91
Desmed/Desmed+Clopyralid	0.25/0.33+0.19	0	97	92
<pre>Desmed/Desmed+Clopyral</pre>		15	98	99
Desm+Endothall/Desm+Endothall	0.25+0.25/0.33+0.25	0	93	85
EPTC+Cycloate(ppi)/Desmedipha		43	99	99
<pre>EPTC+Cycloate(ppi)/Desm/Desm</pre>	1.5+2/0.25/0.33	41	99	99
<pre>EPTC+Cycloate(ppi)/Des/Des/De</pre>	s 1.5+2/0.25/0.33/0.5	54	99	99
Diethatyl(pre)/Desmedipham	4/0.33	0	68	85
Diethatyl(pre)/Desm/Desm	4/0.25/0.33	5	95	96
<pre>Diethatyl(pre)/Desm/Desm/Desm</pre>		24	99	99
<pre>Diet(pre)/Desm/Desm+Clpy/Desm</pre>				
4/0.25	/0.33+0.14/0.5+0.2+0.25G	25	98	99
HIGH MEAN		54	99	99
LOW MEAN		0	0	0
EXP MEAN		10	63	59
C.V. %		56	10	11
LSD 5%		8	9	9
LSD 1%		11	11	12
# OF REPS		4	4	4
				7

<sup>\*</sup> OC = Agway Booster Plus E

Three applications of desmedipham+phenmedipham caused greater sugarbeet injury than three applications of desmedipham. Sugarbeets treated with EPTC+cycloate and desmedipham were injured more than sugarbeets only treated with desmedipham. Also, diethatyl followed by desm/desm/desm caused greater sugarbeet injury than desm/desm/desm. Two applications of desmedipham gave common lambsquarters and redroot pigweed control superior to one application of desmedipham but similar to three applications of desmedipham. Clopyralid at 0.19 lb/A gave 10% control of common lambsquarters while clopyralid+sethoxydim+Booster Plus E at 0.19+0.2 lb/A + 1 qt/A gave 54% control.

<sup>\*</sup> Dash = BASF surfactant

Soil applied plus postemergence herbicides, Hillsboro, 1988. Preplant incorporated herbicides were applied and rototiller incorporated four inches deep April 25. 'Maribo Ultramono' sugarbeet was seeded 1.25 inches deep in 22 inch rows April 25. Preemergence herbicides were applied April 25. All soil applied herbicides were applied in 17 gpa water at 40 psi through 8002 nozzle tips to the center four rows of six row plots. Postemergence treatments containing desmedipham at 0.25 lb/A and desmedipham at 0.33 lb/A were applied May 23 and May 27 respectively. All treatments containing clopyralid were applied May 27. The final portion of treatments with three postemergence applications was applied June 16.

	Date of Application				
	April 25	May 23	May 27	June 16	
Time of day		2:00 pm	11:45 am	11:00 am	
Sky	au ca ea	sunny	sunny	sunny	
Air temperature (OF)	48	84	84	88	
Soil temp. (OF at 6 in)	43	63	66	74	
Relative Humidity (%)		30	47	35	
Wind Velocity (mph)	20	8-12	0-5	15	
Wind Direction	north	north	east	south	
Dry Soil (inches)	0-1	0-1	0-1	0-2	
Moist Soil (inches)	1-3	1-3	1-4	2-3	
Wet Soil (inches)	below 3	below 3	below 4	below 3	
Sugarbeet	ecs Chir can	coty1-4 1f	cotyl-4 lf	4-10 lf	
Green Foxtail	COS COST 1980	2 lf(1" tall)-	emerging to	5 lf(1.5"	
		4 1f(2" tall)	5 lf(3" tall)	tall)-5" tall	
Redroot pigweed		2-4 1f	cotyl-6 lf	4 1f-2.5"tall	

Postemergence herbicides were applied in 8.5 gpa water at 40 psi through 8001 nozzle tips to the center four rows of six row plots. Green foxtail and redroot pigweed control and sugarbeet injury were evaluated June 20.

		Sgbt	Grft	Rrpw
Treatment*	Rate	inj	cntl	cntl
	(1b/A)		(%)	ME ON ME CON COL ME
Desmedipham/Desmedipham	0.25/0.33	0	3	59
Desmed/Desmed	0.25/0.33/0.5	25	40	90
Desm/Desm+Sethoxydim/Desm	0.25/0.33+0.2/0.5	33	76	87
Desm/Desm+Flua-P/Desm	0.25/0.33+0.19/0.5	29	60	90
Des/Des/Des+Seth+Dash	0.25/0.33/0.5+0.2+0.25G	34	92	80
Des/Des/Des+Flua-P+OC 0	.25/0.33/0.5+0.19+0.25G	31	64	81
De/De+Clpy/De+Seth+Dash 0.25/0	0.33+0.09/0.5+0.2+0.25G	43	87	81
Des/Des/Sethoxydim+Dash	0.25/0.33/0.2+0.25G	5	98	52
Des/Des/Fluazifop-P+OC	0.25/0.33/0.2+0.25G	5	79	56
Des+Endothall/Des+Endothall	0.25+0.25/0.33+0.25	0	11	26
Clopyralid+Endothall	0.09+0.5	0	15	0
Clopyralid+Endothall	0.19+0.5	0	15	3
Clopyralid+Endothall	0.09+0.75	0	34	0

Soil applied plus postemergence herbicides, Hillsboro, 1988. (continued)

		C 1 1	0.51	
T		Sgbt	Grft	Rrpw
Treatment*	Rate	inj	cntl	cntl
	(1b/A)		(%)	
Clopyralid+Sethoxydim+Dash	0.09+0.2+0.25G	0	89	0
Clopyralid+Sethoxydim+Dash	0.14+0.2+0.25G	0	91	0
Clopyralid+Sethoxydim+Dash	0.19+0.2+0.25G	3	91	0
Clopyralid+Sethoxydim	0.09+0.2	0	86	0
Clopyralid+Sethoxydim	0.14+0.2	0	88	0
Clopyralid+Sethoxydim	0.19+0.2	0	85	0
<pre>EPTC+Cycl(ppi)/Desmedipham</pre>	1.5+1.5/0.33	45	92	91
Diethatyl(pre)/Desmedipham	4/0.33	0	40	68
<pre>EPTC+Cycl(ppi)/Desm/Desm</pre>	1.5+1.5/0.25/0.33	49	96	95
Diet(pre)/Desm/Desm	4/0.25/0.33	0	54	63
EPTC+Cyc(ppi)/Des/Des/Des	1.5+1.5/0.25/0.33/0.5	66	98	98
Diet(pre)/Desm/Desm/Desm	4/0.25/0.33/0.5	30	59	88
EPTC+Cyc(ppi)/Des/Des+Clpy/Des	s+Seth+Dash	70	99	98
1.5+1.5/0.25/0.	.33+0.09/0.5+0.2+0.25G			
Diet(pre)/Des/Des+Clpy/Des+Set		33	94	87
4/0.25/0	.33+0.09/0.5+0.2+0.25G			
HIGH MEAN		70	99	98
LOW MEAN		0	3	0
EXP MEAN		18	67	51
C.V. %		37	16	14
LSD 5%		10	15	10
LSD 1%		13	20	14
# OF REPS		4	4	4
		MALE SALES		

<sup>\*</sup> OC = Agway Booster Plus E

Postemergence herbicides injury and gave better weed control than postemergence herbicides alone. Postemergence herbicides over preemergence diethatyl gave sugarbeet injury and redroot pigweed control similar to postemergence herbicides alone possibly because rainfall was inadequate to activate the diethatyl. Three applications of desmedipham gave greater sugarbeet injury and redroot pigweed control than two applications.

<sup>\*</sup> Dash = BASF surfactant

Soil applied plus postemergence herbicides, Mooreton, 1988. Preplant incorporated herbicides were applied and rototiller incorporated four inches deep April 21. 'Maribo Ultramono' sugarbeet was seeded 1.25 inches deep in 22 inch rows April 21. Preemergence herbicides were applied April 21. All soil applied herbicides were applied in 17 gpa water at 40 psi through 8002 nozzle tips to the center four rows of six row plots. Postemergence treatments containing desmedipham at 0.25 lb/A and desmedipham at 0.33 lb/A were applied May 13 and May 20 respectively. All treatments containing clopyralid were applied May 20. The final portion of treatments with three postemergence applications was applied June 1.

		D-+£	Annlination	
			Application	
	April 21	May 13	May 20	June 1
Time of day		2:00 pm	3:00 pm	1:00 pm
Sky	000 min min	sunny	partly cloudy	mostly sunny
Air temperature (OF)	59	58	70	88
Soil temp. (OF at 6 in	n) 49	60	64	78
Relative Humidity (%)	gen man min	52	64	55
Wind Velocity (mph)	8-12	3-6	10-15	15
Wind Direction	northwest	southeast	northeast	south
Dry Soil (inches)	0-2	0-2	0-1	0-2
Moist Soil (inches)	2-4	2-6	1-4	2-4
Wet Soil (inches)	below 4	below 6	below 4	below 4
Sugarbeet	an eo eo	cotyl-2 lf	2 1f	6-8 lf
Green Foxtail	ua en en	emerging to		3 to 5 inches
		1" tall(21f)	(1-1.5" tall)	tall
Wild Oats	600 000 600	1 to 2" tall	2 to 4" tall	
Common Lambsquarters		cotyl-2 lf	6-8 leaf	3 to 6" tall

Postemergence herbicides were applied in 8.5 gpa water at 40 psi through 8001 nozzle tips to the center four rows of six row plots. Green foxtail, wild oats, and common lambsquarters control and sugarbeet injury were evaluated June 28.

0		Sgbt	Grft	Wioa	Colq
Treatment*	Rate	inj	cntl	cnt1	cntl
	(1b/A)	aa aa aa aa aa aa	(	%)	
Desmedipham/Desmedipham	0.25/0.33	0	16	0	50
Desmed/Desmed	0.25/0.33/0.5	0	38	0	86
Desm/Desm+Sethoxydim/Desm	0.25/0.33+0.2/0.5	0	79	98	89
Desm/Desm+Flua-P/Desm	0.25/0.33+0.19/0.5	0	74	99	93
Des/Des/Des+Seth+Dash	0.25/0.33/0.5+0.2+0.25G	0	76	66	86
Des/Des/Des+Flua-P+OC	0.25/0.33/0.5+0.19+0.25G	0	54	50	87
Des/Des+Clpy/Des+Seth+Dash	0.25/0.33+0.09/.5+.2+.25G	0	78	76	90
Desm/Desm/Sethoxydim+Dash	0.25/0.33/0.2+0.25G	0	97	96	48
Desm/Desm/Fluazifop-P+OC	0.25/0.33/0.2+0.25G	0	92	99	58
Desm+Endothall/Desm+Endotha		0	2	9	53
Clopyralid+Endothall	0.09+0.5	0	9	0	15
Clopyralid+Endothall	0.19+0.5	0	18	0	60
Clopyralid+Endothall	0.09+0.75	0	18	8	0

Soil applied plus postemergence herbicides, Mooreton, 1988. (continued)

	and the little and Alexander				
		Sgbt	Grft	Wioa	Colq
Treatment*	Rate	inj	cnt1	cntl	cntl
	(1b/A)		(	%)	
Clopyralid+Sethoxydim+Dash	0.09+0.2+0.25G	0	88	99	32
Clopyralid+Sethoxydim+Dash	0.14+0.2+0.25G	0	84	99	49
Clopyralid+Sethoxydim+Dash	0.19+0.2+0.25G	0	84	99	55
Clopyralid+Sethoxydim	0.09+0.2	0	83	93	20
Clopyralid+Sethoxydim	0.14+0.2	0	79	99	46
Clopyralid+Sethoxydim	0.19+0.2	0	76	83	58
EPTC+Cycl(ppi)/Desmedipham	1.5+1.5/0.33	4	95	91	95
Diethatyl(pre)/Desmedipham	4/0.33	0	9	0	11
EPTC+Cycl(ppi)/Desm/Desm	1.5+1.5/0.25/0.33	3	96	97	98
Diet(pre)/Desm/Desm	4/0.25/0.33	0	45	33	53
<pre>EP+Cy(ppi)/Desm/Desm/Desm</pre>	1.5+1.5/0.25/0.33/0.5	10	97	99	99
<pre>Diet(pre)/Desm/Desm/Desm</pre>	4/0.25/0.33/0.5	0	48	23	90
<pre>EP+Cy(ppi)/Desm/Desm+Clpy/Desi</pre>		10	98	99	99
	/0.33+0.09/0.5+0.2+0.25G				
<pre>Diet(pre)/Desm/Desm+Clpy/Desm</pre>		0	79	66	91
4/0.25	/0.33+0.09/0.5+0.2+0.25G				
HIGH MEAN		10	98	99	99
LOW MEAN		0	9	0	0
EXP MEAN		1	63	62	63
C.V. %		386	12	17	19
LSD 5%		5	11	15	17
LSD 1%		7	15	19	23
# OF REPS		4	4	4	4

<sup>\*</sup> OC = Agway Booster Plus E \* Dash = BASF surfactant

Sethoxydim and fluazifop-P gave similar control of green foxtail and wild oats when not tank mixed with desmedipham. However, green foxtail and wild oats control from fluazifop-P was antagonized more than sethoxydim when tank mixed with desmedipham.

Soil applied plus postemergence herbicides, Raymond, 1988. Preplant incorporated herbicides were applied and rototiller incorporated four inches deep April 20. 'Maribo Ultramono' sugarbeet was seeded 1.25 inches deep in 22 inch rows April 20. Preemergence herbicides were applied April 20. All soil applied herbicides were applied in 17 gpa water at 40 psi through 8002 nozzle tips to the center four rows of six row plots. Postemergence treatments containing desmedipham at 0.25 lb/A and desmedipham at 0.33 lb/A were applied May 12 and May 18 respectively. All treatments containing clopyralid were applied May 18. The final portion of treatments with three postemergence applications was applied June 2.

		Date of A	Application	
	April 20	May 12		June 2
Time of day		1:30 pm	8:30 am	10:00 am
Air temperature (OF)	59	90	54	85
Soil temp. (OF at 6 in		65	55(at	4 in) 74
Relative Humidity (%)		27	48	40
Wind Velocity (mph)	8-12	20	0-8	0
	northwest	west	southeast	
Soil Moisture	good		fair	poor
Sugarbeet		cotyl-2 lf	2-4 1f	6-8 lf
Common Lambsquarters	000 000 <b>000</b>	4 lf(1" tall)	4-6 1f	8-10" tall
Redroot pigweed		2 1f	4 1f	8-10" tall
		cotyl-2 lf	2-4 lf	6-9" tall
Marshelder Green and Yellow Foxta		2-31f(.5-1"tall)		6-8" tall
Green and Terrow Toxca	11	2011	water at 40 pci	through 8001

Postemergence herbicides were applied in 8.5 gpa water at 40 psi through 8001 nozzle tips to the center four rows of six row plots. Common lambsquarters, redroot pigweed, marshelder, and green and yellow foxtail control and sugarbeet injury were evaluated June 17.

		0 1 1	0 7	D	M 7	F., + 1
					Mael	
Treatment*	Rate	inj	cnt1	cntl	cntl	cntl
T CA CHIEFT C	(1b/A)			- (%)		
Desmedipham/Desmedipham	0.25/0.33	0	92	88	94	10
Desmed/Desmed/Desmed	0.25/0.33/0.5	0	95	96	99	30
Desm/Desm+Sethoxydim/Desm	0.25/0.33+0.2/0.5	0	96	77	98	74
Desm/Desm+Flua-P/Desm	0.25/0.33+0.19/0.5	4	96	65	99	48
Des/Des/Des+Seth+Dash	0.25/0.33/0.5+0.2+0.25G	8	97	90	98	84
Des/Des/Des+Flua-P+OC	0.25/0.33/0.5+0.19+0.25G	0	98	92	97	64
	/0.33+0.09/0.5+0.2+0.25G	11	97	89	99	94
Des/Des/Sethoxydim+Dash	0.25/0.33/0.2+0.25G	0	92	59	97	95
Des/Des/Fluazifop-P+OC	0.25/0.33/0.2+0.25G	0	60	61	70	64
Desmed+Endothall/Desm+Endo	0.25+0.25/0.33+0.25	0	33	77	96	0
Clopyralid+Endothall	0.09+0.5	0	0	0	99	0
Clopyralid+Endothall	0.19+0.5	0	8	0	99	0
Clopyralid+Endothall	0.09+0.75	0	3	0	99	8

Soil applied plus postemergence herbicides, Raymond, 1988. (continued)

		Sgbt	Colq	Rrpw	Mael	Fxt1
Treatment*	Rate		cntl			
	(1b/A)			- (%)		
Clopyralid+Sethoxydim+Dash	0.09+0.2+0.25G	0	10	0	99	95
Clopyralid+Sethoxydim+Dash	0.14+0.2+0.25G	0	49	0	99	96
Clopyralid+Sethoxydim+Dash	0.19+0.2+0.25G	0	51	0	99	89
Clopyralid+Sethoxydim	0.09+0.2	0	0	0	99	87
Clopyralid+Sethoxydim	0.14+0.2	0	8	0	99	92
Clopyralid+Sethoxydim	0.19+0.2	0	50	4	99	71
EPTC+Cycloate(ppi)/Desmedipha	m 1.5+1.5/0.33	6	95	50	99	56
Diethatyl(pre)/Desmedipham	4/0.33	0	68	83	82	8
EPTC+Cycloate(ppi)/Desm/Desm	1.5+1.5/0.25/0.33	3	97	43	99	66
Diethatyl(pre)/Desm/Desm	4/0.25/0.33	0	90	75	98	39
EPTC+Cycl(ppi)/Des/Des/Des	1.5+1.5/0.25/0.33/0.5	16	98	73	99	73
Diethatyl(pre)/Des/Des/Des	4/0.25/0.33/0.5	0	97	86	97	21
<pre>EPTC+Cycl(ppi)/Des/Des+Clpy/D</pre>						
1.5+1.5/0.25/0	.33+0.09/0.5+0.2+0.25G	21	99	93	99	96
Diethatyl(pre)/Des/Des+Clpy/D						
4/0.25/0	.33+0.09/0.5+0.2+0.25G	3	96	93	99	88
HIGH MEAN		21	99	96	99	96
LOW MEAN		0	0	0	70	0
EXP MEAN		3	66	51	97	57
C.V. %		229	11	22	5	23
LSD 5%		8	10	16	7	19
LSD 1%		11	13	21	9	25
# OF REPS		4	4	4	4	4

<sup>\*</sup> OC = Agway Booster Plus E

Two applications of desmedipham followed by fluazifop-P+oil concentrate gave less control of broadleaf weeds than two applications of desmedipham. Two applications of desmedipham followed by sethoxydim+Dash gave less control of redroot pigweed than two applications of desmedipham. Desmedipham + endothall applied twice gave less control of common lambsquarters than desmedipham applied twice. All treatments including clopyralid gave essentially complete control of marshelder.

<sup>\*</sup> Dash = BASF surfactant

Soil applied plus postemergence herbicides, Robbin, 1988. Preplant incorporated herbicides were applied and rototiller incorporated four inches deep May 3. 'Interstate Puressa II' sugarbeet was seeded 1.25 inches deep in 22 inch rows May 3. Preemergence herbicides were applied May 3. All soil applied herbicides were applied in 17 gpa water at 40 psi through 8002 nozzle tips to the center four rows of six row plots. Postemergence treatments containing desmedipham at 0.25 lb/A and desmedipham at 0.33 lb/A were applied June 7 and June 13 respectively. All treatments containing clopyralid were applied June 13. The final portion of treatments with three postemergence applications was applied June 20.

		Date	of Application	
	May 3	June 7	June 13	June 20
Time of day		8:30 am	3:00 pm	1:00 pm
Sky	gua aus aus	sunny	mostly sunny	sunny
Air temperature (OF)	72	81	88	92
Soil temp. (OF at 6 in)	53	73	76	78
Relative Humidity (%)		48	33	26
Wind Velocity (mph)	10-15	5-8	5-10	10-15
Wind Direction	north	south	north	south
Dry Soil (inches)	0-2	0-2	0-2	0-2
Moist Soil (inches)	2-4	2-5	2-5	2-6
Wet Soil (inches)	below 4	below 5	below 5	below 6
Green Foxtail		emerging to	3 lf(.5"	1-4" tall
		3 lf(1" tall)	tall)-2" tall	
Redroot pigweed		cotyl-4 lf	4-8 1f	1-4" tall

Postemergence herbicides were applied in 8.5 gpa water at 40 psi through 8001 nozzle tips to the center four rows of six row plots. Green foxtail and redroot pigweed control were evaluated June 29.

		Redroot Pigweed	Green Foxtail
Treatment*	Rate	control	control
	(1b/A)		(%)
	0.05/0.00	01	46
Desmedipham/Desmedipham	0.25/0.33	81	
Desmed/Desmed	0.25/0.33/0.5	89	48
Desm/Desm+Sethoxydim/Desm	0.25/0.33+0.2/0.5	92	97
Desm/Desm+Flua-P/Desm	0.25/0.33+0.19/0.5	91	69
Des/Des/Des+Seth+Dash	0.25/0.33/0.5+0.2+0.25G	94	97
	0.25/0.33/0.5+0.19+0.25G	94	75
Des/Des/Des+Flua-P+OC		91	96
20,20	.25/0.33+0.09/0.5+0.2+0.25G		
Des/Des/Sethoxydim+Dash	0.25/0.33/0.2+0.25G	74	95
Des/Des/Fluazifop-P+OC	0.25/0.33/0.2+0.25G	83	64
Desm+Endothall/Desm+Endotha	all 0.25+0.25/0.33+0.25	68	56
Clopyralid+Endothall	0.09+0.5	3	40
	0.19+0.5	Ō	50
Clopyralid+Endothall		ő	45
Clopyralid+Endothall	0.09+0.75	0	45

Soil applied plus postemergence herbicides, Robbin, 1988. (continued)

		Redroot	Green
Treatment*		Pigweed	Foxtail
11 ea chieff C	Rate	control	control
	(1b/A)		%)
Clanusalidicalian			,,,
Clopyralid+Seth+Dash	0.09+0.2+0.25G	0	99
Clopyralid+Seth+Dash	0.14+0.2+0.25G	Ō	99
Clopyralid+Seth+Dash	0.19+0.2+0.25G	Ö	99
Clopyralid+Sethoxydim	0.09+0.2	ŏ	98
Clopyralid+Sethoxydim	0.14+0.2	ő	98
Clopyralid+Sethoxydim	0.19+0.2	ő	98
EPTC+Cycl(ppi)/Desmedipham	1.5+1.5/0.33	78	93
Diethatyl(pre)/Desmedipham	4/0.33	64	
EPTC+Cyc1(ppi)/Desm/Desm	1.5+1.5/0.25/0.33	95	58
Diet(pre)/Desm/Desm	1/0 25/0 22	93	96
EPTC+Cyc(ppi)/Desm/Desm/Desm	1.5+1.5/0.25/0.33/0.5	98	77
Diec(bre)/Desm/Desm/Desm	4/0 25/0 33/0 5		99
EPTC+Cyc(ppi)/Des/Des+Clpy/De	s+Seth+Dash	95	82
1,5+1,5/0,25	/0.33+0.09/0.5+0.2+0.25G	98	99
Diet(pre)/Des/Des+Clpy/Des+Se	th+Dash	00	
	/0.33+0.09/0.5+0.2+0.25G	93	96
1, 0.25	70.5510.0370.540.240.256		
HIGH MEAN			
LOW MEAN		98	99
EXP MEAN		0	40
C.V. %		58	79
LSD 5%		7	9
LSD 1%		6	10
# OF REPS		8	13
" OI MEI'O		4	4
4 00			

<sup>\*</sup> OC = Agway Booster Plus E \* Dash = BASF surfactant

Clopyralid did not antagonize green foxtail control from sethoxydim. Clopyralid+sethoxydim with no oil additive gave excellent green foxtail control. Sethoxydim gave greater control of green foxtail than fluazifop-P. EPTC+cycloate or preemergence diethatyl followed by two or three desmedipham applications gave greater redroot pigweed control than desmedipham alone.

Common cocklebur control, Raymond, 1988. 'KW 3394' sugarbeet was seeded in 22 inch rows May 19. The first half of split application treatments was applied 9:00 am June 9 when the air temp. was 72F, soil temp. at four inches was 65F, relative humidity was 24%, wind was southeast at 10-12 mph, soil was dry, sugarbeets were 4-6 leaf, and cocklebur were 4-6 leaf. The second half of split treatments and all single application treatments were applied 9:00 am June 14 when the air temp. was 69F, soil temp. at four inches was 62F, relative humidity was 32%, wind was northwest at 8-10 mph, soil was very dry, sugarbeets were 6-10 leaf, and cocklebur were 8-10 leaf. All treatments were applied in 8.5 gpa water at 40 psi through 8001 nozzle tips. Sugarbeet injury and common cocklebur control were evaluated June 17.

		Sugarbeet	Common Cocklebur
Treatment*	Rate	injury	control
	(1b/A)	(%	,)
Clanymalid	0.06	0	96
Clopyralid	0.125	0	96
Clopyralid	0.19	ő	96
Clopyralid	0.06+0.25G		96
Clopyralid+OC		3	96
Clopyralid+OC	0.125+0.25G	0 3 3	96
Clopyralid+OC	0.19+0.25G	25	98
Des/Des+Clopyralid	0.33/0.5+0.06		98
Des/Des+Clopyralid	0.33/0.5+0.125	26	98
Des/Des+Clopyralid	0.33/0.5+0.19	28	98
De&Ph/De&Ph+Clopyralid	0.33/0.5+0.06	27	
De&Ph/De&Ph+Clopyralid	0.33/0.5+0.125	28	98
De&Ph/De&Ph+Clopyralid	0.33/0.5+0.19	27	98
Endothall+Clopyralid	0.5+0.06	11	97
Endothall+Clopyralid	0.5+0.125	8	97
Endothall+Clopyralid	0.5+0.19	10	97
Endothall+Clopyralid	0.75+0.125	16	98
HICH MEAN		28	98
HIGH MEAN		0	96
LOW MEAN		13	97
EXP MEAN		37	1
C.V. %		7	1
LSD 5%			1
LSD 1%		9 4	4
# OF REPS		4	

<sup>\*</sup> OC = Agway Booster Plus E

# Summary

All treatments gave excellent control of common cocklebur. Desmedipham or desmedipham&phenmedipham caused greater sugarbeet injury than other treatments.

Effect of herbicides on sugarbeet root rot, Hector, 1988. incorporated herbicides were applied in 17 gpa water at 40 psi through 8002 nozzle tips to the center four rows of six row plots and rototiller incorporated April 20 when the air temp. was 50F, soil temp. at six inches was 46F, wind was north at 10-15 mph, soil was dry at 0-1 inch, moist at 1-3inches, and wet below 3 inches. The rototiller was set four inches deep for treatments containing EPTC and cycloate and two inches deep for all other PPI treatments. 'Maribo Ultramono' sugarbeet was seeded 1.25 inches deep in 22 inch rows April 20. The first half of split application postemergence treatments were applied 5:00 pm May 12 when the sky was cloudy, air temp. was 85F, soil temp. at six inches was 71F, relative humidity was 23%, wind was west at 20-25 mph, soil was dry at 0-1 inch, moist at 1-4 inches, wet below 4 inches, sugarbeets were cotyledon to 2 leaf, common lambsquarters were 2-4 leaf, redroot pigweed were cotyledon to 2 leaf, and green and yellow foxtail were 1-2 leaf. The second half of split postemergence applications and all single application postemergence treatments were applied 2:00 pm May 18 when the air temp. was 80F, soil temp. at six inches was 70F, soil was dry, sugarbeets were 2-4 leaf, common lambsquarters were 2-6 leaf, redroot pigweed were 2-4 leaf, and green and yellow foxtail were 1-3 leaf. All postemergence treatments were applied in 8.5 gpa water at 40 psi through 8001 nozzle tips to the center four rows of six row plots. Common lambsquarters, redroot pigweed, and green and yellow foxtail control and sugarbeet injury were evaluated June 23. Ten sugarbeets per plot were rated June 23 for root rot using the following 0-5 values. 0) No apparent symptoms. 1) Slightly brown or discolored hypocotyl but firm under pressure. Very little root pruning. Dark and or discolored hypocotyl with slight girdling of the hypocotyl with evident root pruning and slightly soft root tissue. 3) Very dark hypocotyl and roots with collapsing root tissue but not to the degree of 4. Very evident hypocotyl girdling. Main tap root intact but very soft and deteriorated. Severe root pruning. 4) Very darkly discolored hypocotyl and roots. Root tissue completely collapsed or main tap root severed as a result of severe hypocotyl girdling. No evidence of secondary root system. 5) Dead or dying plant.

Treatment* Ra	te	Sgbt	Colq	Rrpw	G&Y Fxt1	Disease
110	/A)	inj	cntl	cntl	<u>cntl</u>	index
EPTC	2			(%)		
EPTC	3	0	0	5	68	2.10
Cycloate	4	0	0	18	70	2.07
EDTO: 0		0	0	10	65	1.72
Diethatyl	2+2	0	45	36	86	1.55
FILE 6	6	0	0	59	60	2.01
Metolachlor	3.5	0	23	49	20	2.05
Metolachlor	2	0	0	41	89	1.80
	3	0	15	56	85	1.80
Untreated Check	0	0	0	0	0	1.86
Des&Phen/Des&Phen 0.33/		1	85	69	34	2.08
Sethoxydim+Dash 0.2+0.2		0	0	0	94	1.76
Fluazifop-P+OC 0.19+0.2	25G	0	0	0	89	1.81
Clopyralid 0.	. 19	0	29	Ō	0	1.85
Endothall 0.	. 75	0	0	Ö	20	1.95
Desmedipham/Desmed 0.33/0	0.5	0	88	76	23	
				, 0	23	2.11
C.V. %		775	88	45	22	17 56
LSD 5%		NS	24	18	33	17.56
LSD 1%		NS	32		25	NS
# OF REPS		4		24	34	NS
* OC = Agway Booster Plus	F. Dach -	DACE -	4	4	4	4

\* OC = Agway Booster Plus E; Dash = BASF surfactant

SUMMARY: EPTC+cycloate gave common lambsquarters control and redroot pigweed control superior to EPTC alone or cycloate alone. Metolachlor at 3 lb/A gave less common lambsquarters control, greater redroot pigweed control, and similar foxtail control compared to EPTC+cycloate. Herbicides had no influence on the disease level on the sugarbeet roots.

Trifluralin application at different growth stages, Amenia, 1988. Diethatyl + cycloate+diallate (3+3+1 lb a.i./A) was applied to entire plot area and incorporated with an Alloway 'Seed better' April 15. 'Mitsui Monohikari' sugarbeet was seeded 1.25 inches deep in 22 inch rows May 5. Counter 15G (10 lb/A) was applied during planting using a modified in-furrow applicator. Sugarbeets were hand weeded June 10 and maintained weed free throughout the growing season. Herbicide treatments were applied in 17 gpa water at 40 psi through 8002 nozzle tips to the center four rows of six row plots on the dates listed below.

Date	Sky	Time of Day		Soil Temp	Rel	Wind Veloc (mph)	Wind Direct	Soil Moisture (inches)	Sgbt leaf Stage
June 3 June 10 June 17 June 24 July 1	sunny sunny sunny  cloudy	10:30am 9:00am 9:30am 9:00am 9:45am	95 81 75 78	78 72 74 78 72	33 28 59 67 56	5-8 10-15 15-20 0-5 2-5	south SE south south west	0-3=dry;3-5=moist 0-4=dry;4-6=moist 0-1=dry;1-5=moist 0-3=moist;3-6=wet 0-3=dry;3-6=moist	ctyl-2 2-8 4-10 6-12 8-14

All herbicide treatments were incorporated within two hours of application using an Alloway cultivator with knives and rolling cut-away disks. A John Deere two bar rotary hoe and a Melroe five bar spring tooth harrow were used for their respective treatments. Rotary hoe and harrow incorporation treatments for the June 3, June 10, June 17, and June 24 herbicide applications were done June 24. The July 1 incorporation treatments were done following herbicide application and cultivation July 1. The center two rows of each 40 foot plot were counted and harvested September 30.

		Mathad	Herbicide			Loss			
		Method		Sgbt		to	Root	Impur	Extract
		of	Application		Sucrose	Mol			Sucrose
Treatment	Rate	Incorpor.	. Date	#/80ft)					(1b/A)
	(1b/A)		,	#/0016)	(10)	(10)	( 0011, 11)		
		6 U	(1,000 2)	57	15.8	1.9	14.9	890	4068
Trifluralin	1(0.75)	Rotary Hoe	(June 3)	63	15.8	1.8	14.8	830	4088
Trifluralin			(June 3)		16.2	1.8	12.9	823	3675
Trifluralin				54		2.0	12.8	931	3414
Trifluralin	1 (1.5)	Rotary Ho	e (June 3)	55	15.4	2.0	10.9	954	2813
Trifluralin	1 (1.5)	Harrow	(June 3)	41	15.0		12.9	830	3578
Trifluralin	n(1.5)	Cult. Alo	ne (June 3)	60	15.6	1.8	12.3	870	3365
Trifluralin			e (June 10)	52	15.8	1.9		915	
Triflurali			(June 10)	66	15.4	1.9	14.6		
Triflurali		Cult. Alo	ne(June 10)	53	15.4	2.6		1246	
Triflurali			e (June 10)	39	15.2	2.1	9.9	1002	
Triflurali			(June 10)	52	15.2	1.9		921	
Triflurali			ne(June 10)	50	15.7	2.0	11.2		
Triflurali			e (June 17)	54	15.2	1.9		896	
Triflurali			(June 17)	50	16.0	1.7	12.2		
Triflurali			ne(June 17)	59	15.5	1.9	14.1	882	
			e (June 17)	69	16.0	1.9	14.2	847	
Triflurali			(June 17)	45	15.4	1.9	11.5	893	
Triflurali			ne(June 17)	66	16.1	1.8		828	4155
Triflurali	n (1.5)	cuit. Alc	me (oune 17)						

Trifluralin application at different growth stages, Amenia, 1988. (continued)

							,	
	Method	Herbicide			Loss			
	of A	Application			to		T	
Treatment Rate	Incorpor.	Date		Sucrose	Mo 1	Viol	Impur	Extract
(1b/A)			(#/80ft)	(%)	-	Tield	Index	Sucrose
				(10)	(%)	(ton/A)		(1b/A)
Trifluralin(0.75)	Rotary Hoe	(June 24)	51	16.0				
iriliuralin(U./5)	Harrow	(.luna 2/1)	CA	16.0	1.7	12.5	761	3560
Trifluralin(0.75)	Cult Alone	(June 24)	64	15.7	1.8	15.0	853	4104
Trifluralin (1.5)	Rotary Hoo	(June 24)	59	15.9	1.8	14.4	825	4020
Trifluralin (1.5)	Hannow	(June 24)		15.6	1.8	13.2	853	3600
Trifluralin (1.5)	Cult Alama	(June 24)	72	15.5	1.9	14.7	879	4012
Trifluralin(0.75)	Cuit. Alone	(June 24)	63	16.1	1.8	14.5	812	4120
Trifluralin(0.75) Trifluralin(0.75)	Rocary Hoe	(July 1)		15.5	1.8	12.6	861	3402
Triflunalin(0.75)	narrow	(July 1)	63	16.1	1.8	13.5	810	3822
Trifluralin(0.75)	Cuit. Alone	(July 1)	63	16.0	1.9	14.6	903	4076
Trifluralin (1.5)	Rotary Hoe	(July 1)	59	15.7	1.9	13.5	881	3678
Trifluralin (1.5)	Harrow	(July 1)	58	15.4	1.8	13.1	859	3566
Trifluralin (1.5)	Cult. Alone	(July 1)	63	15.6	1.8	13.5	871	
					1.0	13.3	0/1	3713
HIGH MEAN			72	16.2	2.6	15.0	1046	4155
LOW MEAN			39	15.0	1.7		1246	4155
EXP MEAN			57	15.7		9.9	761	2591
C.V. %			30		1.9	13.2	882	3617
LSD 5%			NS		15.9	19.7	18	23
LSD 1%				NS	NS	NS	NS	NS
# OF REPS			NS	NS	NS	NS	NS	NS
			4	4	4	4	4	4

The treatments had no significant effect on sugarbeet yield or population.

Simulated herbicide drift on sugarbeets, Amenia, 1988. Diethatyl+cycloate+diallate (3+3+1 lb a.i./A) was applied to entire plot area and incorporated with an Alloway 'Seed better' April 15. 'Hilleshog 5135' sugarbeet was seeded 1.25 inches deep in 22 inch rows April 21. 'Hilleshog 5135' sugarbeet was reseeded May 23 due to poor emergence of April 21 seeding. Counter 15G (10 1b/A) was applied during planting using a modified in-furrow applicator. Sethoxydim+Dash (0.2 lb a.i./A+1 qt/A) was applied to entire experiment May 24 to control wild oats and foxtail species. The early application herbicide treatments were applied 1:30 pm June 10 when the air temp. was 92F, soil temp. at six inches was 78F, relative humidity was 20%, wind was south at 15-20 mph, soil was dry at 0-4 inches, and sugarbeets were in the 2-8 leaf stage. Late application treatments were applied 10:00 am June 24 when the sky was mostly sunny, air temp. was 81F, soil temp. at six inches was 78F, relative humidity was 63%, wind was south at 0-5 mph, soil was dry at 0-3 inches, moist at 3-6 inches, and sugarbeets were in the 6-12 leaf stage. All herbicide treatments were applied in 8.5 gpa water at 40 psi through 8001 nozzle tips to the center four rows of six row plots. Plots were cultivated June 27. Sugarbeets were hand weeded June 27 and maintained weed free thoughout the growing season. Sugarbeet injury was evaluated July 1. Sugarbeets from the center two rows of 30 foot plots were counted July 18 and harvested September 29.

				Loss			
	Sgbt	Sgbt		to			Extrac
Theatment Rate	inj	Pop1	Sucrose	Mol		Index	
Treatment (1b/A)	-	#/60f1		(%)	(T/A)		(1b/A)
				0.0	0.2	121/	61
DPX-M6316+X-77 (June 10) 0.016+0.25%	99	2	12.4	2.2	0.3	1314	184
DPX-M6316+X-77 (June 10) 0.008+0.25%	99	2	12.6	2.3	0.9	1357	
DPX-M6316+X-77 (June 10) 0.004+0.259	99	11	12.8	2.2	2.2	1238	452
DPX-M6316+X-77 (June 10) 0.002+0.259		21	13.5	2.0	3.2	1089	733
DPX-M6316+X-77 (June 10) 0.001+0.259		52	14.9	1.9	5.9		1512
AC 222,293 (June 10) 0.2		23	14.0	2.0			919
AC 222,293 (built 10)		83	15.7	1.8	8.6		
AC 222,293 (built 10)		68	15.2	1.9	8.3		2188
AC 222,293 (daile 10)		73	15.5	1.7	8.8		
AC 222,293 (June 10) 0.029 Bromoxynil&MCPA (June 10) 0.12+0.1	Contract of the Contract of th	50	15.2	1.8	6.8		
BI OHOX YITT TOTAL (Sales	No. of the contract of the con	55	15.3	1.9	8.1		
Brolliox yill ration it (oatho = )		V=40=	16.1	1.8	10.2	839	
BI OHIOX VIII I dillor II (Odillo 20)	7.02		15.2	2.0	5.9	937	
Brollioxyll I III (bulle 20)			15.4	1.8	8.4		
Bromoxynil-RP (June 10) 0.0 Bromoxynil-RP (June 10) 0.0			15.7	1.8	9.6	837	2644

Simulated herbicide drift on sugarbeets, Amenia, 1988. (continued)

							•	
Treatment  DPX-M6316+X-77 (lune 24) a con	Rate (lb/A)	Sgbt inj (%)	Sgbt Popl (#/60f	Sucros		Root	Impur Index	Extrac Sucros (1b/A)
Bromoxynil&MCPA (June 24) 0.0 Bromoxynil&MCPA (June 24) 0.0 Bromoxynil-RP (June 24) Bromoxynil-RP (June 24) Bromoxynil-RP (June 24)	2+0.25% +0.25%	44 40 31 23 11 39 31 14 1 60 46 25 35 28 23	40 72 72 69 72 65 48 74 78 57 54 61 63 67 107	13.1 14.5 14.5 15.2 15.6 14.5 15.1 15.5 16.1 15.8 15.7 15.5 16.0	2.0 1.8 1.9 1.8 1.8 1.8 1.8 1.9 1.9 1.7	1.9 4.5 5.5 9.1 9.4 5.3 8.0 9.2 9.3 6.7 7.8 8.9 9.9 8.2 8.1	904 932 908 851 936 884 854 807 908 901 860 810 855 795	417 1141 1368 2383 2549 1325 2110 2498 2648 1756 2044 2454 2725 2232 2282
HIGH MEAN LOW MEAN EXP MEAN C.V. % LSD 5% LSD 1% # OF REPS		99 1 45 20 13 17 4	107 2 57 32 25 33 4	16.1 12.4 14.9 3.7 0.8 1.0	2.3 1.7 1.9 6.8 0.2 0.2	10.2 0.3 6.8 18.3 1.7 2.3	1357 795 937 9 114 151 4	2869 61 1800 20 496 657 4

This experiment did not include a true untreated check. AC 222,293 at 0.025 lb/A caused negligible sugarbeet injury so this treatment was used as a basis of comparison. DPX-M6316 caused significant sugarbeet yield losses at rates of 0.001 lb/A or higher when applied on June 10 or June 24. AC 222,293 at at 0.2 lb/A applied June 10 or 0.1 and 0.2 lb/A applied June 24 caused significant yield loss. Bromoxynil+MCPA at 0.12+0.12 lb/A applied June 10 or per acre, percent sucrose and extractable sucrose per acre were all affected by the herbicides. Injury symptoms from DPX-M6316 and AC 222,293 were identical.

Simulated herbicide drift on sugarbeets, Amenia, 1988. Diethatyl+cycloate (3+3 lb a.i./A) was applied to entire plot area and incorporated with an Alloway 'Seed better' April 18. Maribo Ultramono sugarbeet was seeded 1.25 inches deep in 22 inch rows April 18. Counter 15G (10 lb/A) was applied during planting using a modified in-furrow applicator. Plots were cultivated June 27. Sugarbeets were hand thinned to an eight inch spacing June 28. Sugarbeets were hand weeded June 28 and maintained weed free throughout the growing season. Herbicides were applied in 8.5 gpa water at 40 psi through 8001 nozzle tips to the center four rows of six row plots 1:30 pm June 30 when the air temp. was 84F, soil temp. at six inches was 78F, relative humidity was 30%, wind was east at 5-10 mph, soil was dry at 0-3 inches, moist at 3-6 inches, and sugarbeets were in the 12-18 leaf stage. Sugarbeet injury was evaluated July 16. Sugarbeets from the center two rows of 35 foot plots were counted and harvested September 27.

Countra and								
			Harves	t	Loss			F 1
		Sgbt	Sgbt		to		Impur	Extract
Twoatmant	Rate	inj	Popl	Sucrose	Mol			Sucrose
Treatment	(1b/A)	(%)(	#/70ft	(%)	(%)	(T/A)		(1b/A)
DPX-M6316+X-77 DPX-M6316+X-77 DPX-M6316+X-77 DPX-M6316+X-77 DPX-M6316+X-77 AC 222,293 AC 222,293 AC 222,293 DPX-M6316+Diclofop DPX-M6316+Diclofop DPX-M6316+Diclofop DPX-M6316+Difenzoquat DPX-M6316+Difenzoquat DPX-M6316+Difenzoquat DPX-M6316+Difenzoquat DPX-M6316+Difenzoquat DPX-M6316+Difenzoquat DPX-M6316+Difenzoquat DPX-M6316+Difenzoquat DPX-L5300 Untreated Check	0.0001+0.25% 0.00025+0.25% 0.0005+0.25% 0.001+0.25% 0.002+0.25% 0.025 0.05 0.1 0.0001+0.004 0.00025+0.008 0.0005+0.015 0.001+0.03 0.00025+0.008 0.00025+0.008	0 4 44 70 81 0 8 39 0 1 1 11 11 0 6 16	56 59 48 27 5 61 53 54 58 57 52 42 62 64 64	16.1 15.7 14.1 12.6 12.0 15.4 15.1 14.8 15.5 15.5 15.5 15.5 15.4 14.7 15.8 16.1	1.6 1.9 1.9 1.9 1.7 1.8 1.7 1.6 1.8 1.7 1.7 1.7	13.1 11.0 13.8 12.3 13.1 11.0 12.7 14.9 13.2 13.4 14.4 13.8	842 799 814 958 798	2821 3750 3395 3575 2848 3399 4183 3582 3360 4002 3945
C.V. %		21		4.0	7.6		130	585
LSD 5%		ě		1.1	0.3			
LSD 1% # OF REPS			4	4	4	1 4		1 4

#### Summary

DPX-M6316 at 0.0005 lb/A or greater caused a reduction in % sucrose, T/A, and extractable sucrose/A. AC 222,293 at 0.1 lb/A caused less yield loss than DPX-M6316 at 0.0005 lb/A. DPX-M6316 in combination with diclofop or difenzoquat caused less sugarbeet injury than DPX-M6316. However, DPX-M6316 was applied with X-77 surfactant while the combinations were applied without surfactant. DPX-L5300 caused less sugarbeet injury than DPX-M6316 but DPX-L5300 was applied without surfactant. The lack of surfactant probably reduced sugarbeet injury from DPX-L5300 and may have reduced sugarbeet injury from DPX-M6316 in combination with diclofop or difenzoquat.

Herbicides on weed free sugarbeets, Fargo, 1988. Diethatyl+cycloate (4+3 lb a.i./A) was applied to entire plot area and incorporated with an Alloway 'Seed better' April 12. 'Mitsui Monohikari' sugarbeet was seeded 1.25 inches deep in 22 inch rows May 6. Plots were cultivated June 1. Herbicide treatments were applied in 8.5 gpa water at 40 psi through 8001 nozzle tips to the center four rows of six row plots 9:30 am June 1 when the sky was mostly cloudy, air temp. was 75F, soil temp. at six inches was 70F, relative humidity was 70%, wind was south at 10-15 mph, soil was dry at 0-1 inch, moist at 1-3 inches, wet at 3-6 inches, and sugarbeets were in the 4-6 leaf stage. Sugarbeets were hand thinned to an eight inch spacing June 2. Sugarbeets were hand weeded June 2 and maintained weed free throughout the growing season. injury was evaluated June 30. The center two rows of 28 foot plots were counted and harvested October 3.

	Sgbt	Harves		Loca			
	Saht		-	Loss			
<u>Treatment* Rate</u>	inj	Pop1	Sucros	to	Root	Impur	Extrac Sucros
(1b/A	(%)	(#/56ft	t) (%)	(%)	(T/A)	Tildex	(1b/A)
Sethoxydim+Dash       0.25+0.256         Sethoxydim+Dash+28%N       0.25+0.25G+10         Sethoxydim+Dash+28%N       0.5+0.25G+10         Clopyralid       0.125         Clopyralid       0.25         Clopyralid       0.25         Seth+Clpy+Dash       0.25+0.25+0.256         Seth+Clpy+Dash       0.25+0.5+0.256         Seth+Clpy+Dash+28%N       .25+.25+.25G+10         Seth+Clpy+Dash+28%N       .25+.5+.25G+10         Seth+Clpy+Dash+28%N       .0.75         Endothall       0.75         Endothall+Clopyralid       0.5+0.19         Desmedipham       1         Desmedipham+Clopyralid       1+0.19	G 0 0 0 0 3 3 8 5 16 5 18 4 3 1 26	74 65 59 72 69 73 65 65 64 67 65 65 70 55	16.3 16.2 15.7 16.2 16.1 16.5 16.2 16.1 16.3 16.1 16.2 16.4 16.2 16.4	2.2 2.3 2.2 2.1 2.2 2.3 2.3 2.3 2.3 2.2 2.3 2.2 2.1	24.6 24.7 22.7 25.2 23.5 24.4 22.6 22.7 22.3 20.6 21.8 21.0 23.8 22.7 22.6 21.0 19.8	984 972 1060 1016 1010 912 1005 988 1043 1020 1043 995 1010 991 974 977 967	6819 6825 6014 6908 6386 6940 6190 6206 6091 5696 5898 5795 6620 6248 6320 5763 5531
C.V. % LSD 5% LSD 1%	45 4 6	8 8 10	3.7 NS	5.5 NS	7.8 2.5	8 NS	8 714
# OF REPS	4	4	NS 4	NS 4	3.4	NS 4	953 4

<sup>\* 28%</sup>N = 28% N solution containing urea and  $NH_4NO_3$ 

\* Dash = adjuvant from BASF

#### Summary

Sugarbeets treated with sethoxydim+Dash at 0.5 1b/A+1 qt/A had a lower plant population and less extractable sucrose/A than untreated sugarbeets. However, sugarbeets treated with sethoxydim+Dash+28%N had plant population and yield similar to untreated sugarbeets. Since 28%N would not be expected to safen the herbicide treatment, the observed yield loss from sethoxydim+Dash may be due to some unknown factor rather than treatment effect. All treatments including sethoxydim+clopyralid+Dash caused reduced extractable sucrose/A compared to the untreated check. Desmedipham and desmedipham + clopyralid also caused yield loss.

Herbicides on weed free sugarbeets, St. Thomas, 1988. Preplant incorporated herbicides were applied in 17 gpa water at 40 psi through 8002 nozzle tips to the center four rows of six row plots May 9 when the air temp. was 76F, soil temp. at six inches was 58F, wind was north at 10-15 mph, soil was dry at 0-1 inch, moist at 1-2 inches, and wet at 3-4 inches. A rototiller set two inches deep was used to incorporate PPI herbicide treatments. 'Hilleshog Monoricca' sugarbeet was seeded 1.25 inches deep in 22 inch rows May 9. Temik (1.5 lb a.i./A) was applied during planting using a modified in-furrow applicator. Postemergence herbicide treatments were applied in 8.5 gpa water at 40 psi through 8001 nozzle tips to the center four rows of six row plots 3:00 pm May 31 when the sky was sunny, air temp. was 94F, soil temp. at six inches was 84F, relative humidity was 44%, wind was south at 15-20 mph, soil was dry at 0-1 inch, moist at 1-2 inches, wet at 3-4 inches, and sugarbeets were in the 2 leaf stage. Sugarbeets were hand weeded June 7 and maintained weed free throughout growing season. Sugarbeets were hand thinned to an eight inch spacing June 7. Plots were cultivated June 3 and June 20. Sugarbeet injury was evaluated June 10. The center two rows of each 34 foot plot were counted and harvested September 20.

and narrosses say								
			Harvest		Loss			
		Sabt	Sgbt		to	Root		Extrac
+	Rate	inj	Pop1	Sucr	Mo1	Yield	Index	Sucros
Treatment*	(1b/A)	(%)(	#/68ft)	) (%)	(%)	(T/A)		(1b/A)
	(10/11)	(10)(						
2017 2664106	0.14+0.25G	0	40	14.8	2.3	20.9	1128	5126
R017-3664+0C	0.11.0.200	0	65	14.1	2.3	23.2	1211	5360
Untreated Check	0.75+0.25	13	49	14.4	2.6	22.0	1302	5100
Endothall+Clopyralid	0.25+0.25G	0	57	15.1	2.4	20.1	1147	5033
Sethoxydim+Dash	0.5+0.25G	0	55	15.3	2.3	21.5	1103	5483
Sethoxydim+Dash	0.25+0.25G	0	60	15.2	2.4	19.0	1125	4797
Sethoxydim+OC	0.5+0.25G	Ö	58	15.2	2.4	22.3	1147	5619
Sethoxydim+OC	0.25+0.25G	ő	60	14.4	2.4	22.3	1197	5254
Sethoxydim+Sun-It	0.5+0.25G	2	63	15.0	2.5	22.6	1204	5542
Sethoxydim+Sun-It	0.125	ō	60	15.0	2.3	22.5	1140	5600
Clopyralid	0.25	7	53	15.3	2.4	22.5	1160	5695
Clopyralid	0.5	18	47	14.6	2.5	19.6	1263	4625
Clopyralid	0.25+0.25G+1G	0	55	15.1	2.5	23.8		5928
Sethoxydim+Dash+28%N	0.5+0.25G+1G		55	15.5		20.2		5254
Sethoxydim+Dash+28%N	0.2+0.25+0.25G		41	14.1	2.6	20.8	1365	4664
Sethox+Clpy+OC	0.2+0.5+0.25G		50	14.0	2.7	23.4	1441	5133
Sethox+Clpy+OC	0.2+0.25+0.256		44	15.2		21.9	1225	
Sethox+Clpy+Dash	0.2+0.5+0.250		46	14.1		19.3	1325	
Sethox+Clpy+Dash	0.2+0.25+0.256		54	14.4			1273	
Sethox+Clpy+Sun-It	0.2+0.5+0.250		45	15.1		17.9	1237	
Sethox+Clpy+Sun-It	0.2+0.25+0.25G+1G		39	14.7		19.3	1254	
Seth+Clpy+Dash+28%N	0.2+0.5+0.25G+1G			14.7	The second second		1323	
Seth+Clpy+Dash+28%N	0.2+0.510.250110			15.4		20.8	1093	
Untreated Check				14.0			1397	
Metolachlor (ppi)				14.4			1328	
Metolachlor (ppi)				14.9			1260	4909
Metolachlor (ppi)	HISTORY STREET, ST.							

Herbicides on weed free sugarbeets, St. Thomas, 1988. (continued)

Treatment*	Rate (lb/A)	Sgbt inj	Harve Sgbt Popl #/68f	Sucr	Loss to Mol (%)	Root	Impur Index	Extrac Sucros (1b/A)
Diethatyl (ppi) Trifluralin (ppi) Ethalfluralin (ppi) 2,4-D AC 222,293 AC 222,293 AC 222,293	6 0.25 0.25 0.06 0.025 0.05 0.1	12 32 42 22 5 20 43	48 35 33 40 56 52 51	14.7 15.0 14.6 14.1 15.1 15.4 14.2	2.5 2.5 2.4 2.5 2.5 2.5 2.4	18.7 16.9 14.5 16.1 20.7 20.2 19.8	1237 1225 1200 1312 1216 1170 1262	4484 4104 3473 3654 5110 5112 4559
HIGH MEAN LOW MEAN EXP MEAN C.V. % LSD 5% LSD 1% # OF REPS		43 0 12 33 7 9 3	65 33 50 14 11 15 3	15.5 14.0 14.8 4.1 1.0 NS 3	2.7 2.3 2.5 7.0 NS NS	23.8 14.5 20.3 10.6 3.5 4.6 3	1441 1092 1229 10 192 NS 3	5928 3473 4901 12 923 1225 3

<sup>\*</sup> OC = BASF oil concentrate (Booster Plus E)

\* Dash = adjuvant from BASF

# Summary

The average of the two untreated check means was used as a basis of comparison in this experiment. Sugarbeets treated with metolachlor at 3 lb/A yielded less extractable sucrose/A than the untreated check while sugarbeets treated with metolachlor at 4 lb/A yielded similarly to the untreated check. This is not logical so perhaps unknown factors other than the treatment caused the observed reduction in yield. Trifluralin at 0.25 lb/A, ethalfluralin at 0.25 lb/A, and 2,4-D at 0.06 lb/A caused a reduction in root yield and extractable sucrose/A. Sethoxydim+Dash+28%N at 0.5 lb/A+ 1 qt/A+ 1 gal/A and sethoxydim+clopyralid+Booster Plus E at 0.2+0.5 lb/A + 1 qt/A did not affect sugarbeet yield. However, sethoxydim+clopyralid+Dash at 0.2+0.5 lb/A+ 1 qt/A, sethoxydim + clopyralid + Sun-It at 0.2 + 0.5 lb/A + 1 qt/A, and sethoxydim + clopyralid+Dash+28%N at 0.2+0.5 lb/A + 1 qt/A + 1 gal/A reduced extractable sucrose/A as compared to the untreated check.

<sup>\*</sup> Sun-It = Agsco sunflower methyl ester

<sup>\* 28%</sup>N = 28% N solution containing urea and  $NH_4NO_3$ 

Time of application of sethoxydim and clopyralid plus additives, Amenia, 1988. Diethatyl+cycloate+diallate (3+3+1 lb a.i./A) was applied to entire plot area and incorporated with an Alloway 'Seed better' April 15. 'Mitsui Monohikari' sugarbeet was seeded 1.25 inches deep in 22 inch rows May 5. Counter 15G (10 lb/A) was applied during planting using a modified in-furrow applicator. Sugarbeets were hand weeded June 10 and maintained weed free throughout the growing season. Herbicide treatments were applied in 8.5 gpa water at 40 psi through 8001 nozzle tips to the center four rows of six row plots on the dates listed below.

			Time of	Air	5 inch Soil	Rel	Wind	Wind Direct	Soil Moisture	Sgbt leaf Stage
Date	Day	Sky	Day	Temp	Temp	(%)	(mph)		(inches)	
June	3 (D1)	sunny	12:30		78	33	5-8		0-3=dry;3-5=moist	ctyl-2
June		sunny	10:00	81	72	28		south	0-4=dry;4-6=moist 0-1=dry;1-5=moist	
June		sunny	2:00	76	74	42	5-8	SE south	0-3=moist;3-6=wet	
June					78 72	57 56	0-5 2-5	west	0-3=dry;3-6=moist	
July	1 (D5)	cloudy	10:30	70	12	30				

Plots were cultivated June 10 and June 24. The center two rows of each 35 foot plot were counted July 25 and harvested September 30.

							CONTRACTOR OF THE PARTY
				oss			
		Sgbt		to	Root	Impur	Extrac
Tuestmont*	Rate F	opl	Sucrose		Yield		Sucros
Treatment.	(1b/A)(#/	/70ft	) (%)	(%)(	(ton/A)		(1b/A)
Clopyralid (D1) Clopyralid (D1) Sethoxydim+Sun-It (D1) Sethoxydim+Dash (D1) Sethoxydim+Dash (D1) Sethoxydim+Dash (D1) Sethoxydim+Dash+28%N (D1) Clpy+Seth+Dash+28%N (D1) Clpy+Seth+Dash+28%N (D1) Untreated Check Desmed (D1,2,3,4,5) 0.25/0 Desmedipam (D1,3,5) Desmedipham (D3) Clopyralid (D2)	(1b/A)(#/ 0.19 0.38 0.25+0.25G 0.5+0.25G 0.5+0.25G 0.5+0.25G 0.5+0.25G+1G .19+0.5+0.25G+1G				14.6 11.5 12.1 13.2 13.6 13.3 12.8 14.1 13.1 11.7 13.4 13.4	839 924 910 823 783 849 834 909 822 856 869 891 899	(1b/A)  4051 2984 3244 3715 3788 3660 3526 3856 3557 3214 3660 3635 3646 3449
Clopyralid (D2) Sethoxydim+Sun-It (D2)	0.25+0.25G 0.5+0.25G	69 71	15.7 15.8	1.8 1.7	13.8	774	3837
Sethoxydim+Sun-It (D2) Sethoxydim+Dash (D2)	0.25+0.25G 0.5+0.25G	63 64	15.6 16.0	1.8			
Sethoxydim+Dash (D2) Sethoxydim+Dash+28%N (D2) Clpy+Seth+Dash+28%N (D2)	0.5+0.25G+1G	58 55		1.7	13.2	753	3806

Time of application of sethoxydim and clopyralid plus additives, Amenia, 1988.

Treatment*		Sgbt		Loss		Tmm	
11 Cacinett	Rate	Pop1	Sucros	e Mol	Vield	Impur	Extrac
	(1b/A)(	#/70f	t) (%)	(%)	(ton/A)	Index	
Untreated Check				(10)	( con/ A,		(1b/A)
Clopyralid (D3)		54	15.5	1.9	12.6	901	2264
Clopyralid (D3)	0.19	53	15.0	1.9	10.6	947	3364
Sethoxydim+Sun-It (D3)	0.38	56	15.2	1.9	11.5	932	2737
Sethoxydim+Sun-It (D3)	0.25+0.25G	61	15.6	1.9	13.8	908	3022
Sethoxydim+Dash (D3)	0.5+0.25G	76	15.7	1.8	13.1	823	3742
Sethoxydim+Dash (D3)	0.25+0.25G	60	15.8	1.8	14.3		3609
Sethoxydim+Dash (D3)	0.5+0.25G	62	15.5	1.9	13.9	845	3942
Sethoxydim+Dash+28%N (D3)	0.5+0.25G+1G	51	15.8	1.7	11.7	878	3753
Clpy+Seth+Dash+28%N (D3) 0.19- Clopyralid (D4)	0.5+0.25G+1G	66	15.9	1.8	13.2	789	3269
Clopyralid (D4)	0.19	70	15.7	1.8	12.7	823	3651
Sethorydim Com II (D4)	0.38	60	15.5	1.9	12.7	850	3471
Sethoxydim+Sun-It (D4)	0.25+0.25G	56	15.5	1.9	12.7	911	3189
Sethoxydim+Sun-It (D4)	0.5+0.25G	57	15.7	1.8	11.8	896	3432
Sethoxydim+Dash (D4)	0.25+0.25G	68	15.7	1.9	13.3	829	3251
Sethoxydim+Dash (D4)	0.5+0.25G	54	15.4	1.8	12.0	873	3663
Sethoxydim+Dash+28%N (D4)	0.5+0.25G+1G	61	15.6	1.8	12.7	857	3250
Clpy+Seth+Dash+28%N (D4) 0.19+	0.5+0.25G+1G	61	15.2	1.9	11.8	847	3497
riopyraliu (DD)	0.19	64	15.5	1.9	11.7	925	3117
Clopyralid (D5)	0.38	60	14.9	2.0	9.5	890	3161
ethoxydim+Sun-It (D5)	0.25+0.25G	67	15.7	1.9	12.4	959	2422
ethoxydim+Sun-It (D5)	0.5+0.25G	64	15.7	1.8	11.5	861	3382
ethoxydim+Dash (D5)	0.25+0.25G	49	15.1	1.9	11.6	831	3156
ethoxydim+Dash (D5)	0.5+0.25G	88	15.9	1.7	13.4	926	3016
ethoxydim+Dash+28%N (D5)	TIO OFFI	56	15.7	1.8	12.5	802	3758
lpy+Seth+Dash+28%N (D5) 0.19+0	).5+0.25G+1G	80	15.0	2.0	11.4	837	3412
IGH MEAN			13.0	2.0	11.4	977	2935
OW MEAN		88	16.2	2.0	14.6	077	1054
XP MEAN		49		1.7	9.5	977	4051
.V. %		62			12.7	753	2422
SD 5%		25			15.2	868	3447
SD 1%		NS		0.2	NS	9	18
OF REPS		NS	NS	NS	NS NS	111	NS
OI NEPS		4	4	4	4	145	NS
Sun-It = Agsco sunflower moth					4	4	4

The treatments had no significant effect on sugarbeet yield or population.

<sup>\*</sup> Sun-It = Agsco sunflower methyl ester \* 28%N = 28% N solution containing urea and NH<sub>4</sub>NO<sub>3</sub> \* Dash = BASF surfactant

Multispecies screening of preplant incorporated, preemergence, and postemergence herbicides, Fargo (NW section 22), 1988. Preplant incorporated herbicide treatments were applied and rototiller incorporated May 18 when the sky was sunny, air temp. was 88F, soil temp. at six inches was 68F, wind was southwest at 25-35 mph, soil was dry at 0-2 inches and moist at 2-4 inches. The rototiller was operated four inches deep for EPTC and cycloate and two inches deep for all other PPI herbicides. 'Wheaton' wheat, 'Robust' barley, 'ND810104' oats, 'Cando' durum, 'Linton' flax, 'S541' safflower, common lentils, 'Topaz' canola, 'Kirby' mustard, 'Siberian' foxtail millet, 'McCall' soybean, 'Hyden' navy bean, 'Intersate 201' corn, 'Interstate 3001' sunflower, 'Maribo Ultramono' sugarbeet, wild buckwheat, redroot pigweed, kochia, and common lambsquarters were seeded in strips across the herbicide plots May Preemergence herbicide treatments were applied May 20 when the air temp. 73F, soil temp. at six inches was 68F, relative humidity was 54%, wind was northwest at 5-8 mph, soil was dry at 0-1 inch and moist at 1-6 inches. All soil applied treatments were applied in 17 gpa water at 40 psi through 8002 nozzle tips to the center 7 feet of 11 foot plots. Postemergence herbicide treatments were applied in 8.5 gpa water at 40 psi through 8001 nozzle tips to the center 7 feet of 11 foot plots 9:00 am June 9 when the sky was sunny, air temp. was 70F, soil temp. at six inches was 70F, relative humidity was 33%, wind was south at 12-18 mph, soil was dry at 0-2 inches and moist at 2-4inches, green foxtail was 0.5 to 2 inches tall, and kochia was 0.5 to 2.5 inches tall. Kochia and green foxtail control were evaluated in all three experiments July 14. All other species either did not emerge because of dry soil conditions or were crowded out by the high density of green foxtail and kochia.

Preplant incorporated herbicides

	brated Herbicides	Vanhin	Green Foxtail	
		Kochia	control	
Treatment	Rate	control	(%)	
	(1b/A)	(%)	82	
SAN-582H	1.25	0	95	
SAN-582H	1.75	10		
Isoxaben	0.08	63	0	
Imazethapyr	0.03	99	94	
Imazethapyr	0.06	99	97	
Cyanazine-L	2	87	47	
Alachlor		0	82	
Metolachlor	3	0	96	
	3 3 4	18	99	
EPTC	4	0	91	
Cycloate	0.75	72	53	
Clomazone	0.75	93	95	
Trifluralin	0.75	91	97	
Ethalfluralin	0.125	98	87	
A-1237		91	50	
Metribuzin-DF	0.375	50	18	
AC 222,293	0.25	86	27	
DPX-A7881-22	0.02	7	0	
DPX-V9360-34	0.02	98	78	
CGA-136872	0.02	90		
		18	14	
C.V. %		17	16	
LSD 5%			21	
LSD 1%		22 3	3	
# OF REPS		3	<u> </u>	

# (continued) Preemergence herbicides

<u>Treatment*</u>	Data	Kochia	Green Foxtail
	Rate	control	control
Propachlor	(1b/A)	(%)	(%)
Metolachlor	5 3 3 3	0	83
Alachlor	3	0	68
Chloramben	3	27	57
Pendimethalin		99	97
A-1237	1.5	48	
A-1237	0.062	88	32
SAN-582H	0.125	97	23
	1.25	13	82
SAN-582H	1.75	15	70
		15	71
C.V. %		30	
LSD 5%		29	27
LSD 1%		22	31
# OF REPS		30	42
Postemergence herb	Dicides	3	3
JPX-M6316+X-77	0.016+0.25%		
DPX-L5300+X-77	0.016+0.25%	93	0
DPX-A7881-22+X-77	0.016+0.25%	99	0 7
PX-A7881-22+X-77	0.016+0.25%	0	ó
PX-V9360-34+X-77	0.031+0.25%	28	32
PX-V9360-34+X-77	0.016+0.25%	32	93
mazethapyr+X-77	0.031+0.25%	45	90
mazethapyr-X=//	0.03+0.25%	68	
mazethapyr+X-77	0.06+0.25%	80	85
lopyralid	0.1	0	83
luroxypyr	0.1	96	0
omesafen+X-77	0.2+0.25%	15	0
actofen+X-77	0.18+0.25%	33	0
entazon+PO	0.75+0.25G		0
cifluorfen+X-77	0.375+0.25%	82	0
ent+Acif+X-77 (	0.6+0.25+0.25%	0	13
romoxynı	0.25	43	0
icamba	0.12	80	0
,4-D	0.25	86	0
,4-DB	0.25	58	0
yphosate+X-77		0	Ō
araquat+X-77	0.25+0.25%	80	97
1237	0.5+0.25%	62	90
1237	0.016	88	0
4243	0.031	98	Ö
4243	0.03	73	0
23121	0.06	81	0
23121	0.004	0	
63596+X-77	0.018	ő	0
	0.044+0.05%	Ö	0
63596+X-77	0.088+0.05%	0	0
A-136872	0.016	12	67
A-136872	0.031		35
		45	60
1. %		20	
5%		28	44
1%		22	18
)F RFPS		29	23
-77	urfactant from Chevr	3	3

<sup>\*</sup> X-77 = non-ionic surfactant from Chevron Chemical Co. \* PO = petroleum oil adjuvant from BASF (Booster Plus E)

Injury to sugarbeets and other crops from herbicide carryover. Dexter, Alan G., John D. Nalewaja, and Dallas E. Peterson. The objective of these experiments was to detect phytotoxic soil residual in 1988 from herbicides applied in 1987. Experiments were established on a silty clay with 4.0% organic matter and pH 8.0 approximately 5 miles north of Fargo, ND. "Marshall' wheat was seeded at 75 lb/A in one experiment and "McCall' soybean was seeded in 6-inch rows at 69 lb/A in the other experiment on June 2, 1987. Herbicides were applied to both experiments from 3:00 to 4:00 pm June 24, 1987 when the air temperature was 73 F, six-inch soil temperature was 71 F, relative humidity was 54% and the soil was dry at 0 to 1 inch, moist at 1 to 2 inches, and wet at 2 to 4 inches. At application, soybeans were from the cotyledon to two trifoliolate stage and wheat was six inches tall. Herbicides were applied in 8.5 gpa of water at 40 psi through 8001 nozzles to the center 7 feet of 14 by 45 foot plots arranged in a randomized complete block design with four replications. A chisel plow was used for fall tillage in 1987 and a field cultivator was used for seedbed preparation in the spring of 1988. "McCall' soybean, 'N.D. 810104' oats, 'Linton' flax, and 'Maribo Ultramono' sugarbeet were seeded in the soybean experiment across the herbicide treated blocks on May 18 when the soil was dry at 0 to 2 inches and moist at 2 to 4 inches. Rain totalling 0.9 inches fell from May 19 through May 22, 1988. Total rain from April through August of 1987 was 10.3 inches compared to normal of 12.7 inches. A natural dense population of redroot pigweed and yellow foxtail emerged in the experiments. Crop injury and weed control were evaluated June 11 and June 27 except yellow foxtail was evaluated only on June 11.

DPX-L5300 at 0.03 lb/A applied in 1987 caused slight visible injury to soybeans and controlled redroot pigweed and yellow foxtail in 1988. Soybeans and redroot pigweed exhibited less injury on June 11 than on June 27. AC 222,293 at 0.3 lb/A applied in 1987 injured soybean, sugarbeet, and oats, and controlled redroot pigweed in 1988 while AC 222,293 at 0.6 lb/A injured or controlled all species. DPX-M6316 applied in 1987 did not significantly injure the species evaluated in 1988.

Imazethapyr at 0.06 and 0.12 lb/A applied in 1987 injured flax, wheat, sugarbeet and controlled redroot pigweed and yellow foxtail in 1988. Imazethapyr at 0.12 lb/A injured corn while 0.06 lb/A did not. Injury from imazethapyr to flax and wheat tended to be greater on June 27 than on June 11. Fomesafen at 0.5 lb/A applied in 1987 injured flax and sugarbeet and controlled redroot pigweed and yellow foxtail in 1988. Fomesafen at 0.25 lb/A only affected redroot pigweed and yellow foxtail. Lactofen and acifluorfen did not injure the species evaluated in 1988. (Dept. of Crop and Weed Sciences, published with the approval of the Agric. Exp. Stn., North Dakota State Univ. Fargo)

Table 1. Crop injury and weed control in 1988 from herbicides applied to wheat in 1987. (Dexter, Nalewaja, and Peterson).

1987 treatment	Rate (1b/A)	Soyb 6/11	ean 6/27		ax 6/27 njury		6/27	0a 6/11		Rr 6/11 (% con	6/27	Yeft 6/11 1988)
DPX-M6316 DPX-M6316 DPX-L5300 DPX-L5300 AC 222,293 AC 222,293 Untreated Check	0.015 0.03 0.015 0.03 0.3 0.6	0 0 0 13 3 28	0 0 0 0 14 24	0 0 0 5 4 50	0 0 0 0 0 0 26 0	0 0 0 0 39 99	0 0 0 0 65 91	0 0 0 0 0 8	0 0 0 0 11 40 0	0 0 0 68 86 96	0 5 0 30 85 95	0 0 0 23 15 30 0
Mean LSD (0.05)		6	5 14	8 9	4 NS	20 25	22 15	1 NS	7 11	36 10	31 22	10 23

Table 2. Crop injury and weed control in 1988 from herbicides applied to soybean in 1987. (Dexter, Nalewaja, and Peterson).

1987 treatment	Rate (1b/A)	6/11 	6/27	6/11	lax 6/27 injury	6/11		Suga 6/11	rbeet 6/27	Rr 6/11	pw 6/27 ntrol,	Yeft 6/11 1988)
Imazethapyr Imazethapyr Fomesafen Fomesafen Lactofen Lactofen Acifluorfen Acifluorfen Untreated Check	0.06 0.12 0.25 0.5 0.2 0.4 0.38 0.75	1 23 0 0 0 0 0 0	4 21 0 0 0 0 0 0 0	91 97 3 43 0 0 0	96 99 0 19 0 0 0	0 24 0 4 0 0 0	19 31 4 5 0 0 0	99 99 8 63 0 0 0	99 99 3 50 0 0	99 99 65 91 0 0 0	99 97 57 88 0 3 0	97 99 28 74 0 0 0
Mean LSD (0.05)		3 3	3 8	26 13	24 18	3 9	7 14	30 12	28 13	39 9	38 5	33 18

Three species evaluation of the effect of additives on antagonism, Amenia, 1988. 'Siberian' foxtail millet, 'Wheaton' wheat, and 'ND810104' oats were seeded in adjacent six foot strips across the herbicide plots May 23. Herbicide treatments were applied in 8.5 gpa water at 40 psi through 8001 nozzle tips to a 7.33 foot strip through the center of 11 foot plots and across the three grass species. Herbicide application was between 10:00 am and 1:00 pm June 22 when the air temp. was 84F, soil temp. at six inches was 79F, relative humidity was 35%, wind was north at 10-20 mph, sky was partly cloudy, soil was dry at 0-1 inches, moist at 1-6 inches, foxtail millet was 8 inches tall, wheat was 6 inches tall, and oats was 7 inches tall. Foxtail millet, wheat, and oats control was evaluated July 11.

nillet, wheat, and dats control the		Ftmi	Wheat	Oats
	Rate	inj	inj	in;
reatment	(1b/A)		(%)	
	0.05	37	82	48
)PX-Y6202		68	94	88
PX-Y6202	0.1	0	Ö	0
1017-3664+Booster Plus E	0.035+0.25G	0	10	0
R017-3664+Booster Plus E	0.07+0.25G	13	50	0
R017-3664+Booster Plus E	0.105+0.25G	40	58	10
2017-3664+Rooster Plus E	0.14+0.25G	45	92	79
Fluazifop-P+Booster Plus E	0.09+0.25G	57	97	98
Fluazifop-P+Booster Plus E	0.188+0.25G	83	92	91
Sethoxydim+Dash	0.1+0.25G	96	. 98	98
c-thewidim+Nach	0.2+0.25G	65	67	60
cathogradim+Clonyralid+Booster Plus L	0.1+0.2+0.25G	93	95	9:
Sethoxydim+Clopyralid+Booster Plus L	0.2+0.2+0.25G	85	95	8
Sethoxydim+Clopyralid+Dasn	0.1+0.2+0.25G	96	98	9
Sethoxydim+Clopyralid+Dash	0.2+0.2+0.25G	81	88	8
Sethoxydim+Clopy+Sun-It	0.1+0.2+0.25G	92	94	9
	0.2+0.2+0.25G	85	94	9
	.1+0.2+0.25G+1G	92	98	9
seth+Clony+Dash+28%N	.2+0.2+0.25G+1G 0.1+1+0.25G	42	55	3
Seth+Desmedipham+Booster Plus E	0.1+1+0.25G 0.2+1+0.25G	67	77	7
coth+Desmedipham+Booster Plus E	0.2+1+0.25G 0.3+1+0.25G	89	88	8
Seth+Desmedipham+Booster Plus E	0.4+1+0.25G	92	95	Ġ
soth+Desmedinham+Booster Plus E	0.5+1+0.25G	92	94	9
Seth+Desmedipham+Booster Plus E	0.5+1+0.25G	52	70	(
Seth+Desmedipham+Dash	0.1+1+0.25G	68	75	
Seth+Desmedipham+Dash	0.2+1+0.25G	86	90	
Seth+Desmedipham+Dash	0.4+1+0.25G	95	96	
Seth+Desmedipham+Dash	0.5+1+0.25G	93	94	
Seth+Desmedipham+Dash	0.1+1+0.25G	65	73	
Seth+Desmedipham+Sun-It	0.2+1+0.25G	83	83	
Seth+Desmedipham+Sun-It	0.3+1+0.25G	90	94	
Seth+Desmedipham+Sun-It	0.4+1+0.25G	94	97	
Seth+Desmedipham+Sun-It	0.5+1+0.25G	95	97	
Seth+Desmedipham+Sun-It	0.1+1	33	28	
Sethoxydim+Desmedipham	0.2+1	45	60	
Sethoxydim+Desmedipham	0.3+1	67	75	
Sethoxydim+Desmedipham	0.4+1	80	83	
Sethoxydim+Desmedipham	0.5+1	86	89	
co+boyydim+Desmedipham	0.09+1+0.25G	23	78	
Flua-D+Desmedipham+Booster Plus E	0.188+1+0.25G	40	92	
Elua-D+Desmedipham+Booster Plus E	0.28+1+0.25G	48	96	
Flua-P+Desmedipham+Booster Plus E	<b>7.29 -</b>			

Three species evaluation of the effect of additives on antagonism, Amenia, 1988. (continued)

<u>Treatment</u>	Rate	Ftmi	Wheat	Oats
	(1b/A)	inj	<u>inj</u>	inj
Fluazifop-P+Herbimax			(%)	
Fluazifop-P+Herbimax	0.09+0.25G	33	96	98
Fluazifop-P+LI-700	0.188+0.25G	57	98	99
Fluazifop-P+LI-700	0.09+0.25G	23	85	86
Fluazifop-P+Stollers	0.188+0.25G	45	94	97
Fluazifop-P+Stollers	0.09+0.25G	22	74	48
Fluazifop-P+Desmedinham	0.188+0.25G	32	92	91
Fluazifop-P+Desmedinham	0.09+1	8	67	50
Fluazifop-P+Desmedipham	0.188+1	27	86	78
Sethoxydim+Sun-It	0.28+1	45	92	99
Sethoxydim+Sun-It	0.1+0.25G	79	88	83
Sethoxydim+Booster Plus F	0.2+0.25G	97	97	99
sethoxydim+Booster Plus F	0.1+0.25G 0.2+0.25G	83	85	79
sethoxydim+Herbimax	0.2+0.25G 0.1+0.25G	94	96	96
Sethoxydim+Herbimax	0.2+0.25G	80	77	78
Sethoxydim+LI-700	0.1+0.25G	95	95	93
Sethoxydim+LI-700	0.2+0.25G	60 88	62	43
Sethoxydim+Stollers	0.1+0.25G	77	90	93
Sethoxydim+Stollers	0.2+0.25G	90	80 94	76
Sethoxydim	0.1	45	25	96
ethoxydim Tuazifop-P	0.2	82	77	10
luazifop-P	0.09	38	83	82
017-366/+Dogmod Dogg 4	0.188	48	95	67 95
017-3664+Desmed+Booster Plus E ethoxydim+Desmedipham+Herbimax	0.105+1+0.25G	8	10	13
ethoxydim+Desmedipham+LI-700	0.1+1+0.25G	62	62	53
ethoxydim+Desmedipham+Stollers	0.1+1+0.25G	28	30	17
luazifon-P+Clonyralid+Poorter D1	0.1+1+0.25G	62	62	42
luazifop-P+Clopyralid+Booster Plus	s E 0.09+0.2+0.25G	35	93	77
IGH MEAN				
OW MEAN		97	98	99
XP MEAN		0	0	0
.V. %		62	79	72
SD 5%		14	8	11
SD_1%		14	10	13
OF REPS		19	13	17
		3	3	3

DPX-Y6202 at 0.1 lb/A and fluazifop-P at 0.188 lb/A + Herbimax gave less foxtail millet control than sethoxydim+Dash at 0.1 lb/A + 1 qt/A. R017-3664 + Booster Plus E gave poor control of all three species. Sethoxydim+clopyralid +Dash gave grass control similar to sethoxydim+Dash. Sethoxydim+desmedipham + Dash at 0.3 + 1 lb/A + 1 qt/A gave grass control similar to sethoxydim+Dash at control than sethoxydim + other additives. Desmedipham antagonized grass control from sethoxydim and fluazifop-P regardless of additive. Fluazifop-P + to fluazifop-P at 0.09 lb/A + 0ther additives.

Broadleaf herbicides plus additives, Amenia, 1988. Bioassay strips of 'Linton' flax, 'McCall' soybean, 'Beta 6186' sugarbeet, and 'Kirby' mustard were seeded in adjacent six foot strips across the herbicide plots May 23. Sethoxydim (0.2 lb ai/A) plus Bio-Veg crop oil (1 qt/A) was applied to all bioassay strips June 10 to control foxtail and wild oats. Herbicide treatments were applied between 9:00 am and 1:00 pm June 16 when the sky was mostly sunny, air temp. was 76F, soil temp. at six inches was 74F, relative humidity was 42%, wind was southeast at 2-6 mph, soil was dry at 0-1 inches, moist at 1-5 inches, wet below 5 inches, common lambsquarters were 6 leaf to 3 inches tall, mustard was 4 leaf (1.5 inches tall) to 8 leaf (3 inches tall), flax was 2-3 inches tall, and soybeans had 2 to 3 trifoliolate leaves. The sugarbeet seed did not emerge due to dry soil conditions at planting. All herbicide treatments were applied in 8.5 gpa water at 40 psi through 8001 nozzle tips to the center 7.33 feet of 11 foot plots. Common lambsquarters, flax, tame mustard, and soybean control were evaluated July 2.

flax, tame mustard, and so	Sypean const .				
		Colq	Flax	Tamu	Soyb cntl
_	Rate	cntl	cntl	cntl	CHUI
Treatment*	(1b/A)		(%)	)	
				4.4	3
Ddinham	0.25	30	9	44	3
Desmedipham	0.5	22	12	35	9
Desmedipham	1	48	24	60	56
Desmedipham	0.25+1	29	28	53	56
Desmedipham+Dalapon	0.5+1	26	19	53	
Desmedipham+Dalapon	1+1	65	65	71	64
Desmedipham+Dalapon	0.25+1.1	63	60	72	65
Desmedipham+Ethofumesate	0.5+1.1	83	65	79	68
Desmedipham+Ethofumesate	1+1.1	92	78	87	78
Desmedipham+Ethofumesate	0.25+0.56	53	38	59	53
Desmedipham+Ethotumesate	0.5+0.56	73	45	63	56
Desmedinham+Ethotumesate	1+0.56	70	65	89	70
Desmedipham+Ethotumesate		13	13	43	36
Desmedipham+Ethotumesate		36	23	60	45
Desmedinham+Ethorumesate	0.510.15	70	61	76	59
Desmedipham+Ethotumesate	1.0.13	6	0	18	3 3
Desmedipham+Spraymate	0.2310.234	21	14	41	3
Desmedipham+Spraymate	0.5+0.25G	49	15	54	9
Desmedipham+Spraymate	1+0.25G	0	0	10	4
Pyrazon	3.7		5	44	18
Pyrazon+Dash	3.7+0.25G	11	20	58	39
Pyrazon+Dash+28%N	3.7+0.25G+1G	30	5	36	18
Desmedipham+Pyrazon	0.25+3.7	0	8	26	13
Desmedipham+Pyrazon	0.5+3.7	10	39	64	24
Desmedipham+Pyrazon	1+3.7	66		59	25
Desm+Pyrazon+Dash	0.25+3.7+0.25G	29	14	68	29
Desm+Pyrazon+Dash	0.5+3.7+0.25G	55	31	88	53
Desm+Pyrazon+Dash	1+3.7+0.25G	73	63		60
Desm+Pyra+BAS-090025	0.25+3.7+0.25G	54	31	76	55
Desm+Pyra+BAS-090025	0.5+3.7+0.25G	74	60	80	59 59
Desm+Pyra+BAS-090025	1+3.7+0.25G	80	75	85	30
Desm+Pyra+BAS-090025	0.25+3.7+0.25G	9	11	58	45
Desm+Pyrazon+Sun-It	0.5+3.7+0.25G	38	33	72	
Desm+Pyrazon+Sun-It	1+3.7+0.25G	85	66	78	66
Desm+Pyrazon+Sun-It					

Broadleaf herbicides plus additives, Amenia, 1988. (continued)

Treatment*		Cold te cntl		Tamu cntl	Soyb
	(16	/A)			<u>cnt1</u>
Desm+Pyrazon+Etho Desm+Pyrazon+Etho Desm+Pyrazon+Etho Desm+Pyrazon+Dalapon Desm+Pyrazon+Dalapon Desm+Pyrazon+Dalapon Desm+Pyrazon+Dalapon Desm+Pyra+Dash+28%N Desm+Pyra+Dash+28%N Desm+Pyra+Dash+28%N Clopyralid Clopyralid Clopyralid Clopyralid+OC Clopyralid+OC Clopyralid+OC Clopyralid+OC+28%N Desmedipham+Trigger Desmedipham+Trigger Desmedipham+Trigger Desmedipham+Nu-Film-P Desmedipham+Nu-Film-P Desmedipham+Nu-Film-P	0.25+3.7+0 0.5+3.7+0 1+3.7+0 0.25+3. 0.5+3. 1+3. 0.25+3.7+0.25G 0.5+3.7+0.25G 1+3.7+0.25G	7/A)  1.56	40 58 71 16 30 35 16 46 55 0 0 0 0 0 0 5 13 38 6 28 45	(%) 84 92 92 53 71 71 62 73 75 0 0 0 0 0 0 26 28 60 27 53 60	85 82 83 55 68 65 24 43 50 87 96 99 90 99 96 99 13 9 16 0 23 39
HIGH MEAN  OW MEAN  EXP MEAN  EV. %  SD 5%  SD 1%  OF REPS		92 0 43 32 19 26 4	78 0 29 34 14 18	92 0 52 20 14 19 4	99 0 47 22 14 19 4


<sup>\*</sup> OC = Agway Booster Plus E

Sugarbeet populations were insufficient for evaluation. Trigger did not affect test species control from desmedipham. Ethofumesate increased test species control from desmedipham more than dalapon. Ethofumesate at 1.1 increased or tended to increase control from desmedipham more than ethofumesate at 0.56 lb/A and 0.56 lb/A generally was more effective than 0.19 1b/A. Desmedipham+pyrazon gave control similar to desmedipham alone. BAS-090025 increased or tended to increase control from desmedipham+pyrazon more than Dash or Sun-It. Desmedipham+pyrazon+ethofumesate tended to give greater control than desmedipham+pyrazon+BAS-090025.

<sup>\*</sup> Sun-It = Agsco sunflower methyl ester

<sup>\*</sup> 28%N = 28% N solution containing urea and  $NH_4NO_3$ 

<sup>\*</sup> Dash = BASF surfactant



Wild oats control in Wheat, Fargo 1988. 'Wheaton' Hard Red Spring wheat was seeded 2.5 inch deep in 7 inch spaced rows on April 12. Emergence of wheat was poor because of dry soil conditions. Treatments (S1) were applied to 2-to 3-leaf wheat and wild oats on May 13 with 45F, 40% RH, no wind, and a clear sky. Treatments (S2) were applied to 5-leaf wheat and wild oats on May 26 with 72F, 65% RH, and 8 mph wind. All treatments were applied in 8.5 gpa at of 10 by 30 ft plots. The experiment was a randomized complete block with four replications. Evaluation was on July 22 and wild oats density exceeded 10 plants/sq. ft.

Treatment	Rate	Wheat	Wild Oat
	(oz/A)	injury	control
	(02/14)	(%)	(%)
Diclofop(S1)	12		
Diclofop(S1)	16	0	45
Diclofop+PO(S1)		0	79
Diclofop+MS(S1)	12+0.125G	1	90
AC222293(S1)	12+0.125G	3	89
10222293 (SI)	4	3 1	
AC222293(S1)	5	4	87
AC222293 (S1)	6	4	90
AC222293+PO(S1)	4+0.125G		93
C222293+MS(S1)	4+0.125G	0 3 0	96
Diclofop(S2)	16	3	98
Diclofop+PO(S2)	16+0.125G	Ü	58
liclofop+MS(S2)	16+0.125G	0	69
C222293(S2)	4	4 3 9 6 5 5 5 6 3	75
C222293 (S2)	6	3	94
C222293 (S2)	8	9	94
C222293+PO(S2)		6	98
C222293+MS(S2)	4+0.125G	5	92
ifenzoquat(S2)	4+0.125G	5	95
ifonzaguat (S2)	10	5	94
ifenzoquat(S2)	12	6	
0E-7113-04H(S2)	2.6	3	98
OE-7125-01H(S2)	11	1	90
ntreated	0	Ô	87
		0	0
IGH MEAN		0	
OW MEAN		9	98
(P MEAN		0	0
.V. %		3	82
SD 5%		142	11
SD 1%		NS	13
OF REPS		NS	17
The state of the s		4	4

# Summary

Adjuvants enhanced or tended to enhance wild oats control with diclofop and AC-222293 applied at the 2-leaf stage. AC-222293, difenzoquat, HOE-7113-04H, and HOE-7125-01H all gave 87% or more wild oats control without important injury to wheat. At the 4-leaf stage adjuvants did not enhance wild oats control with AC-222293, but tended to enhance diclofop. Grain yield was not obtained because the drought stress caused variable stands and limited growth.

Wild oats control in wheat, Langdon 1988. 'Coteau' Hard Red spring wheat was seeded on May 6. S1 treatments were to 3-leaf wheat 2-leaf wild oats and green foxtail on May 26. S2 treatments were to 4.5-leaf wheat and wild oats on June 1. All treatments were applied in 8.5 gpa at 35 psi with a bicycle wheel type sprayer to an 8 ft wide area the length of 10 by 25 ft plots. The experiment was a randomized complete block with four replications. Entire plot area was treated with cholosulforon + 2,4-D in early June. Evaluation was on July 20

was on oary to						
	Rate	Wheat injury	Yield	Twt	Wioa	Grft
<u>Treatment</u>	(oz/A)	(%)	(bu/A)	(1b/bu)	(% coi	ntrol)
	(02/1/)	(10)				
Diclofop(S1)	12	0	26.4	59.0	76	88
Diclofop(S1)	16	0	30.1	59.5	81	86
Diclofop+PO(S1)	12+0.125G	0	23.6	60.8	85	89
Diclofop+MS(S1)	12+0.125G	0	25.8	60.0	83	85
AC 222,293(S1)	4	0	21.7	60.4	95	30
AC 222,293(S1)	5	0	25.5	62.4	96	33
AC 222,293(S1)	6		22.2	60.6	97	39
AC 222,293+PO(S1)	4+0.125G	0	28.7	59.5	98	25
AC 222,293+MS(S1)	4+0.125G	1	25.2	58.8	98	46
Diclofop(S2)	16	0	22.2	58.4	61	69
Diclofop+PO(S2)	16+0.125G	1	22.2	58.3	66	71
Diclofop+MS(S2)	16+0.125G	0	23.6	58.3	78	75
AC 222,293(S2)	4	0	22.7	58.6	88	34
AC 222,293(S2)	6	3	23.9	58.6	91	35 48
AC 222,293(S2)	8	0	26.6	59.1	95	46
AC 222,293+PO(S2)	4+0.125G	3	25.3	57.6	90 88	58
AC 222,293+MS(S2)	4+0.125G	0	24.7	58.1	88	70
Difenzoquat+Feno&2,4	-D&MCPA(S2) 5+5.5	5	20.0	58.6	98	5
Difenzoquat(S2)	12	13	20.2	58.3	67	91
HOE-7113-04H(S2)	2.6	0	25.2	58.3		92
HOE-7125-01H(S2)	11	0	23.7	58.9	72 0	0
Untreated	0	0	22.5	59.9	U	0
	Q Q	13	30.1	62.4	98	92
HIGH MEAN		0	20.0	57.6	0	0
LOW MEAN		1	24.2	59.2	81	55
EXP MEAN		226	20.6	2.6	12	35
C.V. %		4	NS	2.1	14	27
LSD 5%		5	NS	2.8	19	36
LSD 1%		4	4	4	4	4
# OF REPS						

a G in the rate column is for gallons per acre.

#### Summary

Wild oats control with diclofop was enhanced by methylated seed oil (MS=Sunit) at the second application and tended to be enhanced by both petroleum oil (PO-11N+17% Atplus 300F) and MS at the first application. Green foxtail control with diclofop was not enhanced by adjuvants. HOE-7113-04H(Puma) and HOE-7125--1H(tiller) gave greater than 90% control of green foxtail. The apparent green foxtail control with AC 222,293 indicates a possible interaction with the chlorsulforon and 2,4-D which was applied for broadleaf weed control. Adjuvants did not enhance wild oats control with AC 222,293. However MS tended to enhance green foxtail control with AC 222,293. Yield was not increased by weed control because wild oats infestation was sparce and the foxtail was partly controlled by the chlorsulfuron applied over the entire plot area.

Wild oats control in Wheat, Minot 1988. 'Stoa' Hard Red Wheat was seeded on April 28. Treatments (S1) were applied to 4-leaf wheat and 2- to 4-leaf wild oats and green foxtail on May 25 with 80 F, and 10 mph wind at application. Treatments (S2) were applied to 5-leaf wheat, 3.5- to 5.5-leaf wild oats, and 3 to 4 inch tall common lambsquarter on June 2 with 70 F, 3 mph wind, and cloudy sky at application. All treatments were applied in 8.5 gpa at 35 psi with a bicycle wheel type plot sprayer to an 8 ft wide area the length of 10 by 25 ft plots. The experiment was a randomized complete block with four replications. Evaluation was on July 19. Green foxtail density was variable and other weeds exceeded 3 plants/sq. ft.

Treatment	Rate	Wh in		Coft	C-1-	V: 7.1	
	(oz/A)	(%		Grft % conti	Colq	Yield (bu/A)	TestWt (1b/bu)
Diclofop(S1) Diclofop+PO(S1) Diclofop+PO(S1) Diclofop+MS(S1) AC 222,293(S1) AC 222,293(S1) AC 222,293+PO(S1) AC 222,293+PO(S1) AC 222,293+MS(S1) Diclofop(S2) Diclofop+PO(S2) Diclofop+MS(S2) AC222293(S2) AC222293(S2) AC222293(S2) AC222293+PO(S2) AC222293+PO(S2) Difenzoquat(S2) Difenzoquat(S2) Difenzoquat(S2) Untreated  HIGH MEAN LOW MEAN EXP MEAN C.V. % LSD 5% LSD 1% # OF REPS	12 16 12+0.125G 12+0.125G 4 5 6 4+0.125G 16 16+0.125G 16+0.125G 4 6 8 4+0.125G 4-0.125G 10 12 2.6 11 0	0 0 0 0 0 1 2 1 1 0 0 1 2 6 1 1 20 38 0 0 0 0 3 1 24 6 8 4	45 63 66 68 92 94 97 97 43 54 56 80 89 96 80 85 92 93 51 14 0	61 79 77 71 11 10 0 60 66 74 0 0 14 0 0 0 65 53 0 79 0 30 46 19 26 4	0 0 0 0 0 0 0 4 13 0 0 0 9 0 23 0 13 0 0 9 0 0 13 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	(bu/A)  11.3 10.6 10.0 11.4 13.1 10.9 12.2 13.5 11.3 10.1 9.8 8.9 8.8 9.8 11.9 8.2 9.2 9.0 9.2 9.3 7.3 13.5 7.3 10.2 24.3 NS NS NS 4	(1b/bu) 56.5 58.0 56.4 57.6 57.6 57.2 56.5 57.8 57.8 57.4 55.5 56.8 57.3 56.9 56.0 57.3 56.9 59.8 55.1 57.0 2.7 NS NS 4

# Summary

Difenzoquat caused visiable injury to the Stoa wheat in this experiment. Petroleum oil with 17% Atplus emulsifier (PO) and methylated seed oil (MS) tended to enhance wild oats and green foxtail control from diclofop. Wild oats control with AC 222,293 tended to be enhanced by both adjuvants with the (S1) applications and only with MS at the (S2) application. Yields were low because of the drought, but generally related to wild oats control.

Wild oats control in wheat, Williston 1988. 'Len' Hard Red Spring wheat was seeded on April 29. Treatments (S1) were applied to 2- to 3-leaf wheat and wild oats, 2- to 3-leaf green foxtail and broadleaf weeds less than 2 inch tall on May 19 with 45 F, 55% relative humidity, clear sky, and a 2 mph wind at treatment and a high of 75 F. Treatments (S2) were to 4-leaf wheat and wild oats, 2- to 6-leaf green foxtail, and broadleaf weed less than 2 inch tall on May 267 with 72 F, 30% relative humidity, and clear sky at treatment and a high of 92 F. Treatments were applied with a plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft wide area the length of 10 by 24 ft plots. The experiment was a randomized complete block with four replications Evaluation was on July 12.

								Wheat	TU
Treatment	Rate	Wioa	Grft	Ruth	KOCZ	Colq	Inj	Yield	TW
	(oz/A)		(%	contr	01)		(%)	(bu/A)(	(lb/bu)
								2.6	F7 0
Diclofop(S1)	12	40	49	0	0	0	0	3.6	57.8
Diclofop (S1)	16	49	61	0	0	0	0	3.9	58.6
Diclofop+PO(S1)	12+0.125G	82	81	0	0	0	0	4.3	58.4
Diclofop+MS(S1)	12+0.125G	86	88	0	0	0	0	3.4	58.7
AC 222,293(\$1)	4	97	0	36	55	27	0	4.8	58.6
AC 222,293(S1)	. 5	96	5	36	36	7	0	4.6	59.5
AC 222,293(S1)	6	99	10	61	75	33	0	4.9	60.1
AC 222,293+PO(S1)	4+0.125G	99	0	60	49	7	0	4.9	60.0
AC 222,293+MS(S1)	4+0.125G	99	36	59	68	3	4	5.3	60.1
Diclofop(S2)	16	30	51	0	0	0	0	3.3	59.0
Diclofop+PO(S2)	16+0.125G	45	74	0	0	0	0	3.6	58.3
Diclofop+MS(S2)	16+0.125G	39	65	0	0	0	0	3.9	68.3
AC 222,293(S2)	4	97	8	0	0	5	0	3.0	58.7
AC 222,293(S2)	6	97	0	23	25	17	0	4.2	59.2
AC 222,293(S2)	8	98	0	38	29	0	0	4.3	59.4
AC 222,293+PO(S2)	4+0.125G	99	13	24	20	33	3	4.3	59.6
AC 222,293+MS(S2)	4+0.125G	97	15	24	5	10	0	4.3	59.2
Difenzoquat(S2)	10	82	10	0	0	22	13	2.7	57.6
Difenzoquat(S2)	12	91	0	0	0	. 0	10	2.4	57.9
HOE-7113-04H(S2)	2.6	80	95	0	0	33	0	4.2	59.0
HOE-7125-01H(S2)	11	5	89	54	69	96	0	3.7	60.0
Untreated	0	0	0	0	0	0	0	3.5	58.8
HIGH MEAN		99	95	61	75	96	13	5.3	60.1
LOW MEAN		0	0	0	0	0	0	2.4	57.6
EXP MEAN		73	34	19	20	13	1	3.9	58.9
C.V. %		26	49	51	120	127	289	26.3	
LSD 5%		26	23	14	33	28	5	1.5	
LSD 1%		35	31	18	44	37	7	NS	
# OF REPS		4	. 4	4	4	4	4	4	1

#### Summary

Len wheat was injured by difenzoquat which tended to reduce yield. However, yield was low regardless of treatments because of drought and high temperature. Wild oats and green foxtail control with diclofop was enhanced by petroleum oil (PO) and methylated seed oil (MS) adjuvants. AC 222,293 gave 95% or more wild oats control regardless of rates or adjuvants. HOE-7113 gave 80% or more wild oats and green foxtail control and HOE-7125 green foxtail control.

Antagonism of wild oats control, Williston-1988. 'Len' Hard Red Spring wheat was seeded on April 29. All treatments, except difenzoquat, were applied to 3.5-leaf wheat and wild oats, 3-to 5-leaf green foxtail, and less than 2 inch tall Russian thistle, kochia, and common lambsquarters on May 24 with 62 F, clear sky and a 2 mph wind at treatment and a high of 89 Difenzoquat treatments were to 4-leaf wheat, 3.5-leaf wild oats, 3-to 6-leaf green foxtail and broadleaf weeds less than 2 inch tall on May 26 with 65 F, 3-to 6-leaf 39% RH, cloudy sky, and 5 mph wind at treatment and a high of 85 F. Treatments were applied in 8.5 gpa at 35 psi to a 8 ft wide area the length of 10 by 24 ft plots. The experiment was a randomized complete block with four replications. Evaluation was on July 12. Infestation of wild oats, kochia, and common lambsquarters was sparse and other weed moderate. Harvest for yield was from 80 sq ft on July 25.

Treatment		Rate	Wina	Gf	Duth	VOC7	Co.1 -	T '	Wheat	
DPX-R9674+X-77 DPX-L5300+X-77 DPX-M6316+X-77 HOE-7125-01H HOE-7125-01H+DPX-R9674 HOE-7125-01H+DPX-M6316 HOE-7125-01H+DPX-M6316 HOE-7125-01H+Chlorsulfuron HOE-7125-01H+Chlorsulfuron HOE-7113-04H HOE-7113-04H+DPX-R9674 HOE-7113-04H+DPX-M6316 HOE-7113-04H+DPX-M6316 HOE-7113-04H+Chlorsulfuron Diclofop+PO Diclofop+DPX-R9674+PO Diclofop+DPX-R9674+MS Diclofop+DPX-R9674+MS Diclofop+DPX-R9674+MS Diclofop+DPX-M6316+MS AC 222,293 AC 222,293+DPX-R9674 AC 222,293+DPX-R9674 Difenzoquat Difenzoquat Difenzoquat+DPX-R9674 Difenzoquat+DPX-R9674 Difenzoquat+DPX-R9674 Difenzoquat+DPX-R9674 Difenzoquat+DPX-R9674 Difenzoquat+DPX-R9674 Difenzoquat+DPX-R9674 Difenzoquat+DPX-R9674 Difenzoquat+DPX-M6316 Untreated C.V. % LSD 5%	12+( 12+( 12+( 12+( 12+(	Rate (oz/A) 0.45+0.25% 0.25+0.25% 1.375+0.25% 1.1 11+0.45 11+0.25 11+0.1 11+0.25 2.6 2.6+0.45 2.6+0.25 2.6+0.1 2.6+0.25 2.6+0.1 2.6+0.1 2.6+0.1 2.6+0.25 0.45+0.125G 0.375+0.125G 0.375+0.125G 0.375+0.125G 0.375+0.125G 0.45+0.125G 0.45+0.125G 0.45+0.125G 0.45+0.125G 0.25+0.125G 0.45+0.125G 0.25+0.125G 0.45+0.125G 0.45+0.25 0.45+0.25	0 0 0 0 8 3 0 4 29 4 56 15 15 24 16 31 65 6 0 40 40 13	Gf (% 21 28 9 65 55 69 68 48 53 70 60 88 40 91 30 25 28 41 11 23 50 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	Ruth cont 97 94 98 95 98 97 99 97 08 98 97 97 81 76 97 81 16	KOCZ (rol)- 99 99 88 35 99 99 99 99 99 99 99 99 99 9	Colq 99 99 72 99 98 98 99 99 99 99 99 99 99 99 99 99	(%) 000000000000000000000000000000000000	Y1d bu/A 5.12 4.7 3.8 4.6 4.3 4.9 4.2 4.2 4.5 4.6 4.9 4.6 4.7 4.6 4.7 1.6 2.7 2.9 5.3 1.2	TW 1b/bu 60 60 60 60 60 61 60 60 61 60 60 60 60 60 60 60 60 60 60 60 60 60

Summary

Wild oats and green foxtail control with HOE-7113-04H, and diclofop was or tended to be reduced when any of the sulfonylurea broadleaf controlherbicides were in the spray mixture. Wild oats control with difenzoquat and green foxtail with HOE-7125-01H also tended to be reduced by the sulfonylureas. Sulfonylurea control of broadleaf weeds generally was not influenced by wild oats control herbicides, except for Russian thistle control with AC 222,293, and difenzoquat with DPX-M6316 for common lambsquarters. Wheat yields were low because of the drought.

Antagonism of Wild oats control, Minot 1988. 'Stoa' Hard Red Spring wheat was seeded on April 28. Treatments were applied to 3.5- to 4-leaf wheat, 2- to 4-leaf wild oats, and 2- to 5-leaf green foxtail on May 25 with 80 F, 10 mph wind, and clear sky at application. All treatments were applied in 8.5 gpa at 35 psi with a bicycle wheel type plot sprayer to an 8 ft wide area the length of 10 by 25 ft plots. The experiment was randomized complete block with four replications. Evaluation was on July 19. All weeds at about 1 plant per sq ft.

A STREET WAS ASSESSED.									
		Wht		0.0		C-1.	- VID	DVC	TW
Treatment	Rate	inj	Wioa	GT_	Rrpw	<u> </u>	(hu/A	DKG	
	(oz/A)	(%)	(%	% CC	ntro	1)	-(bu/A	(%)(	ib/bu)
					0.7	00	0.0	2 0	EO 0
DPX-R9674+X-77	0.45+0.25%	0	6	0	97	98	9.8	3.8	58.8
DPX-L5300+X-77	0.25+0.25%	0	35	14	36	99	11.6	2.4	57.5
DPX-M6316+X-77	0.375+0.25%	0	0	0	97	94	9.5	5.0	58.0
HOE-7125-01H	11	5	46	91	91	99	9.4	3.5	58.5
HOE-7125-01H+DPX-R9674	11+0.45	0	18	62	95	99	8.9	5.7	58.2
HOE-7125-01H+DPX-L5300	11+0.25	1	20	62	96	99	8.5	5.9	58.6
HOE-7125-01H+DPX-M6316	11+0.25	5	28	89	87	99	9.0	6.2	58.5
HOE-7125-01H+Metsulfuron	11+0.1	0	26	68	98	99	12.0	2.9	59.5
HOE-7125-01H+Chlorsulfuron	11+0.25	4	34	30	94	99	10.6	3.9	58.6
HOE-7113-04H	2.6	1	81	94	0	0	9.3	1.3	56.0
HOE-7113-04H+DPX-R9674	2.6+0.45	0	50	66	99	99	13.0	3.0	58.9
HOE-7113-04H+DPX-L5300	2.6+0.25	3	46	63	8	99	10.4	3.0	58.1
HOE-7113-04H+DPX-M6316	2.6+0.25	3	60	86	96	98	11.5	2.7	59.4
HOE-7113-04H+Metsulfuron	2.6+0.1	1	40	74	96	99	12.3	2.6	59.0
HOE-7113-04H+Chlorsulfuron	2.6+0.25	0	48	51	99	98	12.7	2.8	59.7
Diclofop+PO	12+0.125G	3	62	68	0	5	9.1	2.7	56.9
Diclofop+DPX-R9674+PO	12+0.45+0.125	G 3	33	45	92	99	11.6	2.5	59.8
Diclofop+DPX-L5300+PO	12+0.25+0.125		45	36	40	99	11.5	2.9	59.5
Diclofop+DPX-M6316+PO	12+0.375+0.12	5G 0	45	38	96	96	10.0	3.4	59.1
Diclofop+DPX-R9674+MS	12+0.45+0.125		51	51	94	99	11.0	1.6	59.0
Diclofop+DPX-L5300+MS	12+0.25+0.125		31	38	13	99	12.0	2.7	60.0
Diclofop+DPX-M6316+MS	12+0.375+0.12			43	91	99	9.9	3.1	59.9
AC 222,293	6	0		6	20	6	12.3	. 5	57.2
AC 222,293+DPX-R9674	6+0.45	0		8	90	78	11.8	.9	58.9
AC 222,293+DPX-L5300	6+0.25	0		8	0	61	11.7	1.2	58.3
AC 222,293+DPX-M6316	6+0.25	0		10	95	74	12.4	1.0	59.2
Difenzoquat	12	4	87	8	0	0	10.6	2.1	57.0
Difenzoquat+DPX-R9674	12+0.45	6		5	99	98	12.4	2.5	59.3
Difenzoquat+DPX-L5300	12+0.25	3		3		99	10.3	1.6	58.7
Difenzoquat+DPX-M6316	12+0.25	3		4	84	90	10.7	3.3	59.4
Untreated	0	0		0	0	0	9.3	1.8	58.7
On or eaced									
C.V. %		245	30	31	16	6		44.4	1.6
LSD 5%		NS		17		7	NS	1.8	1.3
# OF REPS		1		4	. 3	4	4	4	4
III t III and indication	Wiga-Wilde	atc		Gro	en f	oxta	il.	Cola:	-Common

a Wht=Wheat, inj=injury, Wioa=Wildoats, Gf=Green foxtail, Colq=Common lambsquarter, YLD=Yield, DKG=Dockage, TW=Test Weight;

Summary

All sulfonylurea herbicides generally antagonized wild oats and foxtail control from HOE-7126-01H(Tiller), HOE-7113-04H(Puma), and diclofop. DPX-M6316 generally was the least antagonistic of sulfonylurea, especially when considering antogonism of green foxtail. Wild oats control from difenzoquat tended to be reduced by the sulfonylurea. Common lambsquarters control was less when sulfonylurea herbicides were applied with AC 222,293 than when alone or with other wild oats control herbicides. DPX-L5300 did not adequately control redroot pigweed.

Antagonism of wild oats control, Fargo 1988. 'Wheaton' Hard Red Spring wheat was seeded 2.5 inches deep in 7 inch spaced rows on April 12. Emergence of wheat was poor because of dry soil conditions. Treatments were applied to 4-leaf wild oats and wheat on May 24 with 72 F, 50% RH, and a 3 mph wind. All treatments were applied in 8.5 gpa at 35 psi with a bicycle wheel type plot sprayer to an 8 ft wide area the length of 10 by 30 ft plots. The experiment was randomized complete block with four replications. Evaluation was on June 22 and 27 and wild oats exceeded 10 plants/sq. ft.. Wheat yield or wheat injury ratings were not obtained because drought stress masked any crop injury and what stand was variable.

Treatment	D.	-June 22-	-June 27-
Treatment	Rate	Wild Oats	Wild Oats
	(oz/A)	(% cor	itrol)
DPX-R9674+X-77	0.45+0.25%	Asset Control	Name and State of Sta
DPX-L5300+X-77		4	1
	0.25+0.25%	10	20
DPX-M6316+X-77	0.375+0.25%	4	0
HOE-7125-01H	11	60	44
HOE-7125-01H+DPX-R9674	11+0.45	23	14
HOE-7125-01H+DPX-L5300	11+0.25	29	18
HOE-7125-01H+DPX-M6316	11+0.25	54	43
HOE-7113-04H	2.6	85	79
HOE-7113-04H+DPX-R9674	2.6+0.45	69	61
HOE-7113-04H+DPX-L5300	2.6+0.25	64	50
HOE-7113-04H+DPX-M6316	2.6+0.25	76	70
Diclofop+PO	12+0.125G	30	20
Diclofop+DPX-R9674+PO 12	2+0.45+0.125G	20	
	2+0.25+0.125G	24	21
	2+0.375+0.125G	38	15
	2+0.45+0.125G		29
	2+0.25+0.125G	28	16
	2+0.375+0.125G	28	16
AC222293		40	13
AC 222,293+DPX-R9674	6	89	81
AC 222,293+DPX-L5300	6+0.45	89	79
AC 222,293+DPX-M6316	6+0.25	87	77
Diforzoguat	6+0.25	85	73
Difenzoquat	12	95	84
Difenzoquat+DPX-R9674	12+0.45	94	83
Difenzoquat+DPX-L5300	12+0.25	93	84
Difenzoquat+DPX-M6316	12+0.25	96	88
C.V. %		19	25
LSD 5%			25
# OF REPS		14	16
		4	4

## Summary

Wild oats control with HOE-7125-01H and HOE-7113-04H was antagonized by DPX-R9674 and DPX-L5300 and only tended to be antagonized by DPX-M6316. Wild oats control with AC 222,293 or difenzoquat was not antogonized by the three DPX compounds. Diclofop gave too little wild oats control for detection of antagonism from the DPX compounds in the experiment.

AC 222,293 plus bromoxynil and surfactant. Fargo 1988. 'Wheaton' Hard Red Spring wheat was seeded 2.5 inch deep in 7 inch spaced rows on April 12. Emergence of wheat was poor because of dry soil conditions. Two experiments were conducted, one with all treatments and one half the bromoxynil rate of that used in experiment 2. Treatments were applied to 4-leaf wheat and wild oats on May 23 with 72 F, 70 RH, and 3 mph wind at application for the complete experiment. The second experiment was to 5-leaf wheat and wild oats on May 26 with 70 F, 65% RH and 8 mph wind. All treatments were applied in 8.5 gpa at 35 psi with a bicycle wheel type plot sprayer to an 8 ft wide area the length of 10 by 30 ft plots. The experiment was randomized complete block with four replications. Evaluation of the complete experiment was on June 27 and experiment 2 on June 22. Wild oats density exceeded 3 plants/sq ft in both experiments and wheat was drought stressed and variable.

					-Exp. 2
[reatment	Rate	Brox rate	Wild Oats	Wht	Wild Oats
		Exp. 2	control	inj	control
	(oz/A)			(%)	
AC 222,293	5		78	0	90
AC 222,293+X-77	5+0.25%	6	84	1	98
Bromoxynil	2	4	3	0	0
AC 222,293+Brox	5+1	2	73	0	90
AC 222,293+Brox	5+1.5	3	73	3	90
AC 222,293+Brox	5+2	4	67	1	96
AC 222,293+Brox+X-77	5+1+0.25	5% 2	81	0	94
AC 222,293+Brox+X-77	5+1.5+0		77	1	93
AC 222,293+Brox+X-77	5+2+0.2!		70	1	94
AC 222,293+Brox+X-77	5+1+1%	2	75	0	96
AC 222,293+Brox+X-77	5+1.5+	1% 3	72	3	95
AC 222,293+Brox+X-77	5+2+1%	4	71	0	96
AC 222,293+24-Ddea	5+8		44	-	
AC 222,293+24-Ddea+Na	HC03 5+8+24		58	-	wo
AC293+24-Ddea+NaHC03+			48	-	-
AC293+24-Ddea+AMS	5+8+32		31	-	
AC 222,293+NaHC03	5+24		74	-	
AC 222,293+AMS	5+32		75	-	
AC 222,293+24-Dbee	5+8		68	-	-
Untreated	0		1		
C.V. %			10	271	4
LSD 5%			9	NS	3
# OF REPS			4	4	4

Summary

Bromoxyonil (Buctril) did not antagonize wild oats control with AC 222,293 in either experiment. However wild oats control tended to decrease as bromoxyonil rate increased regardless of surfactant in the complete experiment. Wild oats control was more then 90% with all treatment in experiment 2. Wild oats control with AC-222293 was antagonized by 2,4-D(dea), but not by sodium bicarbonate, ammonium sulfate(AMS), or 2,4-D butoxyethyl ester(bee). Sodium bicarbonate and/or AMS did not overcome the antogonism caused by 2,4-Ddea.

Diclofop plus adjuvants, Fargo 1988. 'Wheaton' Hard Red Spring wheat was seeded 2.5 inch deep in 7 inch spaced rows on April 12. Emergence of wheat was poor because of dry soil conditions. Treatments were applied to 4-leaf wheat and wild oats on May 23 with 72 F, 70% RH, and a 3 mph wind at application. All treatments were applied in 8.5 gpa at 35 psi with a bicycle wheel type plot sprayer to an 8 ft wide area the length of 10 by 30 ft plots. The experiment was randomized complete block with four replications. Evaluations were on June 27. Wheat injury was not evaluated because drought stress masked any treatment effects and what stand was variable.

Treatment	Data	Wild Oats
	(OZ/A)	(%)
Diclofop Diclofop Diclofop Diclofop Diclofop+PO Diclofop+PO Diclofop+PO Diclofop+MS Diclofop+MS Diclofop+MS Diclofop+MS Diclofop+AG-2 Diclofop+AG-2 Diclofop+AG-2 Untreated  HIGH MEAN LOW MEAN EXP MEAN	Rate (oz/A)  8 10 12 16 8+0.12G 10+0.12G 12+0.12G 16+0.12G 8+0.12G 12+0.12G 16+0.12G 16+0.12G 10+0.12G 10+0.12G 10+0.12G 10+0.12G 10+0.12G 10+0.12G 10+0.12G	control (%) 9 10 36 43 30 48 50 55 33 39 51 66 44 33 45 53 0
C.V. %		34
LSD 5%		
LSD_1%		19 25
# OF REPS		4

## Summary

The three adjuvants all enhanced wild oats control with diclofop. The petroleum oil with 17% Atplus (PO) was simuliar to the methylated seed oil(MS=Sun-it). AG-2 adjuvant caused diclofop to give simuliar wild oats control at all diclofop rates. Averaged of one diclofop rates wild oats control was 24, 46, 47, and 44 % with none, petroleum oil, methylated seed oil, and AG-2.

Wild oats and broadleaf herbicide combinations, Fargo 1988. 'Wheaton' Hard Red Spring wheat was seeded 2.5 inch deep in 7 inch spaced rows on April 12. Emergence of wheat was poor because of dry soil conditions. Treatments were applied to 4-leaf wheat and wild oats on May 24 with 70 F, 50% RH and 3 mph wind at application. All treatments were applied in 8.5 gpa at 35 psi with a bicycle wheel type plot sprayer to an 8 ft wide area the length of 10 by 30 ft plots. The experiment was randomized complete block with four replications. Evaluation was on June 22. Wild oats density exceeded 3 plants/sq ft and wheat stand was variable and drought stressed.

Twoatmont	Rate	Wild Oats
<u>Treatment</u>	(oz/A)	(% control)
Diclofop	12	34
Diclofop+Brox&MCPA	12+17.5	26
Diclofop+Acif	12+4	25
Diclofop+Fluroxypyr	12+2	35
Diclofop+BAS514	12+3	36
AC222293	5	87
AC222293+Clopyralid	5+1.5	89
AC222293+C1py&2=4-D	5+9.5	73
AC222293+Fluroxypyr	5+2	88
AC222293+Acif	5+4	61
AC222293+BAS514	5+3	73
Difenzoquat	10	69
Difenzoquat+Clopyralid	10+1.5	87
Difenzoquat+Clpy&2=4-D	10+9.5	80 76
Difenzoquat+Acif	10+4	76
Difenzoquat+BAS514	10+3	76 86
Difenzoquat+Fluroxypyr	10+2	0
Untreated	0	O .
C 11 0/		20
C.V. %		17
LSD 5%		23
LSD 1% # OF REPS		4
# UI NEFS		

#### Summary

None of the treatment cause any visable injury to wheat. Wheat was not harvested for grain yield because of drought stress and a variable stand. Wild oat control with diclofop was low which probably masked any possible antagonism from the broadleaf control herbicides. AC 222,293 was or tended to be antagonized by clopyralid plus 2,4-D, aciflurofen, and BAS-514; but not by clopyralid alone or fluroxypyr. Wild oats control with difenzoquat was simuliar in mixture with all broadleaf control herbicides, but tended to be higher then when difenzoquat was applied alone. The low control of wild oats with difenzoquat alone may have been a chance occurance because of an only 40% value in replication three.

Antagonism of diclofop by various compounds, Fargo 1988. 'Wheaton' Hard Red Spring wheat was seeded 2.5 inch deep in 7 inch spaced rows on April 12. Emergence of wheat was poor because of dry soil conditions. Treatments were applied to 4-leaf wheat and wild oats on May 20 with 65 F, 60% RH, and 15 mph wind at application. 2,4-Ddea was diethanolamine and 2,4-Dbee was butoxyethyl ester. All treatments were applied in 8.5 gpa at 35 psi with a bicycle wheel type plot sprayer to an 8 ft wide area the length of 10 by 30 ft plots. The experiment was randomized complete block with four replications. Evaluation was on June 27. Wild oats density exceeded 2 plants/sq ft and wheat stand was variable because of drought stress.

Treatment	Rate (oz/A)	Wild oats (% control)	Wht inj (%)	
Diclofop Diclofop+28%N Diclofop+24-Ddea Diclofop+24-Ddea+28%N Diclofop+24-Dbee Diclofop+24-Dbee+28%N Diclofop+NaHCO3 Diclofop+NaHCO3+28%N Diclofop+24-Ddea+NaHCO3 Dicl+24-Ddea+NaHCO3+28%N Dicl+24-Dbee+NaHCO3 Dicl+24-Dbee+NaHCO3	12+8+16	53 58 14 3 1 9 39 47 18 11 1 6	0 0 0 0 0 0 0 0 0	
HIGH MEAN LOW MEAN EXP MEAN C.V. % LSD 5% LSD 1% # OF REPS		58 0 20 37 11 14 4	0 0 0 0 NS NS	

#### Summary

Wild oats control level was low reducing the potential for antagonism by the various compounds. 2,4-Ddea or 2,4-Dbee antogonism of diclofop was not overcome by the 28% N fertilizer in the spray mixture. Sodium bicarbonate antagonized wild oats control with diclofop and the antagonism was overcome by 28 % N fertilizer. Sodium bicarbonate did not influence the antagonism caused by 2,4-Ddea or bee.

Evaluation of herbicides for wild oats control, Fargo 1988. 'Wheaton' Hard Red Spring wheat was seeded 2.5 inch deep in 7 inch spaced rows on April 12. Emergence of wheat was poor because of dry soil conditions. The 2-3 lf treatments were applied to 3-leaf wild oats and wheat on May 16 with 55 F, 40% RH, no wind and clear sky at application. The 5-leaf treatments were applied on May 26 with 70 F, 65% RH, and 5 mph wind. All treatments were applied in 8.5 gpa at 34 psi with a bicycle wheel type plot sprayer to an 8 ft wide area the length of 10 by 30 ft plots. The experiment was randomized complete block with four replications. Evaluation was June 27. Wild oats density exceeded 5 plants/sq ft and wheat stand was variable and drought stressed which prevented accurate evaluation.

Treatment	Stage	Rate (oz/A)	Wild Oats (% control)
Diclofop+COC Diclofop&Brox&MCPA HOE-7125-01H HOE-7113-04H AC 222,293 RH-0898+PO RH-0898+PO RH-0898+MS RH-0898+MS RH-0898+AG-2 RH-0898+PO RH-0898+PO RH-0898+PO RH-0898+PO RH-0898+PO RH-0898+MS HOE-7125-01H HOE-7113-04H Feno&24-D&MCPA Feno&24-D&MCPA Difenzoquat Untreated	(2-31f) (2-31f) (2-31f) (2-31f) (2-31f) (2-31f) (2-31f) (2-31f) (2-31f) (2-31f) (2-31f) (2-31f) (51f) (51f) (51f) (51f)	12+0.12G 17.4 11 2.6 6 1.2+0.12G 1.2+0.25G 1.2+0.12G 1.2+0.12G 1.6+0.12G 1.6+0.12G 11 2.6 10.6 12.5 10	40 45 77 97 96 41 73 69 89 30 51 78 48 61 45 61 92 0
C.V. % LSD 5% LSD 1% # OF REPS			13 12 15 4

## Summary

HOE-7113-04H at 2.6 oz/A gave greater wild oats control then HOE 7125-01H at 11 oz/A at both stages of application. The methylated seed oil (MS-Sun-it) enhanced wild oats control with RH-0898 more than the petroleum oil adjuvant (PO-Atplus 411F).

Adjuvants with AC 222,293, Williston 1988. 'Len' Hard Red Spring wheat was seeded on April 29. Treatments were applied to 3.5-leaf wheat, 3-leaf wild oats, 2- to 5-leaf green foxtail, and less than 2 inch tall broadleaf weeds on May 23 with 70 F, 13% RH, clear sky, and a 6 mph wind. The herbicides were applied with a tractor-mounted plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft wide area the length of 10 by 25 ft plots. The experiment was a randomized complete block with four replications. Evaluations were on July 13. Wild oats, kochia, and common lambsquarters infestations were sparce, green foxtail moderate, and Russian thistle dense.

Treatment	Rate	Wioa	Grft	Ruth	KOCZ	Colq	ini	Tw Yield
	(oz/A)		(%	conti	rol)-			(1b)(bu)
AC 222,293 AC 222,293+LI-700 AC 222,293+MS AC 222,293+AG-2 AC 222,293+PO AC 222,293+2,4-Ddea+NaHCO3 AC 222,293+2,4-Ddea+NaHCO3+AMS AC 222,293+2,4-Ddea AMS AC 222,293+NaHCO3 AC 222,293+AMS AC 222,293+AMS AC 222,293+AMS	5 5+0.25G 5+0.25G 5+0.25G 5+0.25G 5+8 5+8+16 5+8+16+32 5+8+32 5+16 5+32 5+32+16 0	94 94 94 94 94 88 89 88 64 95 95	21 13 0 6 0 5 15 8 0 0 0	58 60 56 50 56 89 50 91 80 69 63 61	67 38 0 0 38 85 83 70 90 63 48 68	43 39 0 0 5 95 93 89 95 76 44	1 3 3 5 1 1 0 0 0 1 3 3	60.9 5.4 60.5 5.0 60.6 5.1 59.8 4.3 60.3 4.9 61.1 5.2 60.5 5.7 60.9 5.3 60.7 6.0 60.5 5.2 60.2 5.3 60.3 5.9
HIGH MEAN LOW MEAN EXP MEAN C.V. % LSD 5% LSD 1% # OF REPS		95 0 83 12 15 20 4	21 0 5 11 NS NS 4	91 0 59 30 26 34 4	90 0 50 47 51 NS 2	23	5	60.4 5.4 61.1 6.0 59.8 4.3 60.5 5.3 19.7 NS NS 1 4

#### Summary

2,4-D + ammonium sulfate (AMS) reduced and 2,4-D alone or with sodium bicarbonate and ammonium sulfate tended to reduce wild oats control with AC 222, 293. Thus, the sodium bicarbonate overcame the antagonism of the ammonium sulfate with 2,4-D. Sodium bicarbonate with 2,4-D reduced Russian thistle control, but ammonium sulfate overcame the antagonism. 2,4-D with AC 222, 293 enhanced kochia and common lambsquarter control, regardless of other salts present.

Wild oats control evaluation, Langdon 1988. 'Coteau' Hard Red Spring wheat was seeded on May 6. The first treatments (2) were to 3 -leaf wheat and 1 to 3 leaf wild oats and green foxtail on May 26. The second treatments (4) were to 4 to 5 leaf wheat and wild oats on June 1. The entire plot area was treated with chlorsulfuron at 0.19 oz/A + 2,4-D at 4 oz/A on June 12. Treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi through 8001 nozzles to an 8 ft wide area the length of 10 by 25 ft plots. The experiment was a randomized complete block with four replications. Evaluations were on July 20.

					Whe	
Treatment	Rate	Inj	Wioat	Grft	Yield	Tswt
Treatment	(1b/A)	(%)	(% con	trol)	(bu/A)	(1b/bu)
				•	12 E	59.4
Untreated		0	0	0 92	13.5 16.6	58.9
HOE-7125 (41f)	0.66	1	59	96	19.1	58.5
HOE-7125 (41f)	0.78	0	71 92	95	21.8	59.9
HOE-7113 (21f)	0.16	0	64	95	16.4	58.8
HOE-7113 (41f)	0.16 0.12	0	56	59	16.7	58.8
Diclofop&Brax&MCPA (21f)	0.75+1pt	1	72	77	21.7	58.6
Diclofop+PO (21f) Diclofop+DPX-M6316+PO (21f)	0.75+0.016+1pt	1	80	68	18.0	59.0
Difenzoquat (41f)	1.0	13	97	0	14.1	57.8
AC 222,293 (21f)	0.38	3	99	58	22.0	59.5
HOE-7125 (41f)	0.33	0	40	81	16.2	58.9
HOE-7125 (41f)	0.39	1	59	92	17.7 20.6	58.5 58.3
HOE-7113 (21f)	0.08	0	55 38	83 93	16.9	57.6
HOE-7113 (41f)	0.08	0	30	33	10.5	37.0
WEAR		13	99	96	22.0	59.9
HIGH MEAN		0	0	0	13.5	57.6
LOW MEAN EXP MEAN		1	63	71	17.9	58.7
C.V. %		147	19	13	15.9	1.4
LSD 5%		3	17	13	4.1	1.2
LSD 1%		4	23	17	5.5	NS 4
# OF REPS		4	4	4	4	4

#### Summary

Wheat yield was not influenced by the degree of wild oats or green foxtail control. Green foxtail control was greater with HOE-7125 or HOE-7113 than with diclofop; regardless of stage at application. Wild oats control exceeded 90% with HOE-7113, AC 222,293, and diffenzoquat.

Wild oat control in durum wheat, Minot 1988. Postemergence herbicides were evaluated for wild oat and green foxtail control in durum wheat. 'Lloyd' durum was seeded April 29 in Williams loam soil with pH 6.6 and 4% organic matter. Treatments were applied with a bicycle wheel plot sprayer delivering 8.5 gpa at 35 psi on May 23 (S1) to 3 to 3.3 leaf wheat, 1 to 3 leaf green foxtail and wild oats with sunny sky and 80 F and on June 1 (S2) to 5 to 5.5 leaf wheat, 3 to 6 leaf green foxtail and 3 to 5 leaf wild oats with cloudy sky and 70 F. Green foxtail was drought stressed at the time of treatment. Initial rainfall after treatment was 1.06 inches on June 14 and only 3.39 inches of rain fell from May 23 to July 20. Wheat injury and weed control evaluations were made on June 23 (Eval 1) and July 21 (Eval 2). The experiment was a randomized complete block design with four replicates and experimental units were 10 X 18 sq ft. Wheat was harvested from a 4 X 18 sq ft. area on July 28.

-		D	urum wh	Injury	Weed of Wioa	control Grft
Treatment	Rate	Yld	Tswt	1 2	1 2	1 2
	(1b/A)	(bu/A)	(1b/bu)		(%)-	
Diclofop+P0 Dicl+DPX-M6316+P0 Dicl&Brox&MCPA Imazamethabenz Imazamethabenz H0E7113-04H H0E7113-04H H0E7125-02H H0E7125-02H Control	.75+.125G .016+.125G .+0.25+0.05 0.25 0.38 0.16 0.16 0.66 0.78	8.4 8.6 6.9 7.7 10.8 6.9 6.7 5.7 4.6 5.5	57.9 60.0 60.3 59.4 58.6 58.7 58.9 56.5 58.1	6 1 8 6 3 0 1 3 4 3 18 8 10 5 10 8 14 14 34 35 0 0	79 76 76 65 46 34 99 94 99 98 85 71 71 50 43 15 58 29 90 81 0 0	41 30 29 16 6 9 20 0 10 0 68 64 16 18 11 16 16 19 4 0
C.V. % LSD 5% F-TRT		23.0 2.3 4.9	1.0 1.3 7.3	66 83 9 9 9 11	20 16 20 13 19 52	56 41 16 9 12 35

a PO = Paraffin base petroleum oil with 17% surfactant blend; Dicl = diclofop; Brox = bromoxynil; & = formulated mixture which was 1:0.31:0.06 for diclofop & bromoxynil & MCPA;

#### Summary

'Lloyd' durum wheat was injured initially by HOE 7113-04H, HOE 7125-02H, and difenzoquat, however, only HOE 7125-02H and difenzoquat caused wheat injury greater than 10% by the July 21 evaluation. Wheat treated with HOE 7125-02H and difenzoquat yielded less than all other treatments and 5.1 to 6.2 bushels less than the wheat treated with imazamethbenz at 0.38 lb/A. Difenzoquat treated wheat had the lowest test weight, 56.5 lb/bu, indicating crop injury which influenced the grain filling process. Green foxtail control was reduced by heat and drought stress and was generally poor with all treatments. The second wild oat control evaluation was lower than the first indicating an incomplete kill of the wild oats with all treatments except imazamethabenz. Diclofop & bromoxynil & MCPA and HOE 7125-02H gave less than 35% wild oat control on the second evaluation.

Wild oats herbicides with adjuvants, -1-Prosper 1988. 'Wheaton' HRS wheat, ND810104' oats, and siberian foxtail millet were seeded as bioassay species in adjacent 6 ft wide strips at Prosper, ND on May 23. Treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 34 psi to an 8 ft wide area across the bioassay species. Plots were 10 ft wide to provide a 4 ft untreated area between treated areas to aid evaluation. Treatments were applied to 5-to 6-leaf wheat, oats, and foxtail millet on June 20 with 95 F, 50% RH, and clear sky. Evaluations were on July 22 and August 11. Data was not taken on injury to wheat because none was obvious.

		Jul	y 22	Augus	
Treatment	Rate	Ftmi	Oats	Fxtl	Oats
	(%)	(a) (c) (a) (a) (a) (a) (a) (a) (a)	(% 0	control)	
Diclofop	12	47	39	32	36
Diclofop+PO	12+0.12G	57	46	36	37
Diclofop+MSC	12+0.12G	53	54	35	54
Diclofop+AG-2	12+0.12G	58	52	43	40
Diclofop+28%	12+1G	49	52	35	34
Diclofop+SCI-40	12+0.25%	54	48	42	41
Diclofop+Inhance	12+0.12G	47	31	31	26
Diclofop+LI-700	12+0.25%	44	32	29	35
Diclofop+X-77	12+0.25%	53	56	34	37
Untreated	0	0	0	0	0
AC 222,293	4	3	85	0	96
AC 222,293+P0	4+0.12G	10	92	13	97
AC 222, 293+MSC	4+0.12G	11	91	5	97
AC 222,293+AG-2	4+0.12G	13	88	6	94
AC 222, 293+28%	4+1G	9	89	9	98
AC 222, 293+SCI-40	0.25%	6 .	89	0	95
AC 222, 293+Inhand		6	86	5 3 3	95
AC 222, 293+LI-700		4	83	3	96
AC 222,293+X-77	4+0.25%	6	90		99
Untreated	0	0	0	0	0
Difenzoquat	10	6	80	9	96
Difenzoquat+PO	10+0.12G	6	76	9	90
Difenzoquat+MSC	10+0.12G	10	81	4	95
Difenzoquat+AG-2	10+0.12G	10	80	6	96
Difenzoquat+28%	10+1G	10	81	3	96
Difenzoquat+SCI-		8	82	5	94
Difenzoquat+Inham		6	82	0	96
Difenzoquat+LI-7	00 10+0.25%	13	82	9	95
Difenzoquat+X-77	10+0.25%	5	80	1	96
Untreated	0	0	0	0	0
C.V. %		28	11	45	9
LSD 5%		8	10	8	9
# OF REPS		4	4	4	4

#### Summary

Diclofop control of oats and foxtail millet was low probably because of the drought and the advanced growth stage at treatment. However control of oats and foxtail millet tended to be enhanced by both petroleum oil with 17% Atplus 300 F (PO) and methylated seed oil with 15% emulsifier (MSC). AC-222293 gave a high degree of oats control, with certain adjuvants tended to enhance control at the early evaluation. Difenzoquat control of wild oats was not influenced by adjuvants.

Hard Red Spring wheat cultivar response to herbicides, Minot 1988. Triallate at 1.0 lb/A was applied to an 80 inch band of moist soil and incorporated with a rototiller and cultiweeder and Hard Red Spring wheat varieteis were seeded 2.5 inches deep on April 21. Wheat stand reduction was evaluated on May 19. One shot at 3.5 pts/acre was applied for weed control on May 21. Difenzoquat at 1.0 lb/A, DPX-L5300 at 0.5 oz ai/A. and DPX-R9674 at 0.9 oz ai/A were applied to 5.0 leaf wheat on June 1, with a cloudy sky and 70 F. Metsulfuron at 0.06 oz ai/A was applied to 5.2 leaf wheat on June 3, with a sunny sky and 90 F. Difenzoquat injury was evaluated on June 17 and all the postemergence herbicide injury on July 8. Data are for unreplicated cultivars.

		Difenzoquat	DPX-	DPX-	Motaul
Cultivar	Triallate	June 17 July 8		R9674	Metsul-
	(% Std red)		Hard Red S	nring wh	furon
Alex	20	35 10	0		
Kenyon	10	5 0	0	0	0
Roblin	0	25 20	0	0	0
Columbus	0	5 0	0	5	0
Lew	0	75 25	0	0	0
Coteau	0	10 5	0	0	5
Stoa	5	5 5	0	0	0
Butte 86	0	10 10	0		0
Amidon	10	25 10	5	.0	0
Laura	0	35 10		0	0
Baart	Ö		0	0	. 0
Cutless	15		5	0	0
Rambo	0	20 5 10 0	10	5	5
Len	Ö	75 15	0	0	0
Marshall	15		5	0	0
Wheaton	0		10	0	0
Prospect	20	5 0 20 10	10	5	5
2369	0	20 5	10	0	0
2385	Ö	20 30	5	- 0	0
Norak	Ö	25 15	10	10	0
Leif	Ö	35 15	15	0	15
Celtic	Ö	75 20	5 0	0	0
Norseman	Ŏ	55 15	10	0	0
Nordic	Ō	15 5		0	0
Telemark	5	20 5	10	0	0
Tammy	Ö	5 0	5 5	0	0
Westbred 926	Ō	80 90	0	0	5
ND640	5	15 0	0	0	0
ND651	5 5	5 0	0	0 5	0
ND652	0	15 5	0		5
ND618	20	20 5	0	0	0
ND626	0	50 15	0	0	0
ND649	10	25 5		0	0
ND650	5	15 5	0	0	0
Mn81110	10	5 0			0
Mn82354	0	5 0	0	0	0
2375	5	40 50	0 5	0	0
W2501	5	35 15		0	0
W2502	25	15 5	0	10	0
HS84-873	5	35 0	5 5	5 5	0
		Summary	3	3	0
Triallato co	mand and the table of	Juninary			

Triallate caused an initial stand reduction of some varieties however these stand reductions were not detectable at the later evaluation. Cultivars with greater than 20% injury from differzoquat at the late evaluation were Lew,

2385, Westbred 926, and 2375. Sulfonyl urea herbicides did not cause important injury to Hard Red Spring wheat cultivars.

Oats cultivar response to sulfonyl urea herbicides, Minot 1988. Oat cultivars were seeded on April 21. Bromoxynil at 4 oz/A + MCPA at 4 oz/A were applied for weed control on May 21. DPX-L5300 at 0.5 oz ai/A and DPX-R9674 at 0.9 oz ai/A were applied on 8.5 gpa at 35 psi to 5.0 leaf oats on June 1 with a cloudy sky and 70 F. Metsulfuron at 0.06 oz ai/A was applied to 5.2 leaf oats on June 3 with a sunny sky and 90 F. Oat injury evaluations were on June 17 and July 22. Data are for observations of injury to cultivars which were not replicated.

GBC 200 AVEN	npy-	=L5300	DPX-R	9674	Metsu	lfuron
Cultivar	June 17	July 22		July 22	June 17	July 22
OUTCIVAL			(% injury			
Border	60	10	10	0	25	0 5 15
Monida	75	10	25	5	25	5
Otana	65	10	10	5	30	15
Moore	65	15	20	5	35	15
Proat	75	20	35	0	30	0
Pierce	80	20	40	0	35	25 10
Steele	60	15	45	5	35 45	15
Valley	75	15	60	0	40	10
Tibor	60	15 15	40 35	0	40	10
Robert	65 55	20	30	0	50	10
Riel	55	20	30	Ö	35	10
Dumont Fidler	50	15	25	Ö	30	10
Kelsey	50	15	25	0	25	10
Hytest	55	15	30	0	25	10
Sandy	65	15	30	0 -	40	5
Porter	85	40	75	20	80	35
ND810104	60	20	15	0	50	10
Mean	64	16	32	2	37	11

# Summary

Oat cultivars were initially injured by all herbicides tested especially DPX-L5300. The July 22 evaluations indicate that the cultivars had recovered considerably from the initial injury. Porter appears to be the most susceptable cultivar to the sulfonyl urea herbicides.

Durum wheat variety response to herbicides Minot 1988. Triallate at 1.0 lb/A was applied to an 80 inch band of moist soil and incorporated with a rototiller and a cultiweeder and durum wheat varieties were seeded 2.5 inches deep on April 21. Durum stand reduction was evaluated on May 19. One shot (diclofop & bromoxynil & MCPA) at 3.5 pts/acre was applied for weed control on May 21. Difenzoquat at 1.0 lb/A, DPX-L5300 at 0.5 oz ai/A, and DPX-R9674 at 0.9 oz ai/A were applied to 5-leaf durum on June 1 with a cloudy sky and 70 F. Metsulfuron at 0.06 oz ai/A was applied to 5.2 leaf wheat on June 3 with a sunny sky and 90 F. Difenzoquat injury was evaluated on June 17 and all the postemergence herbicide injury evaluations were on July 8. The data represent visual evaluations of cultivars which were not replicated.

Variety	Triallate	<u>Difenz</u> June 17	oquat July 8	DPX-L5300	DPX-R9674	Metsulfuron
	(std.red.)			(% injury	/)	
Rugby	5	20	5	5	5	0
Ward	0	10	0	5	5	ŏ
Arcola	5	25	10	0	Ō	ő
Kyle	0	70	40	0	0	ő
Medora	0	20	10	10	5	ő
Sceptre	0	15	5	10	0	Ö
Monroe	10	15	10	5	0	Ö
Renville	0	10	5	0	0	0
Vic	0	20	10	15	0	Ö
Fjord	0	40	20	0	0	
Stockholm	5	5	0	15	5	5
Laker	0	60	15	10	5	5
Lloyd	0	10	0	25	5	Ö
Cando	0	5	5	0	5	5
D8261	5	20	10	35	10	5
D8291	0	15	10	5	0	0
D8370	10	15	10	5	5	0
D8374	0	40	15	20	5	0
D8380	10	30	5	10	0	0
D83103	0	25	5	0	0	0
D84130	5	25	10	0	0	5
D84134	0	5	5	0	5	5
D8302	10	5	15	5	5	5
D8434	0	20	5	25	20	10
D8458	0	15	5	0	5	0
D8459	5	20	5	0	5	0
D8460	5	10	5	0	5	0
D8475	0	10	0	5	10	0
D8479	0	10	0	5	5	0
FA883-323	5	15	5	5	0	0
Mean	3	20	8	7	4	2

#### Summary

Triallate caused a slight stand reduction of certain durum varieties, however these reductions were not visable late in the season. Difenzoquat caused injury early to many varieties but most varieties recovered with time. Kyle and Fjord were the only varieties to have 20% or greater injury on July 8. DPX-L5300 caused injury of 20% or greater on Lloyd, D8261, D8374, and D8434. D8434 was injured 20% by DPX-R9674 while all other varieties were generally unaffected by DPX-R9674 and Metsulfuron.

Wheat cultivar response to herbicides, Williston 1988. Hard Red Spring wheat varieties were seeded on April 25, durum varieties on April 26. Difenzoquat was applied at 1 lb ai/A to wheat that was in the 3.5 to 4 leaf stage of growth on May 23. Difenzoquat was applied with a tractor mounted sprayer delivering 8.5 gpa at 30 psi. Weather conditions at treatment were as follows: wind SSE at 6 mph, 71 F, 14% RH, clear sky, soil dry. Experiment was on a max loam soil having a PH of 6.4 and an organic matter content of 2.1%. Evaluation was late in June. Severe drought tended to mask the effect of herbicide.

		Difenzo	oquat		
Wheat	Crop	Wheat	Crop	Wheat	Crop
cultivar	injury	cultivar	injury	cultivar	injury
HRS	(%)	HRS	(%)	HRS	(%)
Waldron	15	Coteau	0	Alex	0
Butte 86	0	Stoa	0	Columbus	0
Kenyon	0	Amidon	0	Robin	20
Laura	0	Cutless	0	Lew	0
Lancer	0	Leader	10	Fortuna	0
Thatcher	0	ND 640	0	ND 651	10
ND 652	10	ND 618	0	ND 626	15
ND 649	0	ND650	0	Len	30
Marshall	0	Wheaton	0	2369	20
2385	10	2375	0	Norak	10
Leif	0	Norseman	20	Celtic	0
Nordic	0	Telemark	10	Tammy	10
Prospect	0	Westbred 926	70	Rambo	10
Newana	0	Glenmam	0	MT 7926	0
MT 8402	10	MN 81110	10	MN 82354	10
W 2501	0	W 2502	0	HS 84-873	0
Durum		Durum		Durum	
Ward	0	Crosby	0	Rugby	10
Renville	10	Vic	30	Monroe	20
Medora	10	Septre	10	Kyle	15
Arcola	15	Fjord	0	Cando	0
Lloyd	20	Laker	10	Stockholm	15
D 8261	10	D 8291	0	FA 883-323	0
D 8302	10	D 8370	10	D 8374	30
D 8380	10	D 83130	0	D 8434	30
D 8458	15	D 8459	15	D 8460	25
D 8475	15	D 8479	40	D 84130	30
D 84134	10	Wakooma	20		

#### Summary

Plants were drought stressed which reduced the response to difenzoquat, but injury symptons occurred for certain cultivars as shown in the table. DPX-L5300 and DPX-R9674 were also applied to all the above cultivars of Hard Red Spring wheat and durum on the same day, with the same equipment and weather conditions, and the same growth stages at the difenzoquat application. DPX-L5300 was applied at 0.5 oz ai/A and DPX-R9674 applied at 0.9 oz ai/A. DPX-L5300 appeared to have delayed maturity slightly on only Kyle durum and ND 649 and Rambo HRS wheat. DPX-R9674 caused no visible crop injury to any cultivar.

Small grain response to herbicides, Fargo 1988. The treatments consisting of commercial triallate plus trifluralin granules (Buckle) were applied and once incorporated on October 26, 1987. Treatments were applied with Valmar air flow granular applicator. "Wheaton', "Alex', Alex treated with a protectant from monsanto (MON 5500), and "Coteau' Hard Red Spring wheat; "Rugby' Durum wheat and "Robust' barley were seeded on April 15, 1988. The plot area was treated with 2,4-D at 6 oz/A to control broadleaf weeds when the crops were tillering, June 6. The area was free of wild oats and only had a sparce foxtail infestation. Injury evaluations were on June 22. The herbicides were applied to 20 by 30 ft plots and the crops were seeded across the plots.

Treatment	(oz/A)		Alx	AlxTr (% in	<u>Cot</u>	Ruc	Rbt	Wm	A1x	<u> </u>	<u>Cot</u> /A)	Rug	Rbt
Trif+Triallate Trif+Triallate Trif+Triallate Untreated	6.7+20	9 21 50 0	20 43 78 3	10 23 49 0	8 26 34 0	10 21 40 0	4 16 24 0	15.3	16.2	15.3 12.3	17.4 17.2	13.0 14.1 14.9 12.8	24.6 22.8 20.1 25.2
HIGH MEAN LOW MEAN EXP MEAN C.V. % LSD 5% LSD 1% # OF REPS		50 0 20 67 22 31 4	78 3 36 32 18 26 4	49 0 20 40 13 19 4	34 0 17 82 22 NS 4	40 0 18 59 17 24 4	U	10.8	9.4	12.3	17.1 17.6	14.9 12.8 13.7 21.8 NS NS	20.1

a Wm=Wheaton, Alx=Alex, AlxTr=Alex treated, Cot=Coteau, Rug=Rugby durum, and Rbt=Robust barley;

# Summary

Triallate plus trifluralin at commercial rates (16+5.3 oz/A) did not cause important injury to any of the small grain cultivars. These herbicides at twice the commercial rate caused severe injury to untreated "Alex' wheat, but not to Mon 5500 treated Alex wheat. The soil was extremely dry at seeding which would have increased the amount of herbicide residual, but also reduced "activation" of the herbicides during seed emergence.

Broadleaf weed control in wheat, Williston 1988. 'Len' Hard Red Spring wheat was seeded on April 29. Treatments were applied to 3.5-leaf wheat, 2- to 3-leaf green foxtail, and broadleaf weeds less than 2 inch tall on May 23 and 26 with 55 to 60 F, 30% relative humidity and 5 mph wind at application and a high of 86 F. Treatments were applied with a tractor-mounted plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft wide area the length of 10 by 25 ft plots. The experiment was a randomized complete block with four replications. Evaluation was on June 28 and July 15. Weed infestations were sparce for wild oats and kochia, medium for green foxtail, and dense for Russian thistle. Harvest was from 80 sq ft on July 25.

			June	28	Ju	ıly 15	5		Wheat	
Treatment	Rate	Grft	Ruth	KOCZ	Grft	Ruth	KOCZ	Inj	<u>Yield</u>	Tw
11 Cabinette	(%)			(% cor	ntrol)			(%)(	bu/A)	(1b/bu)
		0	٥٤	20	0	84	25	1	3.1	60.6
2,4-Ddea	6	0	95 58	29	0	24	13	0	2.8	59.8
MCPAda	6	0				91	60	1	4.3	60.5
2,4-Ddea+X-77	6	0	97	63	0		78	4	3.6	61.1
Dicamba+2,4-Ddea	1.5+4	0	92	89	6	86	80	8	3.3	60.6
Dicamba+MCPAdma	1.5+4	0.	89	79	0	80	51	6	3.3	60.7
Picl+2,4-Ddea	0.25+6	0	94	39	13	91		4	2.9	60.6
Bromoxynil&MCPA	6	0	86	45	0	76	66		3.8	60.5
Bromoxyni1&MCPA	8	0	95	85	0	86	88	0	3.4	60.4
DPX-M6316+X-77	0.25	0	98	53	9	95	44	2	3.4	60.8
DPX-L5300+X-77	0.125	0	95	95	0	85	97	4		
DPX-L5300+2,4-Ddea+X-77	0.125+4	0	96	95	8	85	95	3	3.2	60.9
DPX-L5300+Dicamba+X-77	0.125+1+	0	96	99	8	91	97	4	3.1	61.1
DPX-M6316+DPX-L5300+X-77	0.3+0.12		98	92	13	95	84	1	3.6	60.9
Metsulfuron+X-77	0.062	28	99	99	35	99	98	0	4.5	60.8
Metsulfuron+2,4-Ddea+X-77	0.062+4	0	99	98	23	99	93	0	5.0	60.6
Clopyralid&2,4-D	9.5	0	94	26	0	90	20	1	4.2	60.9
Fluroxypyr	1.5	0	55	95	6	41	91	0	3.7	60.6
Fluroxypyr+2,4-Ddea	1.5+6	0	94	97	29	90	96	4	4.0	60.9
Diclofop+Bromoxynil	12+4	45	90	82	68	78	80	0	4.2	60.6
Fenoxaprop&2,4-D&MCPA	6.8	0	66	19	19	48	28	4	3.3	60.4
Untreated Check	0	0	0	0	0	0	0	0	2.3	58.0
HIGH MEAN		45	99	99	68	99	98	8	5.0	61.1
		0	0	0	0	0	0	0	2.3	58.0
LOW MEAN		4	85	66	11	77	66	2	3.6	60.5
EXP MEAN		221	12	26	135	13	31	150	24.2	
C.V. %		20	14	24	21	15	28	5	1.2	
LSD 5%		NS	19	32	28	19	38	NS	NS	
LSD 1%		2	4	4	4	4	4	4	4	1
# OF REPS		_	7							

#### Summary

None of the herbicides cause important injury to wheat. Diclofop + bromoxynil gave the greatest green foxtail control which was only 45% at the first and 67% at the second evaluation. X-77 with 2,4-Ddea enhanced kochia control and tended to enhance Russian thistle control. DPX-M6316 was less effective than the other sulfonylureas in controlling kochia, but equal in controlling Russian thistle. Fluroxypyr gave more than 90% season long kochia control. Wheat yield was low because of drought conditions.

Broadleaf weed control in wheat, Langdom 1988. 'Marshall' Hard Red Spring wheat was seeded on May 6. Treatments were applied to 5-leaf wheat, cotyledon to 4-leaf ladysthumb, and 2.5-leaf wild buckwheat on June 1. Treatments 1 through 15 were applied prior to a 0.09 inch rain between 2:30 and 4:00 PM. Treatments 16 through 20 were applied after the rain at 6:00 PM. All treatments were applied in 8.5 gpa at 35 psi with a bicycle wheel type plot sprayer to an 8 ft wide area the length of 10 by 30 ft plots. The experiment was randomized complete block with four replications. Evaluation was on July 0. The stand of ladysthumb was dense with more than 1 plant/sq ft and wild 0.

<u>Treatment</u>		Whea				Ladys-
·······································	Rate		Yield	Tswt	Wioa	thumb
	(oz/A)	(%)	(bu/A	)(1b/bu)	(%	control)
2,4-Ddea	6					
MCPAdma	6	0	43.0	61.4	0	54
2,4-Ddea+X-77	6+0.25%	0	43.5		0	41
Dicamba+2,4-Ddea	1.5+4	0	40.6	60.1	0	73
Dicamba+MCPAdma	1.5+4	0	40.5	60.5	0	70
Picloram+2,4-Ddea	0.25+6	0	39.4	59.9	0	59
Bromoxynil&MCPA	6	0	40.3	61.1	0	53
Bromoxyni1&MCPA	8	0	39.2 31.5	60.1	5	97
DPX-M6316+X-77	0.25+0.25%	ő	36.2	61.0	0	99
DPX-L5300+X-77	0.125+0.25%	0	33.6	61.3	0	71
DPX-L5300+2,4-Ddea+X-77	0.125+4+0.25%	Ö	37.2	60.8	0	49
DPX-L5300+Dicamba+X-77	0.125+1+0.25%	Ö	36.9	61.4	0	20
DPX-M6316+DPX-L5300+X-77	0.3+0.125+0.25%	0	38.4	61.0	0	45 78
Metsulfuron+X-77	0.062+0.25%	0	35.5	60.4	5	45
Metsulfuron+2,4-Ddea+X-77 Clopyralid&24-D	0.062+4+0.25%	0	33.2	61.0	6	68
Fluroxypyr	9.5	0	39.2	60.1	0	97
Fluroxypyr+2,4-Ddea	1.5	0	38.3	60.6	0	86
Diclofop+Bromoxynil	1.5+6	0	42.2	61.1	0	96
Fenoxyprop&24-D&MCPA	12+4	0	42.6	60.8	82	90
Untreated	6.8	5	42.4	60.8	85	71
	0	0	40.0	60.4	0	0
C.V. %		0.51				
LSD 5%		351	17.4	1.9	39	21
LSD 1%		1	NS	NS	5	19
# OF REPS		2	NS	NS	6	26
		4	4	4	4	4

# Summary

None of the treatments caused important injury to wheat Diclofop + bromoxynil and Fenoxaprop+2,4-D+MCPA gave similar wild oats control. Ladysthumb control was 90% or more with bromoxynil treatments, clopyralid plus 2,4-D, and fluoroxypyr plus 2,4-D. Wheat grain yield and test weight were not influenced by herbicide treatments.

Broadleaf weed control in oats, Minot 1988. 'Dumont' oats was seed on May 7. Treatments were applied to 4-leaf oats; 1- to 2-inch tall common lambsquarter, prostrate pigweed, and green foxtail; and 1- to 3-inch tall Russian thistle, and Kochia on June 1 with 75 F and calm wind. All treatments were applied in 8.5 gpa at 35 psi with a bicycle wheel type plot sprayer to an 8 ft wide area the length of 10 by 24 plots. The experiment was randomized complete block with four replications. Evaluation was on July 18. Density of all weeds exceeded 1 plant/sq. ft.

		Oat						Dat
Treatment	Rate	inj	Grft	KOCZ	Colq	Ruth	YLD	TW
11 Casiments	(oz/A)	(%)		(% co	ntrol	)	(bu/A	)(1b/bu
		0	0	EA	90	84	18.1	25.7
2,4-Ddea	6	8	0	54 30	88	18	12.8	23.8
MCPAdma	6				98	96	19.4	29.5
2,4-Ddea+X-77	6+0.25%	10	0	84	94	84	18.3	26.8
Dicamba+2,4-Ddea	1.5+4	20	3	85	97	64	15.0	25.3
Dicamba+MCPAdma	1.5+4	13	0	93	99	84	17.0	26.2
Pic1+2,4-Ddea	0.25+6	8	0 5	59 70	98	72	16.0	29.9
Bromoxyni1&MCPA	6	5	0	80	99	82	15.3	28.1
Bromoxynil&MCPA	8	9	9	88	78	95	20.1	32.0
DPX-M6316+X-77	0.25+0.25%		14	94	91	81	16.7	30.8
DPX-L5300+X-77	0.125+0.25%	15	13	98	98	91	18.1	29.5
DIX LOCULLY!	0.125+4+0.25%	6	23	96	93	89	24.0	28.1
DPX-L5300+Dicamba+X-77	0.125+1+0.25%	13		96	97	98	20.7	32.2
D. 7	0.3+0.125+0.25%	3	0 5	98	98	99	24.1	31.2
Metsulfuron+X-77	0.062+0.25%	8	10	99	99	99	24.1	32.3
Metsulfuron+2,4-Ddea+X-77	0.002+4+0.23%	15	5	76	98	91	14.1	27.5
Clopyralid&2,4-D	9.5	11	0	96	0	0	13.5	22.7
Fluroxypyr	1.5 1.5+6	13	15	96	99	94	17.0	28.8
Fluroxypyr+2,4-Ddea	12+4	32	49	34	89	83	19.6	26.6
Diclofop+Bromoxynil	6.8	43	51	71	99	74	16.8	28.4
Fenoxaprop&2,4-D&MCPA	0.8	0	0	0	0	0	14.5	23.4
Untreated	19.0	Ö	29	40	98	8	28.0	24.9
Propanil&MCPA	19.0	0	44	65	99	82	39.6	32.0
Propanil&MCPA+Bromoxynil	1372							
C.V. %		93	98	16	5	15	41.4	6.0
LSD 5%		13	16	17	7	15	11.2	2.4
LSD 1%		18	22	22	9	20	14.9	3.2
# OF REPS		4	4	4	4	4	4	4
" OI KLIS						Take!		

#### Summary

All herbicides, except fluroxypyr, gave 90% or more common lambsquarters control. Diclofop and fenoxoprop only gave about 50% green foxtail control probably because of emergence after treatment because of a late season rain. DPX-65300+2,4-D or +DPX-M6316, metsulforan alone or with 2,4-D, and fluroxypyr+2,4-D gave 90% or more control of all broadleaf weeds. DPX-L5300 was more effective than DPX-M6316 in controlling Kochia and common lambsquarter but DPX-M6316 was more effective than DPX-L5300 on Russian thistle. Oat yields and test weight generally related to the degree of weed control.

Broadleaf weed control in wheat, Carrington 1988. 'Stoa' Hard Red Spring wheat was seeded on May 9. Treatments were applied to 5-leaf wheat, 2- to 3-leaf green foxtail, and 3 inch tall common lambsquarters on June 9. Plants were drought stressed and high temperatures were 84 to 96 F for 1 week after treatment. Treatments were applied in 8.3 gpa at 3.5 psi with a bicycle type plot sprayer delivery to 8 by 24 ft plots. The experiment was a randomized complete block with four replications. Evaluation was on July 15. Weed densities were less than 1 plant/2 sq yds.

Treatment	Doto	Wheat	da area	a library
	Rate	injury	Grft	Colq
	(oz/A)	(%)	(% co	ntrol)
2,4-Ddea				Tributh alarm
MCPAdma	6	0	0	82
2,4-Ddea+X-77		0	0	78
Dicamba+2,4-Ddea	6+0.25%	0	0	94
Dicamba+MCPAdma	1.5+4	0	0	86
	1.5+4	0	0	82
Picloram+2,4-Ddea	0.25+6	0	0	99
Bromoxynil&MCPA	6	0	0	96
Bromoxynil&MCPA DPX-M6316+X-77	8	0	0	92
DPX-L5300+X-77	0.25+0.25%	0	0	59
DDY 15300+X-//	0.125+0.25%	1	0	94
DPX-L5300+2,4-Ddea+X-77	0.125+4+0.25%	0	0	91
DPX-L5300+Dicamba+X-77	0.125+1+0.25%	0	0	99
DPX-M6316+DPX-L5300+X-77	0.3+0.125+0.25%	0	0	95
Metsulfuron+X-77	0.062+0.25%	1	1	67
Metsulfuron+2,4-Ddea+X-77	0.062+4+0.25%	i	0	83
Clopyralid&2,4-D	9.5	Ō	Ŏ	99
Fluroxypyr	1.5	ő	Ö	0
Fluroxypyr+2,4-Ddea	1.5+6	ő	0	99
Diclofop+Bromoxynil	12+4	ő	33	
Fenoxyprop&2,4-D&MCPA	6.8	i	86	59
Untreated Check	0	0	0	99
		· ·	0	0
C.V. %		447	43	10
LSD 5%		NS		19
LSD 1%			3	21
# OF REPS		NS	5	28
		4	4	4

## Summary

None of the herbicide treatment caused any visable injury to wheat. Green foxtail control with fenoxaprop+2,4-D+MCPA (Tiller) was more than 80% while with diclofop+bromoxynil only 33%. Common lambsquarter control exceed 80% with most herbicide treatments, except MCPA, DPX-M6316, metsulfuron, fluroxypyr, and bromoxynil applied with diclofop.

Broadleaf weed control in wheat, Casselton 1988. 'Stoa' wheat was seeded on April 14. Treatments were applied to 4-leaf wheat and foxtail; and 1-to 3-inch tall common lambsquarters and wild mustard on June 1, with 88 F, 50% RH, and dry soil at application. All treatments were applied in 8.5 gpa at 35 psi with a bicycle wheel type plot sprayer to an 8 ft wide area the length of 10 by 30 ft plots. The experiment was randomized complete block with four replications. Evaluation was on June 15. Weed densities were sparce and variable. Many weeds had apparently died from the drought as populations were high at treatment. Wheat grain yeild was not taken because the drought stress masked any differences from treatment and caused variation in growth whithin replications.

	Data	Wheat injury	KOCZ	Wimu	Colq
Treatment	Rate	(%)	1%	contr	
	(oz/A)	(10)	10	Contr	01)
	•	3	70	98	99
2,4-Ddea	6	3	50	99	99
MCPAdma	6	0	78	99	99
2,4-Ddea+X-77	6+0.25%	6	93	99	98
Dicamba+2,4-Ddea	1.5+4	8	90	99	99
Dicamba+MCPAdma	1.5+4		60	99	99
Picloram+2,4-Ddea	0.25+6	0		99	99
Bromoxynil&MCPA	6	1	60	99	99
Bromoxynil&MCPA	8	. 0	80		65
DPX-M6316+X-77	0.25+0.25%	0	80	81	75
DPX-L5300+X-77	0.125+0.25%	1	60	82	
DPX-L5300+2,4-Ddea+X-77	0.125+4+0.25%	0	95	99	97
DPX-L5300+Dicamba+X-77	0.125+1+0.25%	1	80	89	86
DPX-M6316+DPX-L5300+X-77	0.3+0.125+0.25%	1	95	93	96
Metsulfuron+X-77	0.062+0.25%	5 3	85	95	82
Metsulfuron+2,4-Ddea+X-77	0.062+4+0.25%	3	0	99	95
Clopyralid&2,4-D	9.5	0	0	99	99
Fluroxypyr	1.5	0	70	50	15
Fluroxypyr+2,4-Ddea	1.5+6	1	95	99	97
Diclofop+Bromoxynil	12+4	1	0	99	98
Fenoxaprop&2,4-D&MCPA	6.8	0	0	99	99
Untreated	0	0	0	0	0
C.V. %		141		13	11
LSD 5%		3		17	13
LSD 1%		4		22	18
# OF REPS		4	1	4	4

## Summary

All herbicides gave 85% or more common lambsquaters control, except DPX-M6316, DPX-L5300, metsulfuron and fluroxypyr. None of the herbicide treatments caused visable injury to wheat. Kochia was only present in one replication, but control only exceeded 90% with dicamba treatments, DPX-L5300+2,4-D, or DPX-L5300 + DPX-M6313, and fluroxypyr+2,4-D.

Broadleaf weed control in wheat with sulf@nylureas, Casselton 1988. 'Stoa' Wheat was seeded on April 14. Treatments were applied to 4-leaf wheat and foxtail; and 1-to 3-inch tall common lambsquarter and wild mustard on June 1, with 88 F, 50% RH, and dry soil at application. All treatments were applied in 8.5 gpa at 35 psi with a bicycle wheel type plot sprayer to an 8 ft wide area the length of 10 by 30 ft plots. The experiment was randomized complete block with four replications. Evaluation on August 1. Weed densities were sparce and variable. Many weeds had apparently died from the drought as populations were high at treatment. Wheat grain yield was not taken because the drought stress masked any differences from treatment and cause variation in growth whithin replications.

		Wheat		
Treatment	Rate	injury	KOCZ	Colq
	(oz/A)	(%)	-(% cor	
DPX-R9674+X-77	0 125.0 25%		0.5	0.4
DPX-R9674+X-77	0.125+0.25% 0.225+0.25%	1	95	94
DPX-R9674+X-77	0.30+0.25%	0	97	98
DPX-R9674+X-77	0.30+0.25%	0	95	98
DPX-R9674+X-77	0.45+0.25%	1	99	97
DPX-R9674+2,4-Ddea+X-77	0.45+0.25%	0	97	98
DPX-R9674+Dicamba+X-77	0.30+4+0.25%	3	99	99
DPX-L5300+X-77	0.125+0.25%	0 1	99	96
DPX-L5300+X-77	0.125+0.25%		95	97
DPX-L5300+X-77	0.25+0.25%	0	97	97
DPX-L5300+2,4-Ddea+X-77	0.125+4+0.25%		81	99
DPX-L5300+Dicamba+X-77	0.125+1+0.25%	0	99	99
Bromoxynil			99	98
Bromoxyni1&MCPA	4	0	83	99
Bromoxyni1&MCPA	8	0	84	95
Bromoxynil+DPX-R9674+X-77	2+0.12+0.25%	0 3	86	99
Bromoxyni1+DPX-R9674+X-77	2+0.12+0.25%	0	94 82	97
Bromoxynil+DPX-R9674+X-77	3+0.12+0.25%	0	95	94
Bromoxynil+DPX-R9674+X-77	3+0.12+0.25%	3	95	98
AC 222,293	5	0	0	96 8
AC 222,293+Bromoxynil&MCPA	5+4	1	95	99
2,4-Dbee	6	4	99	99
Dicamba+MCPA	1.5+6	3	98	99
	1.010		90	33
C.V. %		272	12	4
LSD 5%		NS	15	
LSD 1%		NS	19	5 7
# OF REPS		4	4	4
		THE BEST OF STREET		110000000000000000000000000000000000000

#### Summary

None of the herbicide treatments caused visable injury to wheat. Control of both kochia and common lambsquarter was high with nearly all treatments. The extreme drought caused death to weed even in untreated areas. The extreme stress for moisture plus the herbicide stress may have accented the high level of weed control.

Fluroxypyr for weed control in wheat, Fargo 1988. 'Wheaton' Hard Red Spring wheat was seeded to dry soil on April 14. Treatments were applied to 4-to 5-leaf wheat, 4- to 6-inch tall common lambsquarter and 2 inch tall wild mustard and redroot pigweed on June 2 with 80 F, 60% RH, no wind at application. All treatments were applied in 8.5 gpa at 35 psi with a bicycle wheel type plot sprayer to an 8 ft wide area the length of 10 by 30 ft plots. The experiment was randomized complete block with four replications. Evaluation was on July 21. All weeds exceeded 5 plants/sq. ft.

with the same of t		Wheat				
Treatment	Rate		Vosf	Rrpw	Colq	KOCZ
11 eachenc	(oz/A)	(%)			ntrol)	
	(02/11)	(10)		(70 00	,	
Fluroxypyr	1	0	85	25	0	98
Fluroxypyr	1.5	0	92	41	0	99
Fluroxypyr	2	0	94	34	8	99
Fluroxypyr+2,4-Ddea	1+6	0	99	95	99	99
Fluroxypyr+2,4-Ddea	1.5+6	0	99	99	99	99
2,4-Ddea	6	0	98	96	96	73
Fluroxypyr+MCPA-dma	1+6	0	98	87	99	95
Fluroxypyr+MCPA-dma	1.5+6	3	98	92	97	96
MCPA-dma	6	0	99	87	99	72
Fluroxypyr+Picloram+2,4-Dd		1	99	99	99	99
Fluroxypyr+Picloram+2,4-Dd		0	99	99	99	99
Picloram+2,4-Ddea	0.25+6	0	99	99	99	90
Fluroxypyr+Clopyralid&2,4-		0	99	96	99	99
Fluroxypyr+Clopyralid&2,4-		1	99	98	99	99
Clopyralid&2,4-D	9.5	0	99	98	99	85
Dicamba+MCPA-dma	1+6	0	98	99	99	96
Bromoxynil&MCPA	8	0	99	96	99	87
C V %		402	2	1.4	r	10
C.V. %		493	2	14	5	10
LSD 5%		NS	3	16	6	13
LSD 1% # OF REPS		NS 4	4	21	8	17 4
# UF KEYS		4	4	4	4	4

# Summary

Fluroxypyr at 1 oz/A gave 85% control of volunteer sunflower and 98% control of kochia. Fluroxypyr did not give adequate control of redroot pigweed or common lambsquarter at any rate evaluated. Fluroxypyr in combination with MCPA or 2,4-D gave greater then 95% control of all weeds. However, any of these herbicides applied alone did not give 95% control of all weeds. Dicamba+MCPA also gave 95% or more control of all weeds. Wheat was not harvested for yield because of inadequate growth and stand variabilty caused by an extreme drought.

Preemergence foxtail control in wheat, Fargo, 1988. 'Wheaton' Hard Red Spring was seeded on May 16, and treatment were applied to a moist surface soil which was dry below on May 19 with 60 F, 65 RH, and cloudy sky. Treatments were applied in 17 gpa at 35 psi with a bicycle wheel type plot sprayer to an 8 ft wide strip the length of 10 by 24 ft. plots. A rain occurred within 1 h after treatments were applied. Evaluations for wheat injury and weed control were on June 20, July 26, and August 18. All weeds exceeded 1 plant/sq. ft., except redroot pigweed. The experiment was a randomized complete block with 4 replications. Wheat was not harvested for grain yield because of extreme drought and poor wheat growth.

		Ju	ne 20-		,	July 26	ĵ	Aug 18
Rate	Wht	Grft	KOCZ	Rrpw	Wht i	COCZ (	Grft.	KOCZ
(oz/A)								
0.24	0	38	64	0	0	26	11	0.4
								24
_								49
								74
								51
								41
								72
								48
					· ·	U	U	20
	0	95	94	85	0	89	65	74
								20
	0							47
	0							53
	NS	23						NS
	NS	32						NS
	4	4	4	2				4
	Rate (oz/A) 0.24 0.50 1 0.24 0.50 1 48	(oz/A) 0.24	Rate Wht Grft (oz/A)  0.24	(oz/A)	Rate Wht Grft KOCZ Rrpw (oz/A)(%  0.24	Rate Wht Grft KOCZ Rrpw Wht I (oz/A)	Rate Wht Grft KOCZ Rrpw Wht KOCZ (OZ/A)	Rate   Wht Grft KOCZ Rrpw   Wht KOCZ Grft

# SUMMARY

A-1237 and C-4243 give good early foxtail and kochia control. However, late season weed control was not adequate probably because of lack of competition from the wheat with the drought.

Postemergence foxtail control in wheat, Fargo, 1988. 'Wheaton' Hard Red Spring wheat was seeded to dry soil on May 16. The treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi. The 2-leaf treatments were applied to 3-leaf wheat and green foxtail and 0.75 to 4 inch tall broadleaf weeds on June 6 with 90F, 35% RH, clear sky, 20 mph wind and dry soil conditions. The 5-leaf treatments were applied to 5-leaf foxtail and 4 inch kochia on June 15 with 72F. The experiment was a randomized complete block with four replications. Treatments were applied to an 8 ft wide strip the length of 10 by 24 ft plots. Evaluation was on June 29 and July 26. Weeds exceeded 5 plants/sq. ft. Plants were severely drought stressed throughout the season.

					29			26
Treatment			Wht	Grft	KOCZ	Wht	KOCZ	<u>Fxtl</u>
		(oz/A)			(% co	ntrol	)	
HOE-7113-04H	(21f)	1.28	3	75	35	0	9	68
HOE-7113-04H	(51f)	1.28	3	48	15	1	15	82
Feno&24-D&MCPA	(51f)	5.3	3	44	68	0	63	58
Feno&24-D&MCPA	(21f)	5.3	3 3 3 3	54	56	0	51	43
Feno&24-D&MCPA	(51f)	6.2	3	0	33	5	38	31
Feno&24-D&MCPA	(21f)	6.2	0	13	63	3	39	30
A-1237	(21f)	0.24	26	55	95	36	83	20
A-1237	(21f)	0.50	18	30	82	35	80	10
B-4243	(21f)	0.24	5	9	26	5	23	19
C-4243	(21f)	0.50	0	5	5	1	21	18
Propanil&MCPA	(21f)	19	0	33	46	Ō	17	18
Diclofop+Brox	(21f)	12+4	0	14	5	Ö	20	28
BAS-51400H+BAS-090	(21f)	4+0.25G	11	84	76	14	59	88
BAS-51400H+MSC	(21f)	4+0.25G	3	83	64	1	59	69
BAS514+DPX-M6316+MSC	(21f)	4+0.25+0.25G	4	56	85	3	87	75
		4+0.25+0.25G	15	78	96	0	92	79
BAS514+DPX-M6316+BAS090	(21f)	0	0	0	0	0	0	0
Untreated		U	U	U	U	U	U	U
LITCH MEAN			26	84	96	36	92	88
HIGH MEAN			26	0	0	0	0	0
LOW MEAN			0			6		
EXP MEAN			6	40	50		44	43
C.V. %			121	46	32	124	33	34
LSD 5%			9	26	23	11	21	21
LSD 1%			13	35	30	14	28	28
# OF REPS			4	4	4	4	4	4

#### Summary

BAS-514 applied with MSC or BAS-090 and HOE -7113 controlled green foxtail without causing important injury to wheat. DPX-M6316 tended to reduce green foxtail and increase kochia control when applied with BAS-514. A-1237 caused severe contact injury to wheat shortly after treatment and injury persisted throughout the season.

2,4-D diethanolamine with carbonates Exp 1, Fargo 1988. 'Wheaton' Hard Red Spring wheat was seeded April 14, to dry soil. Treatments were applied to 4-to-5 leaf wheat, 2-to-4 leaf redroot pigweed, and 1-to-6 inch common lambsquarter on June 3 with 88 F, 60% RH, and 10 mph wind. Treatments were on 8.5 gpa at 35 psi applied with a bicycle wheel type plot sprayer to an 8 ft. wide strip the length of 30 by 10 ft. plots. The experiment was a randomized complete block with four replications. Evaluations were on June 21 and July 25. Weed infestations exceeded 5 pr sq. yd., except for kochia which was variable.

				\ <del>.</del>				
Twostmont			lune 2		J	uly 2	25	Wheat
Treatment	Rate	Wht	Rrpw		Wht	KOCZ	Rrpw	yield
	(oz/A)			· (% co	ntrol	)		(Bu/A)
0.4.0.1	13 13 19							
2,4-Ddea	6	1	88	97	0	83	98	20.7
2,4-Ddea+NaHCO3	6+16	0	53	45	0	13	52	19.5
2,4-Ddea+Na2CO3	6+16	0	30	58	0	38	66	19.4
2,4-Ddea+NaHCO3+28%N	6+16+1G	0	93	96	0	73	99	19.9
2,4-Ddea+Na2CO3+28%N	6+16+1G	0	94	99	3	65	98	19.7
2,4-Ddea+NaHCO3+SCI-40	6+16+0.25%	0	54	55	0	35	75	18.5
2,4-Ddea+Na2CO3+SCI-40	6+16+0.25%	0	43	41	0	31	63	17.5
2,4-Ddea+NaHCO3+LI-700	6+16+0.25%	Ö	61	81	0	45	83	
2,4-Ddea+Na2CO3+LI-700	6+16+0.25%	0	73	91	1			20.0
2,4-Ddea+NaHCO3+MS	6+16+0.25G	0	95	97		33	87	17.4
2,4-Ddea+Na2CO3+MS	6+16+0.25G	0	97		0	89	97	20.7
2,4-Ddea+NaHCO3+AG-2	6+16+0.25G	1		100	0	92	98	19.8
2,4-Ddea+Na2CO3+AG-2	6+16+0.25G		99	99	1	85	99	22.8
2,4-Ddea+NaHCO3+PO		0	99	99	1	70	99	19.9
2,4-Ddea+Na2CO3+PO	6+16+0.25G	1	97	99	1	54	99	19.2
2,4 Ddo2+NaUCO2+V 77	6+16+0.25G	0	97	99	0	67	99	18.9
2,4-Ddea+NaHCO3+X-77	6+16+0.25%	0	72	98	0	46	95	18.9
2,4-Ddea+Na2CO3+X-77	6+16+0.25%	0	75	97	1	51	94	18.4
2,4-Ddea+NaHC03+MS+28%N	6+16+0.25G+16		99	99	1	86	99	18.7
2,4-Ddea+Na2CO3+MS+28%N	6+16+0.25G+16	4	99	99	0	92	99	19.3
Untreated	0	0	0	0	0	0	0	17.4
HIGH MEAN		4	99	100	3	92	99	22.8
LOW MEAN		0	0	0	0	0	0	17.4
EXP MEAN		0	77	84	0	59	86	19.2
C.V. %		334	11	11	336	37	15	12.0
LSD 5%		NS	12	13	NS	31	18	NS
LSD 1%		NS	15	17	NS	41	24	NS
# OF REPS		4	4	4	4	4	4	4
			7		_	4	4	4

#### Summary

Redroot pigweed, common lambsquarter, and kochia control with 2,4-Ddea was antagonized by sodium bicorbonate or carbonate in the spray solution. Ammonium nitrate, methylated seed oil (MS), petroleum oil, AG-2, and Inhance all overcame the antagonism caused by the carbonates in the solution.

2,4-D diethanolamine with carbonates Exp 2, Fargo 1988. 'Wheaton' Hard Red Spring wheat was seeded on May 16 to dry soil. Treatments were applied to 4- to 5-leaf wheat, 4- to 6-inch tall redroot pigweed, and kochia on June 13 Treatments were in 8.5 gpa at 35 psi applied with a bicycle wheel type plot sprayer to a 8 ft. wide strip the length of 24 by 10 ft. plots The experiment was a randomized complete block with four replications. Evaluations were on June 30 and August 18. Injury to wheat was not evaluated because of drought stress. Kochia and redroot pigweed infestations generally exceeded 5 plant per square yard, but were variable.

		3	20	Λ	+ 10
Tuestment	Data	<u>June</u> KOCZ	Rrpw	Augus KOCZ	Rrpw
Treatment	Rate (oz/A)	KUCZ		ntrol)	KIPW
	(UZ/A)		- ( // CUI	10101)	
2,4-Ddea	6	64	75	35	63
2,4-Ddea+NaHCO3	6+16	26	25	31	33
2,4-Ddea+Na2CO3	6+16	48	54	08	34
2,4-Ddea+NaHCO3+28%N	6+16+1G	45	54	11	76
2,4-Ddea+Na2CO3+28%N	6+16+1G	55	62	51	69
2,4-Ddea+NaHCO3+SCI-40	6+16+0.25%	13	15	38	44
2,4-Ddea+Na2CO3+SCI-40	6+16+0.25%	15	48	05	56
2,4-Ddea+NaHCO3+LI-700	6+16+0.25%	29	26	18	53
2,4-Ddea+Na2CO3+LI-700	6+16+0.25%	33	51	24	45
2,4-Ddea+NaHCO3+MS	6+16+0.25G	83	73	63	45
2,4-Ddea+Na2CO3+MS	6+16+0.25G	82	76	75	51
2,4-Ddea+NaHCO3+AG-2	6+16+0.25G	72	79	61	65
2,4-Ddea+Na2CO3+AG-2	6+16+0.25G	73	78	60	73
2,4-Ddea+NaHCO3+PO	6+16+0.25G	65	75	60	75
2,4-Ddea+Na2CO3+PO	6+16+0.25G	64	69	51	75
2,4-Ddea+NaHCO3+X-77	6+16+0.25%	44	48	28	28
2,4-Ddea+Na2CO3+X-77	6+16+0.25%	54	61	40	34
2,4-Ddea+NaHCO3+MS+28%N	6+16+0.25G+1G	97	94	90	93
2,4-Ddea+Na2CO3+MS+28%N	6+16+0.25G+1G	95	97	86	90
Untreated	0	0	0	0	0
HIGH MEAN		97	97	90	93
LOW MEAN		0	0	0	0
EXP MEAN		55	60	43	56
C.V. %		30	24	38	28
LSD 5%		23	20	23	22
LSD 5%		31	27	31	29
# OF REPS		4	4	4	4
π OI KEF3					

## Summary

Both kochia and redroot pigweed control with 2,4-Ddea was antagonized by sodium carbonate or sodium bicarbonate. The antongonism generally tended to be greater from bicarbonate than carbonate. The oil adjuvants and Inhance overcame to antagonism of 2,4-D caused by both bicarbonate and carbonate. Methylated seed oil (MS) with 28%N liquid fertilizer tended to enhance weed control with 2,4-Ddea more than any other adjuvants.

Kochia control with 2,4-D, Fargo 1988. 'Wheaton' Hard Red Spring wheat was seeded on May 16 to dry soil. Treatments were applied to 4- to 5-leaf wheat, 4- to 6-inch tall redroot pigweed, and kochia on June 13 with 83 F, 60% RH, and 10 mph wind. Treatments were in 8.5 gpa at 35 psi applied with a bicycle wheel type plot sprayer to a 8 ft. wide strip the length of 24 by 10 ft. plots. The experiment was a randomized complete block with four replications. Evaluations were on June 29 and August 18. Injury to wheat was not evaluated because of drought stress. Kochia and redroot pigweed infestations generally exceeded 5 plant per square yard, but were variable.

<u>Treatment</u>	Rate	Dynu	June	29	Auc	just 18
	(oz/A)	(%	KOCZ	rol)	<u>KOCZ</u> (% c	Rrpw control)
2,4-Ddea 2,4-Ddea+X-77 2,4-Ddea+X-77 2,4-Ddea+PO 2,4-Ddea+PO 2,4-Ddea+MSC 2,4-Ddea+MSC 2,4-Ddea+AG-2 2,4-Ddea+SCI-40 2,4-Ddea+SCI-40 2,4-Ddea+LI-700 2,4-Ddea+LI-700 2,4-Ddea+INHANCE 2,4-Ddea+INHANCE 2,4-Ddea+28%N 2,4-Ddea+28%N Brox&MCPA Untreated	6 6 6 6+0.25% 6+0.25G 6+0.25G 6+0.25G 6+0.25G 6+0.25% 6+0.25% 6+0.25% 6+0.25% 6+0.25G 6+0.25G 6+1G 6+1G 8 0	60 93 76 90 68 94 89 94 81 92 46 88 54 90 66 87 58 83 68	50 93 65 90 72 95 89 94 43 92 58 94 79 93 54 82 83 0	99 99 99 99 99 99 99 99 99 99 99 99	64 84 71 80 61 84 79 83 80 30 83 57 87 69 54 39 79 80 0	60 88 81 80 70 85 92 94 92 89 39 86 76 92 74 63 56 89 70 0
HIGH MEAN LOW MEAN EXP MEAN C.V. % LSD 5% LSD 1% # OF REPS		94 0 74 18 19 25 4	95 0 75 17 18 24	99 0 94 1 1 2 4	87 0 67 22 21 28	94 0 74 22 23 31
			4	4	4	4

#### Summary

2,4-D butoxyethanol ester (bee) gave greater kochia and redroot pigweed control than the 2,4-D diethanolamine (dea). Adjuvants with 2,4-Ddea did not enhance weed control, except 28% nitrogen fertilzer tended to reduce control with the 2,4-Dbee at the early evaluation. 2,4-Ddea applied with methylated seed oil (MSC) or AG-2 gave weed control equal to that with 2,4-Dbee. Wild mustard was controlled by all treatments.

Carbonate antogonism of 2,4-D, Exp 1. Fargo 1988. 'Wheaton' Hard Red Spring wheat was seeded April 14 to dry soil. Treatments were applied to 4-to 5-leaf wheat, 2-to 4-leaf redroot pigweed, and to 6 inch tall common lambsquarter on June 3. Treatments were in 8.5 gpa at 35 psi applied with a bicycle wheel type plot sprayer to an 8 ft. wide strip the length of 30 by 10 ft plots. The experiment was a randomized complete block with four replications. Evaluations were on June 22 and July 25. Weed infestations exceeded 5 pr sq. yd., except for kochia which was variable.

Treatment	Rate	Wht	Jur Rrpw	e 22 Colq	KOCZ	Wht	July 2 Rrpw	5 Colq	Wheat yield
	(oz/A)	<u>inj</u> (%)	- (%	contro	1)-	(%)	(% co	ntrol)	(Bu/A)
2,4-Ddea 2,4-Dbee 2,4-Dbee+NaHCO3 2,4-Dbee+Na2CO3 2,4-Dbee+Na2CO3 2,4-Dbee+NaHCO3 2,4-Dbee+NaHCO3 2,4-Ddea+Na2CO3 2,4-Dbee+Na2CO3 2,4-Dbee+Na2CO3 2,4-Dbee+X-77 2,4-Dbee+X-77 Untreated	6 6 6+1 6+1 6+1 6+1 8 6+16 8 6+16	0 0 0 0 0 0 0 0 0 2 0 2	93 95 85 97 82 97 35 96 33 98 83 97 0	99 99 99 99 99 99 97 91 98 99	87 94 81 98 83 98 27 97 18 96 90 98	0 0 0 0 0 1 0 1 1 0 0	78 91 74 93 76 92 3 86 3 91 89 86	99 99 79 99 96 99 73 99 25 99 98	17.3 16.4 16.6 12.2 16.5 15.7 15.8 13.9 14.0 16.0 15.3 14.8 15.0
HIGH MEAN LOW MEAN EXP MEAN C.V. % LSD 5% LSD 1% # OF REPS		2 0 0 451 NS NS 3	98 0 76 8 10 14 3	99 0 91 3 5 7	98 0 74 11 14 19 3	1 0 0 321 NS NS 4	93 0 66 13 12 16 4	99 0 82 13 16 21 4	17.3 12.2 15.3 13.3 NS NS

#### Summary

Carbonates at 1 oz/A in a 2,4-D diethanolamine spray mixture, tended to reduce weed control, but carbonates at 16 oz/A significantly reduced weed control. Weed control with 2,4-D butoxyethyl ester was not influenced by carbonates. Surfactant X-77 did not enhance weed control with either 2,4-Ddea or 2,4-Dbee.

Carbonate antagonism of 2,4-D, Exp 2. Fargo 1988. 'Wheaton' Hard Red Spring wheat was seeded to dry soil on May 16. Treatments were applied to 4- to 5-leaf wheat, 4-leaf redroot pigweed, and 4- to 6-inch tall kochia on June 10 applied with a bicycle wheel type plot sprayer to 8 ft wide strip the length three replications. Evaluations were on July 1, 27, and August 18. Kochia plants exceeded 1 plant pr sq. ft and redroot pigweed was variable.

Treatment Rat	ce Whi	j	Wht	July 2 KOCZ	27 Rrpw	Wht	ust 18- KOCZ
2,4-Ddea+NaHCO3 6+ 2,4-Ddea+NaHCO3 6+ 2,4-Ddea+NaHCO3 6+	6 0 6 5 +1 0 +1 3 +1 0 +1 0 16 0 16 4 16 0 16 0	73 89 30 86 38 87 5 85 5 71 82 87 0 89 0 57 17 14 18	0 0 0 0 1 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0	51 80 18 74 26 75 6 69 0 65 58 75 0 80 0 46 30 19 26 4	ntrol 69 88 38 87 34 70 25 86 5 89 72 91 0 91 0 58 23 19 25 4	inj ) (%) (% 0 0 0 6 2 4 0 1 0 1 0 1 0 1 2 2 4 NS 4	29 71 18 79 23 80 0 72 5 50 73 64 0 80 0 43 29 18 24 4

# Summary

Weed control with 2,4-D diethanolamine (dea) was antagonized by both sodium bicarbonate and sodium carbonate in the spray mixture at both 1 and 16 oz/A. Weed control with 2,4-D butoxyethyl ester (bee) was not influenced by Carbonates in the spray. Weed control was greater with 2,4-Dbee them 2,4-Ddea. Surfactant X-77 with 2,4-Ddea tended to increase weed control, but not when with 2.4-Dbee.

Sodium bicarbonate antagonism of 2,4-D, Fargo 1988. 'Wheaton' Hard Red Spring wheat, 'ND810104' oats, and Siberian foxtail millet were seeded as bioassay species in adjacent 6 ft wide strips at Fargo, ND on May 31. Treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft wide area across the bioassay species. Plots were 10 ft wide to provide a 4 ft untreated area between treated areas to aid evaluation. Treatments were applied to late tiller grass species and 6 aid evaluation tredroot pigweed on July 1 with 71 F, 63% RH and no wind.

			7,,7,	y 13			Jul	y 21	
[reatment	Rate (oz/A)	Fomi	Wht (	Oats F	Rrpw % con		Wht	Oats	Rrpw
2,4-Ddea 2,4-Ddea+WK 2,4-Ddea+PO 2,4-Ddea+AG2 2,4-Ddea+NaHCO3 2,4-Ddea+NaHCO3+WK 2,4-Ddea+NaHCO3+AG2 2,4-Ddea+WK 2,4-Ddea+WK 2,4-Ddea+WK 2,4-Ddea+WK 2,4-Ddea+WK 2,4-Ddea+NaHCO3 2,4-Ddea+NaHCO3 2,4-Ddea+NaHCO3+WK 2,4-Ddea+NaHCO3+WK 2,4-Ddea+NaHCO3+AG2 Untreated  HIGH MEAN LOW MEAN EXP MEAN C.V. % LSD 5% LSD 1%	4 4+0.25% 4+0.25G 4+0.25G 4+8 4+8+0.25G 4+8+0.25G 16-0.25G 16+0.25G 16+0.25G 16+8 16+8+0.25G 16+8+0.25G 16+8+0.25G	10 10 15 20 0 5 14 9 29 35 65 70 16 30 0 70 0 21 50 15	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	3 4 6 13 0 3 4 3 11 15 21 23 5 3 6 5 0 7 54 6 7 4	45 55 65 66 8 35 56 53 64 71 88 91 21 39 64 71 0 91 26 4	15 13 13 38 1 11 19 33 30 55 73 80 14 9 30 35 0 80 0 27 48 19 25 4	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		0 51 26 19 25

# Summary

In general redroot pigweed control was enhanced more by Ag-2 than PO (petroleum oil with 17% Atplus 300F) or surfactant X-77 as Adjuvants with 2,4-D applied alone or in the presence of sodium bicarbonate. None of the adjuvants overcame the antagonism caused by sodium bicarbonate of 2,4-D for redroot pigweed control. Injury from 2,4-D to wheat ,oats, and millet related to the level of redroot pigweed control.

BAS 514 for weed control in wheat Exp 1. Fargo 1988. 'Wheaton' Hard Red Spring wheat was seeded to dry soil on April 14. Treatments were applied to 5- to 6-leaf wheat, 2- to 3-inch tall kochia and common lambsquarter, and 6- to 8-leaf volunteer sunflower on June 6 with 85F, 35% RH, cloudless sky, and a 12 mph wind. Treatments were applied in 8.5 gpa at 35 psi with bicycle wheel type plot sprayer to 8 ft wide area the length of 10 by 30 ft plots. Plants drought stressed at treatment. The experiment was a randomized complete block with four replications. Evaluations were on June 22 and July 25.

Treatment	l./h.t		June 2	22			-July	26
	Wht inj	Rrpw	Colq	KOCZ	Vost	Wht	KOCZ	Rrpw
(oz/A)	(%)		- (% co	ntrol)		inj		
BAS514 2 BAS514+BAS-090 2+0.25G	0	13	45	40	30	(,		ntrol
DACE14 DE	0	50	53	53	50	•	41 38	60
BAS514+MS 2+0 256	0	50	28	60	35		60	69 85
BAS514+SO 2+0 250	0	58	15	43	38	Ö	46	83
3AS514 4	Ö	15 45	0	35	33	1	34	38
BAS514+BAS-090 4+0.25G	ő	50	15 0	35	15	1	31	45
BAS514+PO 4+0.25G BAS514+MS 4+0.25G		63	43	63 68	38 50	1	49	57
ACETA OF	5 3 3	73	0	48	43	1 1	34	69
AS514+S0 4+0.25G AS514 8		38	55	65	55	0	38 61	80
AS514+BAS-090 8+0 256	0 5	69	48	33	38	Ö	38	64 53
AS514+PO 8+0 25C	3	73 20	48	63	63	2	68	68
AS514+MS 8+0.25G	Ö	35	15 23	55	50	1	42	64
AS514+SO 8+0.25G ntreated 0	5	45	10	45 40	55 48	3	40	52
orregred 0	0	0	0	15	10	1 0	48	60
IGH MEAN					10	U	0	0
DW MEAN	5	73	55	68	63	3	68	85
(P MEAN	0	0 43	0	15	10	0	0	0
V. %	205	49	25 90	47	40	1	42	59
5D 5% D 1%	NS	NS	NS	37 NS	38	150	22	23
D 1% OF REPS	NS	NS	NS	NS	NS NS	1	13	20
OI KLF3	2	2	2	2	2	2	17	26
					-	7	4	4

Summary
Weed species were variable making evaluation difficult. However, none of the treatment indicated any distinct advantages to the level of control obtained.

BAS-514 for weed control in wheat Exp. 2, Fargo 1988. 'Wheaton' Hard Red Spring wheat was seeded to dry soil on May 16. Treatments were applied to 3-leaf wheat and green foxtail, 0.75- to 4-inch tall kochia, and 6-leaf wild mustard on June 6 with 90 F, 35% RH, sunny sky, 20 mph wind and dry soil. Treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft wide area the length of 10 by 24 ft plots. Evaluations were on June 30 and August 5. Kochia and green foxtail exceeded 1 plant/sq. ft. The experiment was a randomized complete block with three replications. Adjuvants were all at 1 qt/A (0.25G). PO=petroleum oil 11N plus 17% Atplus 300 F, MS=Sun-it, and SO=soybean oil with 15% T-Mulz VO.

olus 17% Atplus 30		J	une 30		A Wht	ugust	5
Treatment  BAS514 BAS514+BAS-090 BAS514+P0 BAS514+S0 BAS514+BAS-090 BAS514+P0 BAS514+P0 BAS514+P0 BAS514+P0 BAS514+P0 BAS514+P0 BAS514+BAS-090 BAS514+BAS-090 BAS514+BAS-090 BAS514+P0 BAS514+P0 BAS514+P0	Rate (oz/A)  2 2+0.25G 2+0.25G 2+0.25G 2+0.25G 4+0.25G 4+0.25G 4+0.25G 8+0.25G 8+0.25G 8+0.25G	Wht	Grft	Nocz ntrol) 15 83 38 53 51 25 81 28 71 73 41 86 71 68 58	0 0 3 0 0 0 0 11 1 1 0 1 1 3	KOCZ (% C 4 63 20 28 15 14 67 13 38 35 23 65 41 40 39	Grft ontrol)  36 94 90 88 88 74 96 94 96 78 96 94 96 97
BAS514+S0 C.V. % LSD 5% LSD 1% # OF REPS	8+0.25G	140 6 9 4	16 17 23 4	22 17 22 4	167 3 4 4	30 13 18 4	9 10 14 4

# Summary

BAS-514 AT 2 to 8 oz/A gave greater then 80% control of green foxtail without important injury to wheat. The BAS-090 adjuvant tended to enhance green foxtail control with BAS-514 more then the other adjuvants at the June 30 evaluation. All adjuvants simularly enhanced BAS-514 at the August 5 evaluation. Kochia control did not exceed 67% with any treatment at the evaluation. Green foxtail control increased but kochia control decreased from the early to late evaluation which probably reflects kochia's greater drought tolerance.

S-23121 for broadleaf weed control in wheat, Fargo 1988. 'Wheaton' Hard Red Spring wheat was seeded to dry soil on May 16. The 2-leaf (2-lf) treatments were applied to cotyledonary to 2-leaf weeds on June 2 with 80 F, 60% RH, and no wind. The 5-lf treatments were to 4-to 5-leaf wheat, 3-to 4-leaf foxtail, 4-leaf redroot pigweed, and 2 to 6 inch tall kochia on June 10 with 75 F, 55% RH, and 12 mph wind. Treatments were applied in 8.5 gpa at 35 psi with a bicycle wheel type plot sprayer to an 8 ft wide area the length of 10 by 24 ft plots. The experiment was a randomized complete block with four replications and evaluated June 15, June 24, and July 27. Weeds exceeded 3 plants/sq. ft. except wild mustard stand was sparse and variable. Wheat stand was sparse and variable making the June 29 evaluations meaningless.

#### Summary

S-23121 gave greater control of weeds when applied to small weeds with the June 2 than the larger weeds with the June 10 applications, at the June 16 evaluation. Wild mustard control was not adequate with S-23121, but was controlled when 2,4-D or MCPA were added to S-23121. Redroot pigweed was the most suseptable to S-23121, of the weeds present. None of the treatments adequately controlled all weeds with the 5-leaf applications. Surfactant X-77 generally enhanced weed control with S-23121. The extremly dry conditions probably reduced weed control with all treatments.

Preemergence weed control in wheat, Williston 1988. Preplant incorporated (PPI) treatments were applied, and twice field cultivator incorporated, and 'Butte' Hard Red Spring wheat seeded on May 25 with 60 F, 25% RH, clear sky, 5 mph wind, and a dry soil surface. Preemergence incorporated (PEI) treatments were applied and twice harrow incorporated on May 26 with 56 F, 29% RH, clear sky, a 4 mph wind, and dry soil. Treatments were applied with a tractor-sky, a 4 mph wind, and dry soil. Treatments were applied with a tractor-mounted plot sprayer delivering 17 gpa at 35 psi to an 8 ft width the length of 10 by 25 ft plots. The experiment was a randomized complete block with four replications.

Tout tehticacions.									
						Std		Whea	
		Rate	Grft.	Rrpw	Ruth	Red	Inj	Tw	Yield
Treatment		(oz/A)	1%	contr	01)-	(%)	(%)	(1b/bu	) (bu/A)
		(02/1/	(,,,						
	PPI	0.25	33	84	73	0	0	52.7	3.2
Chlorsulfuron	PPI	0.50	58	95	80	0	0	53.1	5.3
Chlorsulfuron	PPI	5	59	20	5	53	11	47.4	2.2
Trifluralin	PEI	8	50	0	6	16	9	48.5	2.9
Trifluralin	PEI	0.25	0	89	55	3	0	50.9	3.0
Chlorsulfuron	PEI	0.23	18	95	65	0	3	51.3	3.7
Chlorsulfuron Tichunglin	PEI	0.25+8	58	93	56	6	5	50.1	3.2
Chlorsulfuron+Trifluralin	PEI	0.2310	0	0	0	0	0	62.2	3.6
Untreated		0							
			59	95	80	53	11	53.1	5.3
HIGH MEAN			0	0	. 0	0	0	47.4	2.2
LOW MEAN			34	59	43	10	3	50.8	3.4
EXP MEAN			67	27	44	89	77		36.2
C.V. %			34	24	28	13	4		NS
LSD 5%			46	32	38	17	5		NS
LSD 1%			4	4	4	4	4	1	4
# OF REPS									

#### Summary

Chlorsulfuron preplant incorporated gave greater green foxtail and tended to give greater Russian thistle control then when preemergence incorporated. None of the treatments gave adequate green foxtail control which may be a result of the drought. Chlorsulfuron gave 84% or more redroot pigweed control, regardless of incorporation method. Trifluralin at 5 or 8 oz/A applied alone preplant incorporated reduced wheat stand.

Small grain response to postemergence herbicides, Fargo 1988. The crop cultivars were seeded in adjacent 18 ft wide strips. The Alex cultivar was seeded untreated and treated with Mon 5500 at % w/w. Treatments were applied to 3- to 4-leaf crops on June 2 with 80 F, 60% RH and no wind. Treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa to an 8 ft wide area across the crops. Evaluation of crop injury was on June 28. The area was nearly free of weeds. The plots were 10 ft wide which gave a 4 ft area for reference in determining injury.

<u>Treatment</u>	<u>Rate Wh Alx</u> (oz/A)	A1x1 (%	r Cot   injury	Rug Ri	ot Wh	Alx AlxTr	Cot Rug Rbt
S-23121 Feno&2,4-D+MCPA DPX-R9674  HIGH MEAN LOW MEAN EXP MEAN C.V. % LSD 5% LSD 1% # OF REPS	0.23 0 0 12 0 0 0.75 0 0 0 0 0 0 0 0 0 0 NS NS NS NS NS NS	0 0 0 0 0 0 0 NS NS 3	0 3 3 0 1 1 300 6 NS 1 NS 2	0 0 32 7 0 0 1 2 9 300 6 NS	20.6 19.4 20.6 15.4 18.5 22.6 NS	22.2 17.9 25.4 21.4 23.9 21.4 25.4 21.4 22.2 17.9 23.9 20.2 5.7 13.0 NS NS NS NS 3 3	16.6 11.4 22.1 19.8 9.5 31.8 20.1 10.9 31.3 20.1 11.4 31.8 16.6 9.5 22.1 18.9 10.6 28.4 14.4 26.9 14.0 NS NS NS NS NS NS 3 3 3

a= Wh=Wheaton, Alx=Alex, AlxTr=Alex Treated, Cot=Coteau, Rug=Rugby durum, and Rbt=Robust barley;

## Summary

Fenoxaprop & 2,4-D & MCPA (Tiller) caused visual injury to Rugby durum. However, the yield of Rugby durum treated with Tiller was similar to that with other treatments which did not cause injury. The experiment did not contain an untreated treatment for yield because of limited space. The experiment was adjacent to "Small grain response to herbicides" experiment with yields from untreated areas which could be used for comparison.

CGA-131036 fall and spring applied on small grains, Williston 1988. Fall treatments were to bare summerfallow soil with 6.6 pH, and 1.7% organic matter on October 12, 1987 with 45 F, 65% RH, clear sky, 5 mph wind, and dry soil surface. The area was field cultivated prior to seeding 'Monroe'durum, 'Otana' oats, and 'Bowman' barley on May 13. Spring treatments were to 3- to 3.5-leaf crops, 1- to 5-leaf green foxtail, and less than 1 inch tall Russian thistle on June 1 with 61 F, 26% RH, clear sky, and a 4 mph wind. The treatments were applied with a tractor-mounted plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft wide area across the various crops. Evaluation was for weed control across the three crops. Harvest was from 64 sq ft area for each crop on August 3.

				Wh	eat_	0a	its_		<u>ley</u>
		Conft	Duth	TU	Viold	TW	Yield	TW	<u>Yield</u>
Treatment/Time	Rate	Grft		(1h/hu)	/hu/A)	(1h/hu)	(bu/A)	(1b/bu)	(bu/A)
	(gm/ha)	(% con	troi	(ID/Du)	(Du) A)	(10/04)	(24)		
					4.0	22.0	15.8	46.6	11.7
CGA-121036 Fall	7.5	0	5	60.5	4.2	32.0		45.1	12.0
Our t all a	10.0	0	9	59.9	4.4	30.2	17.6		
Our receive	20.0	0	25	60.0	4.4	31.4	20.6	47.3	13.8
CGA-121036 Fall		5	31	60.8	4.9	30.4	21.5	46.5	12.7
CGA-121036 Fall	30.0			60.8	5.1	30.7	20.4	47.0	15.0
Chlorsulfuron Fall	20.0	48	74		4.8	30.4	17.7	46.1	13.6
Control Fall	0.0	0	0	60.2			19.1	47.0	11.4
CGA-121036 Spring	7.5	0	40	61.5	5.1	31.5	the same of the sa	46.9	12.0
CGA-121036 Spring	10.0	5	38	60.4	4.6	29.8	19.0		13.3
		4	74	61.5	5.3	29.7	21.5	47.8	
Our. ===		13	84	60.9	3.9	31.4	19.9	46.3	11.4
CGA-121036 Spring		51	88	61.6	5.7	31.9	16.6	47.5	10.1
Chlorsulfuron Spring				60.8	5.2	32.3	17.5	46.3	10.5
Control Spring	0.0	0	0	00.0	3.2	52.0			
					22 4		20.2		25.3
C.V.%		90	46		33.4				NS
		14	26		NS		NS		
LSD 5%		18	35		NS		NS	Diest I	NS
LSD 1%		4	4	1	4	1	4	1	4
# Of Reps		4	100						

Summary

None of the treatment caused injury to any of the crops so individual treatment ratings are not presented in the table. Crop yield or seed test weight was not influenced by the herbicide treatments. Chlorsulfuron gave greater green foxtail control then CGA-121036 regardless if fall or spring applied. CGA-13036 gave greater Russian thistle control when applied postemergence in the spring then in the previous fall as a preemergence treatment and was similar to chlorsulfuron as a postemergence treatment.

Fenoxaprop and HOE-7113 in Durum and HRS wheat, Langdon 1988. 'Cando' durum and 'Butte 86' Hard Red Spring wheat were seeded on May 16. The wheat was seeded at 95 lb/A and contained 5 lb of 'Robust' barley as a bioassay species. Treatments were applied to 6 -leaf wheat on June 10. Herbicides were applied with a bicycle wheel type plot sprayer delivering 8.3 gpa at 40 psi through an 8001 nozzle. The soil in the area was variable which influenced yields. Two replications for fenoxaprop and three for untreated and HOE-7113.

		I.I.A.	Butte 8	6		Cando		_
<u>Treatment</u>	Rate (oz/A)	Wht inj (%)	yield (bu/A)	Tswt (1b/bu)	Wht inj (%)	yield	Tswt	
Fenoxaprop HOE-7113 Untreated	2.6	25 5 0	27.8 33.7 38.1	58.0 57.8 60.0	87 2 0	3.2 39.4 42.6	(1b/bu)  58.4 58.5	

### Summary

Data was not subjected to statistical analysis because there were only two replications for some treatment. However, fenoxaprop caused more injury to Cando durum than Butte 86. HOE-7113 (Puma) did not cause important injury to either Butte 86 Hard Red Spring wheat or to Cando durum. The area was relatively free of weeds.

Harrowing for weed control, Langdon 1988. 'Coteau' Hard Red Spring wheat was seeded on May 6. The wild oats, wheat, and green foxtail were all at similar leaf stage for a given harrowing treatment. Harrowing was with 10 ft wide Melroe flex tine harrow at 4 mph. The herbicide treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi to a 6.7 ft wide area. The entire area was treated with chlorsulfuron at 0.19 oz/A + 2,4-D at 4 oz/A on June 12. Plots were 12 ft wide. Weed control evaluation on July 26.

<u>Treatment <sup>a</sup></u>	Rate	Inj	Wioa	Grft		Tswt (lb/bu)
Treatment	(oz/A)	(%)	(% COI	11101	(Du) N)	(15/24)
Diclofop+DPX-M6316+PO	16+0.25	0	72 0	45 0	15.5 11.1	58.6 58.3
Check 1 harrow at right angle to rows(2.5 leaf) 1 harrow at right angle to rows(3 leaf)		26 18 21	36 34 36	63 70 60	11.7 11.8 12.0	57.9 57.0 58.1
1 harrow at right angle to rows(4 leaf) 1 harrow at right angle to rows(5 leaf) 1 harrow with rows+dcfp+DPX-M6316+PO(2.51f)	16+0.25	41	46 83	53 64	8.4 14.3	56.4 59.5 58.5
1 harrow with rows(2.51f) 1st with rows(2.5 lf)then across(2.51f)		19 35 46	39 54 53	53 46 64	11.3 10.5 10.2	57.0 56.8
1st with rows(2.5 lf)then across(3lf) 1st with rows(2.5 lf)then across(4lf) 1st with rows(2.5 lf)then across(5lf)		46 39	39 56	53 44	9.2 9.6	56.3 57.3
HIGH MEAN		46	83 0	70 0	15.5	59.5 56.3
LOW MEAN EXP MEAN C.V. %		26 58	46 32	51 32 23	11.3 24.3 4.0	57.6 1.9 1.6
LSD 5% LSD 1%		22 29 4	21 28 4	31		2.1
# OF REPS						

a PO=petroleum oil aft 1 oz/A.

#### Summary

Wheat yield was not influenced by any of the treatments for weed control, regardless of mechanical chemical or both mechanical and chemical. However the harrowing treatments generally injured the wheat. The various times or directions of harrowing did not greatly influence wild oats or green foxtail control, but were generally as effective as diclofop + DPX-M316 in controlling green foxtail. Wild oats control tend to be the highest with the harrowing + herbicide treatment.

Green foxtail control in wheat, Minot 1988. An experiment was conducted to evaluate herbicides for green foxtail control in Hard Red Spring wheat on a Williams loam soil with a pH 6.6 and 4% OM. Preplant incorporated (PPI) treatments were applied and incorporated into the top 2 inches of soil with a S-tine cultiweeder, 'Stoa" Hard Red Spring wheat was seeded, preemergence S-tine cultiweeder, 'Stoa" Hard Red Spring wheat was seeded, preemergence incorporated (PEI) treatments were applied to moist soil and incorporated 2 X with a spring tooth harrow and preemergence (PE) treatments were applied to the soil surface on May 13. The first rain after treatment was 1.06 inch on June 14. The postemergence (S1) treatments were applied to 3.0 leaf wheat, 0.5 to 2.0 inch green foxtail, and 0.5 to 1.0 inch common lambsquarter and Russian thistle. on June 1 with cloudy sky and 75 F. The (S2) treatments were applied to 5 leaf wheat, 0.5 to 4.0 inch green foxtail, and 1.0 to 4.0 inch common lambsquarter and Russian thistle on June 13 with clear sky and 80 Treatments were applied with a bicycle wheel type plot sprayer delivering either 17 or 8.5 gpa at 35 psi for the soil applied or postemergence treatments, respectively. Stand reduction for the preemergence treatments was evaluated on June 7 and green foxtail, common lambsquarter and Russiam thistle control was evaluated on June 28 and on July 21. The experiment was a randomized complete block design with four replications and experimental units were 10 X 18 sq ft. A 4 X 18 sq ft area of wheat was harvested for yield on August 4.

	~								
Treatment		Rate		Wheat td.red.	In.j	Grf: 6/28	-	C-1.	D 11
		(oz/A)	(bu/A)	(%)	(%)	(%	7/21 % con	Colq trol)	<u> </u>
Chlorsulfuron Chorsulfuron Trifluralin Clsu+Trif Clsu+Trif Chlorsulfuron Chlorsulfuron Clsu+R11 Clsu+Dicl Diclofop H0E7113-04H H0E7113-04H H0E7125-02H Check C.V. %	(PPI) (PPI) (PEI) (PEI) (PE) (PE) (S1) (S1) (S1) (S2) (S2) (S2)	0.25 0.50 8.0 0.25+8 0.25+16 0.25 0.50 0.25+.25% 0.25+12 12.0 1.28 1.28 5.25 6.25	6.3 6.5 8.7 8.3 6.9 5.5 4.5 5.2 4.0 6.6 5.0 4.5 3.9	1 0 4 0 5 0 0 0 0 0 0	1 0 0 0 1 3 1 3 8 3 5 10 11 10 0	39 64 66 75 88 13 18 50 63 59 89 89 81 85 0	30 60 66 79 92 19 16 38 54 48 73 81 75 75	80 88 56 79 70 51 64 98 98 1 0 91 93 0	61 66 48 70 55 53 64 98 95 1 0 84 93 0
LSD 5%			34.7 2.9	314	106	25 21	28	20	30
a Rll = surfact	tant with	90% ai;			0		22	17	22

Summary The soil applied treatments caused minimal Chlorsulfuron + diclofop, HOE 7113-04H (S2) wheat stand significant wheat injury but only tended to reduce wheat grain yields. Wheat treated with trifluralin (PEI) at 8 oz/A alone yielded significantly more than and HOE 7125-02H caused the check and several other treatments despite only fair green foxtail and broadleaf weed control. Chlorsulfuron applied postemergence gave the highest control of common lambsquarter and Russian thistle while chlorsulfuron + trifluralin, HOE 7113-04H and HOE 7125-02H gave the highest green foxtail control. Green foxtail control was less with surface applied chlorsulfuron (PE) compared with other applications methods. Minimal rainfall and excessive temperatures during May and June following planting probably accounted for the generally poor weed control.

Weed control in flax, Fargo 1988. 'Linton' flax was seeded to dry soil on May 17. Treatments were applied to 3- to 8-inch tall flax, 2- to 6-leaf redroot pigweed, kochia, wild mustard, and foxtail on June 7 with 85 F, 48% RH and 10 mph wind at application. The second treatment of splits (/) was on June 10 with 75 F, 60% RH, and 10 mph wind at application. Plants were within the stages given for the first treatments. All treatments were applied in 8.5 gpa at 35 psi with a bicycle wheel type plot sprayer to an 8 ft wide area the length of 10 by 24 ft plots. The experiment was randomized complete block with four replications. Evaluations were on June 29 and July 18. Green foxtail density was about 1 plant/sq ft and other weeds about 5/sq ft.

TOXEATT delistey has assure				- 00			17 1	10
			June	e 29			July 1	Flax
		Flax	0 61			Coet	Dwnu	inj
Treatment	Rate	inj	Grtt	Rrpw	Willia	Grft	KIPW	
	(oz/A)	(%)		(%	conti	ro1)		(10)
Diclofop+Bromoxynil Diclofop+Bromoxynil+PO Diclofop+Bromoxynil+MS Rent+MCPA+PO/Seth+MS 84	12+4 12+4+0.12G 12+4+0.12G 12+4+0.12G -4+0.25G/2+0.25G 2+0.25G/4 2+0.25G/4 2+4+0.25G 2+4+0.25G 2+4+0.25G/4 2+4+0.25G/4 2+8+0.25G 2+8+0.25G 2+8+0.25G 2+8+0.25G	(%) 10 9 5 12 8 25 26 18 10 30 34 6 13 4	95 99 99 95 97 97 97 99 95 95 95	85 72 66 74 43 80 89 79 75 87 88 83 91 65	98 99 98 99 99 99 99 99 99 99	97 98 99 99 99 99 99 99 99 98 98 99	64 48 36 82 49 74 77 78 81 93 94 90 94	(%) 2 0 2 9 7 11 17 2 6 17 34 2 1 0
Sethoxydim+Bromoxynil+MS Sethoxydim+Brox&MCPA+BCH Sethoxydim+Brox&MCPA+MS Untreated	2+4+0.25G 2+8+0.25G 2+8+0.25G 0	9 8 5 0	95 98 95 0	46 90 79 0	92 99 99 0	99 99 99 0	21 78 67 0	0 0 0
C.V. % LSD 5% LSD 1% # OF REPS		59 11 14 4	5 6 8 4	23 23 31 4	3 4 5 4	1 2 3 4	16 15 20 4	127 11 15 4

#### Summary

The MCPA bee (butoxyethyl ester) injured flax when applied on June 10 following a June 7 application of sethoxydim or sethoxydim + MCPA bee with BCH (Dash adjuvant from BASF) or MS (Sun-it methylated seed oil + emulsifier). Sethoxydim + MCPA bee with these adjuvants did not injure flax treated on June 7. Green foxtail was controlled by all herbicide treatments. MCPA at 8 oz/A was equal to MCPA as a 4+4 oz/A split application for redroot pigweed control, with either BCH or MS.

Weed control in flax, Carrington 1988. 'Linton' flax was seeded on May 10. Treatments were applied on June 9 except for second applications of split (/) treatments which were on June 15. Conditions for 1 week after treatment were 84 to 96 F and drought stress. Flax was 5- to 6-inch tall, green foxtail 2- to 3-leaf, wild oats 6-leaf, and broadleaf weeds 3 inch tall at treatment. All treatments were applied in 8.5 gpa at 35 psi with a bicycle wheel type plot sprayer to an 8 ft wide area the length of 8 by 20 ft plots. The experiment was randomized complete block with four replications. Evaluation was on July 15. Green foxtail density exceeded 6 plants/sq ft, but other species were less dense and variable.

<u>Treatment</u>	Do.L.	Flax			
	Rate	inj	Wioa	Grft	Rrpw
	(oz/A)	(%)	(%	contro	1)
Sethoxydim+BCH/MCPA-bee Sethoxydim+MS/MCPA-bee Sethoxydim+MCPA-bee+BCH Sethoxydim+MCPA-bee+MS Seth+MCPA-bee+BCH/MCPA-bee	12+4 12+4+0.12G 12+4+0.12G 4+0.25G/2+0.25G 4+0.25G/4 2+0.25G/4 2+4-0.25G 2+4+0.25G 2+4+0.25G/4 2+8+0.25G 2+8+0.25G 2+8+0.25G 2+8+0.25G 2+8+0.25G 2+8+0.25G 2+8+0.25G	1 0 5 0 6 3 0 0 4 3 9 0 9 3 6 1 165 NS NS NS NS NS NS NS NS NS NS NS NS NS	54 29 51 8 9 34 23 16 15 19 10 3 9 15 5 11	61 65 69 10 39 48 58 50 49 51 66 60 54 33 35 18 44 38 24 32 4	50 53 67 33 45 67 53 57 63 77 88 82 65 27 35 55 45

### Summary

None of the herbicides caused any important injury to flax or gave adequate control of wild oats and green foxtail. The drought stress and 300 ppm sodium bicarbonate in the spray water may have contributed to the lack of grass species control with sethoxydim. BCH (adjuvant from BASF-Dash) and MS (methylated seed oil adjuvant Sun-it) were similar as adjuvants with the various herbicide treatments. MCPA butoxyethyl ester (bee) at 4+4 oz/A split application did not exceed MCPA bee at 8 oz/A as a single application for redroot pigweed control.

Weed control in flax, Williston 1988. 'Flor' flax was seeded on May 13. Treatments were applied to 3 inch tall flax, 3-leaf to tillering green foxtail, 2- to 4-inch tall Russian thistle, and other broadleaf weeds 2- to 4-inch tall on June 9 with 69 F, 35% RH, clear sky, and a 7 mph wind. High inch tall on June 9 was 90 F. The second treatments following the / were on June 13 with 58 F, 55% RH, cloudy sky, and a 5 mph wind. The temperature for June 13 was 81 F. Treatments were applied with a tractor mounted plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft wide area the length of 10 by 25 ft plots. The experiment was a randomized complete block with four replication. Weed infestations were dense for Russian thistle, moderate for green foxtail and common lambsquarters, and sparse for redroot pigweed and kochia. Evaluation was on July 18. Harvest was from 80 sq ft on August 16.

<u>Treatment</u>	Rate (oz/A)	Grft	Ruth (%	Rrpw	Colq	KOCZ	Inj (%)	Flax Yield (bu/A)(	<u>Tswt</u> 1b/bu)
Diclofop+Bromoxynil Diclofop+Bromoxynil+PO Diclofop+Bromoxynil+MS Bent+MCPA+PO/Seth+MS Bent+Bromoxynil+PO/Seth+MS Sethoxydim+BCH/MCPA-bee Sethoxydim+MS/MCPA-bee Sethoxydim+MCPA-bee+BCH Sethoxydim+MCPA-bee+MS Seth+MCPA-bee+BCH/MCPA-bee Seth+MCPA-bee+MS/MCPA-bee Sethoxydim+MCPA-bee+BCH	12+4 12+4+0.12G 12+4+0.12G 8+4/2 8+4/2 2/4 2/4 2/4 2+4	15 31 26 11 0 59 94 94 92 94 90	71 86 81 15 70 18 0 35 39 55 33 49	61 56 56 33 40 33 31 66 58 72 48 91	83 85 90 71 93 90 93 86 94 95 94	63 94 78 0 60 54 70 46 39 95 71	4 3 3 0 0 8 0 0 8 10 10 6	1.8 2.4 1.6 1.5	41.7 51.7 51.2 51.5 50.8 50.8 52.5
Sethoxydim+MCPA-bee+MS Sethoxydim+Bromoxynil+BCH Sethoxydim+Bromoxynil+MS Seth+Bromoxynil&MCPA+BCH Seth+Bromoxynil&MCPA+MS Untreated	2+8 2+4 2+4 2+8 2+8 0	93 95 83 95 93 0	66 81 83 74 88 0	34 81 35 49 41 0	95 94 84 95 93	78 85 80 84 59	6 4 6 5 3 0	3.1 3.9 4.5	51.6 51.8 50.4
C.V. % LSD 5% LSD 1% # OF REPS		12 11 15 4	33 24 32 4	55 40 53 4	17 20 27 4	48 43 58 4	129 NS NS 4	22.5 1.0 1.3 4	1

a Missing data was for plots not harvested because of excessive weed competition

Summary

None of the treatments caused important injury to flax. Bentazon as a treatment 4 days before sethoxydim antagonized green foxtail control. Treatments with bromoxynil gave the greatest Russian thistle control and control tended higher when bromoxynil was with methylated seed oil petroleum oil alone. Redroot pigweed and kochia control was too variable to determine differences among treatments.

Weed control in flax, Minot 1988. 'Neche' flax was seeded on May 6. Treatments were applied to 3 inch tall flax, 1- to 3-inch tall kochia, 1- to 4-inch diameter Russian thistle, and 0.5- to 2.5-inch tall green foxtail on June 1 with 70 F, cloudy sky, and no wind. The second treatment of the split applications (after 1) were on June 3. All treatments were applied in 8.5 gpa at 35 psi with a bicycle wheel type plot sprayer to an 8 ft wide area the length of 10 by 24 ft plots. The experiment was randomized complete block with four replications. Evaluation was on July 14. Weed densities exceeded 5 plants/sq ft for all species.

Treatment		Flax			
	Rate	inj	Grft	Ruth	KOCZ
	(oz/A)	(%)		% cont	roll
Diclofop+Bromoxynil					. 01)
Diclofop+Bromoxynil+PO	12+4	3	19	83	55
Diclofop+Bromoxynil+MS	12+4+0.12G	0	23	88	49
Bentazon+MCPA+PO/Sethoxydim+MS	12+4+0.12G	0	26	86	44
Bentazon+Bromoxynil+PO/Sethoxydim+MS	8+4+0.25G/2+0.25G	0	77	79	48
Sethoxydim+BCH/MCPA-bee		0	43	89	88
Sethoxydim+MS/MCPA-bee	2+0.25G/4	0	80	20	31
Sethoxydim+MCPA-bee+BCH	2+0.25G/4	0	83	30	35
Sethoxydim+MCPA-bee+MS	2+4+0.25G	0	68	39	44
Sethoxydim+MCPA-bee+BCH/MCPA-bee	2+4+0.25G	0	75	49	50
Sethoxydim+MCPA-bee+MS/MCPA-bee	2+4+0.25G/4	0	86	66	46
Sethoxydim+MCPA-bee+BCH	2+4+0.25G/4	0	83	76	48
Sethoxydim+MCPA-bee+MS	2+8+0.25G	4	70	68	59
Sethoxydim+Bromoxynil+BCH	2+8+0.25G	6	62	74	61
Sethoxydim+Bromoxynil+MS	2+4+0.25G	0	70	82	64
Sethoxydim+Bromoxynil&MCPA+BCH	2+4+0.25G	0	81	85	63
Sethoxydim+Bromoxynil&MCPA+MS	2+8+0.25G	0	80	88	63
Untreated	2+8+0.25G	4	72	94	65
	0	0	0	0	0
C.V. %					
LSD 5%		275	27	19	38
LSD 1%		4	23	18	27
# OF REPS		NS	31	24	36
		4	4	4	4

#### Summary

None of the herbicides caused injury to flax. Green foxtail control was greater with sethoxydim than with diclofop treatments. BCH (adjuvant from BASF-Dash) was similar to MS (methylated seed oil adjuvant Sun-it) for grass species control with sethoxydim. Bromoxynil and Bentazon treatments tended to give the greatest Russian thistle control. MCPA bee (butoxyethyl ester) 4+4 oz/A as a split application was simuliar to MCPA bee at 8 oz/A as one application for controlling Russian thistle and kochia. Russian thistle control tended to be greater with treatments containing MS than BCH adjuvant. Kochia control was the greatest with the Bentazon plus bromoxynil

Acifluorfen for Flax desiccation, Casselton 1988. Treatment were applied on August 17 with 75 F, 85% RH, and 10 to 15 mph wind. Treatments were to mature 'Foster' flax with green leaves and stems, but no seed balls because of drought and grasshoppers. The 17 gpa treatments were with 8022 and 8.5 gpa with 8001 nozzles on a bicycle wheel type plot sprayer using 35 psi CO2 pressure. Treatments were to 8 ft wide area the length of 30 by 10 ft plots. Evaluation for percut plant desiccation was on August 24 and September 2. The experiment was a randomized complete block with three replications.

Treatment	Spray volume (gpa)	Rate (oz/A)	August Flax (%	24 Se Colq % desid	Flax ccatio	Colq
Acifluorfen+MS Acifluorfen+MS Acifluorfen+MS Acifluorfen+Dash Acifluorfen+MS Acifluorfen+MS Acifluorfen+MS Paraquat+X77  C.V. % LSD 5% LSD 1% # OF REPS	(8.5) (8.5) (8.5) (8.5) (17) (17) (17) (17)	2+0.25G 4+0.25G 8+0.25G 4+0.25G 2+0.25G 4+0.25G 8+0.25G 8+0.25%	94 92 99 89 93 91 98 99 4 7 NS 3	0 0 22 33 0 0 3 77 122 36 50 3	80 81 97 78 77 78 92 98 9 14 NS 3	12 13 17 33 0 0 13 73 114 40 NS 3

### Summary

Acifluorfen at 8 oz/A applied with methylated seed (MS) oil was equal to paraquat for desiccation of flax. The differences between the two evaluations may be because of different evaluators as plants probably did not recover with time. The BCH (Dash from BASF) tended to cause less enhancement than MS for flax and less for common lambsquarters desiccation.

Dessication of flax, Minot 1988. 'McGregor' flax was seeded in mid delivering 8.5 or 17 gpa at 35 psi to 18 to 24 inch flax which was near physicalogical maturity on September 14 with 55 F, a sunny sky and no Evaluations of flax dry down were made on September 23 and October 3.

Treatment		Rate	Eval 1	Eval 2
Aciflourfen+MS Aciflourfen+MS Aciflourfen+MS Aciflourfen+Dash Aciflourfen+MS Aciflourfen+MS Aciflourfen+MS Aciflourfen*MS C.V. %	(8.5 gpa) (8.5 gpa) (8.5 gpa) (8.5 gpa) (17 gpa) (17 gpa) (17 gpa) (17 gpa)	2+0.25G 4+0.25G 8+0.25G 4+0.25G 2+0.25G 4+0.25G 8+0.25G	(% dess 15 15 17 15 12 10 12 32	
LSD 5% # OF REPS F-TRT			29 8 3	18 7
a *MS = The MS was	not include	d in the treatme	6	3 10

a \*MS = The MS was not included in the treatment.

## Summary

Treatments did not adequately dessicate flax with the cool conditions of mid to late September. NaClO3 matured the flax bolles quicker than the other treatments however it did not kile the flax plant.

Weed control in flax, Langdon 1988. 'McGregor' flax was seeded on May 10. Treatments were applied to 3 inch tall flax and ladysthumb on June 9, except for the second application of the split (1) treatment which was on June 13. for the second application of the split (1) treatment which was on June 13. All treatments were applied in 8.5 gpa at 35 psi with a bicycle wheel type plot sprayer to an 8 ft wide area the length of 10 by 20 ft plots. The plot sprayer to an 8 ft wide area the length four replications. Evaluation experiment was randomized complete block with four replications. Evaluation was on July 20. Ladysthumb density exceeded 1 plant/sq. ft.

was on July 20. Ladysthumb density  Treatment	Rate (oz/A)	Flax l	Ladys- thumb control	Yield F )(lb/A)	-lax '	ID/A)
Sethoxydim+Bromoxynil+MS Sethoxydim+Bromoxynil&MCPA+MS Sethoxydim+MCPA-bee+MS/MCPA(1-3d) Sethoxydim+MCPA-bee+MS Sethoxydimoxydim+MS Diclofop+Bromoxynil Picloram+MCPA-dma Picloram+MCPA-dma Picloram+MCPA-dma+Sethoxydim+MS Metsulfuron+X-77 DPX-R9674+X-77 DPX-M6316+MCPAdma	3+4+0.25G 3+8+0.25G 3+4+0.25G/4 3+8+0.25G 3+0.25G 12+4 0.25+4 0.25+4+3+0.25% 0.05+0.25% 0.25+0.25% 0.15+4	0 0 5 4 0 0 1 3 6 20 7	38 53 45 53 25 5 48 52 96 76 88 0	590 873 914 909 950 1003 938 840 762 597 651 1015	68 60 55 48 39 38 65 45 98 64 94 31	393 524 497 436 368 380 606 380 742 389 614 320
Untreated  C.V. % LSD 5% LSD 1% # OF REPS	sugal yiz ni a	113 6 8 4	33 21 29 4	16 188 253 4	15 13 17 4	22 148 200 4

# Summary

DPX-R9674 caused 20% injury to flax and gave 76% ladysthumb control. However, metsulfuron and DPX-M6316+MCPA gave 96 and 88% ladysthumb control respectively without causing important injury to flax. None of the other herbicide treatments gave adequate ladysthumb control. The percent flax in the harvested treatments gave adequate ladysthumb control.

Flax variety response to sulfonyl urea herbicides, Minot 1988. Flax was seeded May 10 on fallow which had been fertilized for a 50 bu/A wheat yield goal. Sethoxydim 4.5 oz/A + Sun-it oil at 1 qt/A was applied for grass weed control. DPX-L5300 at 0.5 oz ai/A and DPX-R9674 at 0.9 oz ai/A were applied with a bicycle wheel type plot sprayer delivering 8.5 metsulfuron was applied with the same method to 2.5 inch flax with a sunny sky and 90 F on June 3. All sulfonyl urea herbicides were applied with 0.25% V/V R-11 surfactant. Injury evaluations were made on June 17. The treatments were not replicated and data represent one observation.

Variety ————————————————————————————————————	DPX-L5300	Flax injury DPX-R9674	Metslufuron
U 605 U 605 Clark Flor Linton McGregor Neche NorLin NorMan Verne	55	60	25
	50	60	25
	60	50	40
	60	50	50
	80	80	40
	40	40	25
	80	70	55
	55	60	30
	80	75	45
	50	50	30

## Summary

All flax varieties were injured initially by the sulfonyl urea herbicides. Metsulfuron caused the least injury of the herbicides tested. Flax was stunted, yellowed, and delayed in maturity 2 to 3 weeks. Late June and early July rains resulted in vigorous branching potential. The plots were not harvested for yield.

Weed control in sunflower, Carrington 1988. Preplant incorporated (ppi) treatments were applied and rototiller incorporated into dry soil and 'Sigco treatments were applied on June 6. Preemergence (PE) treatment was applied on June 7. The first rain after treatment occurred on June 12. The first rain after treatment occurred on June 12. Postemergence (p1) treatments were to 2- to 8-leaf sunflower, green foxtail Postemergence (p1) treatments were to 2- to 8-leaf sunflower, green foxtail reatments were applied on July 5. Postemergence (p2) 3-leaf and 3 inch tall redroot pigweed on July 5. Postemergence (p2) treatments were applied on July 7. Herbicides were applied with a bicycle treatments were applied on July 7 gpa for ppi and pe treatments and 8.5 wheel type plot sprayer delivering 17 gpa for ppi and pe treatments and 8.5 wheel type plot sprayer delivering 17 gpa for ppi and pe treatments and 8.5 the experiment was a randomized complete block with four replications. The experiment was a randomized complete block with four replications. Sunflowers variable in size and weed variable in Evaluated July 15. Sunflowers variable in size and weed variable in varia

density. <u>Treatment</u>	Rate (oz/A)	Sufl inj (%)	Grft (% cor	
EPTC (ppi) Trifluralin (ppi) Ethalfluralin (ppi) Pendimethalin (ppi) Pendimethalin+AC 222,293 (ppi) Trifluralin+Chloramben-W (ppi) EPTC+Chloramben-W (ppi) Trifluralin+RE40885 (ppi) Alachlor(pe) Pendimethalin(ppi)/AC 222,293(p1) Pendimethalin(ppi)/AC 222,293(p2) Pendimethalin(ppi)/Aciflurofen(p1) Pendimethalin(ppi)/Aciflurofen(p1) Pendimethalin(ppi)/Aciflurofen(p1) Sethoxydim+AC 222,293+MS(p1) Sethoxydim+Aciflurofen+MS(p1)  C.V. % LSD 5% LSD 1% # OF REPS	40 12 12 20 20+5 12+32 40+32 12+8 40 20/3 20/3 20/3 20/2 20/3 3+3+0.25G 3+2+0.25G	35 0 0 0 0 0 3 0 0 3 20 4 15 15 45	99 98 98 98 99 98 98 93 89 99 99 98 68 60	99 98 98 99 98 99 95 97 99 99 28 88

## Summary

EPTC (ppi), AC 222,293 (p2), and aciflurofen + MS (p1) MS=(methylated seed oil, Sun-it) all caused 20% or more injury to sunflower. All preplant incorporated and preemergence herbicide treatments gave 89% or more control of green foxtail and redroot pigweed.

Weed control in sunflower, Minot 1988. Preplant soil incorporated (ppi) treatments applied and twice (first immediately and second after 20 minutes) 6440 sunflower seeded and preemergence (pe) treatment applied on May 26. Pl just emerging to 1 or 2 true leaves on June 22 with 75 F, high humidity, and 2- to 8-inch tall weeds in early bud stage on July 8 with 70 F, 10 mph wind, and good growing conditions. Treatments were applied to an 8 ft wide area delivering 17 gpa for the ppi and pe and 8.5 gpa for the postemergence block with four replications. Evaluation was on July 18. All weeds densities were greater then 1 plant/sq ft.

Rate	Sufl ini	Grft	Rrpw	Rrpw	Colq	Tamu
(oz/A)	(%)		(% (	control	)	
12+32 40+32 12+8 40 20/3 20/3 20/2 20/3 +3+0.25G	0 0 1 0 0 15 8 5 0 0 16 0 1 2	97 98 98 98 97 98 97 78 96 98 97 91 88 0	96 97 99 97 99 98 99 86 99 98 99 60 80	92 98 99 95 97 98 97 85 95 94 97 99 34 81 0	91 93 99 89 98 96 97 98 61 99 93 93 97 41 65 0	86 0 16 5 98 89 95 94 25 99 85 56 77 99 95 0
	232 NS NS 4	3 4 5 4	5 7 9 4	10 12 15 4	16 18 24 4	19 18 24 4
	(oz/A)  40 12 12 20 20+5 12+32 40+32 12+8 40 20/3 20/3 20/3 20/2 20/3 +3+0.25G 2+0.25G	inj (oz/A) (%)  40 0 12 0 12 1 20 0 20+5 0 12+32 15 40+32 8 12+8 5 40 0 20/3 0 20/3 16 20/2 0 20/3 1 +3+0.25G 2 2+0.25G 11 d 0  232 NS NS	inj (oz/A) (%) 40 0 97 12 0 98 12 1 98 20 0 97 12+32 15 98 40+32 8 98 12+8 5 97 40 0 78 20/3 0 96 20/3 16 98 20/2 0 98 20/3 16 98 20/2 0 98 20/3 1 97 +3+0.25G 2 91 21 88 d 0 0	inj  (oz/A)  (%)  40  12  0  98  97  12  1  98  99  20  20+5  0  12+32  15  98  99  12+32  15  98  99  40+32  18  8  98  98  12+8  5  97  99  20/3  20/3  20/3  20/3  20/3  20/3  20/3  16  98  98  98  98  98  20/3  20/3  16  98  98  98  98  98  99  20/3  16  98  98  99  20/3  1  97  99  20/3  1  97  99  20/3  1  97  99  20/3  1  97  99  20/3  1  97  99  20/3  1  97  99  20/3  1  35  NS  4  7  NS  5  9	in,j  (oz/A)  (%)	inj (oz/A) (%)(% control)  40

#### Summary

AC 222,293 (Assert) caused greater injury at the second postemergence then the first application. Green foxtail control was 88% or more with all 84% with EPTC, all AC 222,293 treatments, RE-40885, and aciflurofen+MS pigweed, and common lambsquarter were adequately (>85%) controlled by all for common lambsquarter.

Weed control in sunflower, Williston 1988. Preplant soil incorperated (ppi) treatments were applied and field cultivator incorporated twice on May 18 with 60 F, 40% RH, dry soil and a 7 mph wind. 'Pioneer 6440' sunflower was with 60 F, 40% RH, dry soil and a 7 mph wind. 'Pioneer 6440' sunflower was with 60 F, 40% RH, dry soil and a 7 mph wind. 'Pioneer 6440' sunflower was RH, seeded and preemergence (pe) treatment applied on May 19 with 62 F, 18% RH, seeded and preemergence (pe) treatments were applied to dry soil, and a 6 mph wind. Postemergence (mostly 6 leaves), 3-leaf to tillering green foxtail, 0.5 inch tall redroot pigweed, and 1 to 3 inch tall Russian foxtail, 0.5 inch tall redroot pigweed, and 1 to 3 inch tall Russian foxtail, and tame yellow mustard (Tamu) on June 13 with 64 F, 45% RH, and 7 thistle, and tame yellow mustard (Tamu) on June 13 with 64 F, 45% RH, and 7 thistle, and tame yellow mustard, 4 to 6 inch tall redroot sunflowers, 6 inch tall to flowering tame mustard, 4 to 6 inch tall redroot sunflowers, 6 inch tall to flowering tame mustard, 4 to 6 inch tall redroot sunflowers, 6 inch tall to flowering tame mustard, 4 to 6 inch tall redroot sunflowers, 6 inch tall to flowering tame mustard, 4 to 6 inch tall redroot sunflowers, 6 inch tall to flowering 17 green foxtail on June pigweed and Russian thistle, and 2-leaf to tillering green foxtail on June 22 with 68 F, 60% RH, cloudy sky, dry soil, and no wind. Herbicides were applied with a tractor-mounted plot sprayer delivering 17 gpa at 35 psi for the postemergence the preplant and preemergence and 8.5 gpa at 35 psi for the postemergence the preplant and preemergence and 8.5 gpa at 35 psi for the postemergence treatments. Treatment was to an 8 ft area over the two center rows of four row plots the 25 ft length of the plots. Weed control evaluation was on July 24.

July 24.						
Treatment	Rate (oz/A)	Sufl inj (%)	<u>Grft Ru</u> (%	uth Ta	amu Rr	<u>^pw</u> 
EPTC(ppi) Trifluralin(ppi) Ethalfluralin(ppi) Pendimethalin(ppi) Pendimethalin+AC 222,293(ppi) Trifluralin+Chloramben-W(ppi) EPTC+Chloramben-W(ppi) Trifluralin+RE-40885(ppi) Alachlor(pe) Pendimethalin(ppi)/AC 222,293(p1) Pendimethalin(ppi)/AC iflurofen(p1) Pendimethalin(ppi)/Aciflurofen(p1) Pendimethalin(ppi)/Aciflurofen(p1) Sethoxydim+AC 222,293+MS(p1) Sethoxydim+Aciflurofen+MS(p1)	40 12 12 20 20+5 12+32 40+32 12+8 40 20/3 20/3 20/2 20/3 3+3+0.25G 3+2+0.25G	4 0 1 0 1 0 0 1 0 1 0 3 4 5 8	95 89 95 36 79 96 97 96 63 91 94 91 94 89 0	24 60 65 20 66 91 78 92 8 75 80 64 76 73 79 0	55 11 19 0 56 33 85 43 0 94 96 79 71 95 80 0	90 99 94 35 95 97 96 94 24 91 89 90 43 85 0
Untreated  C.V. % LSD 5% # OF REPS		202 NS 4	9	30 25 4	39 28 4	17 28 4

## Summary

None of the herbicides caused important injury to sunflower. AC 222, 293 at 3 oz/A applied postemergence gave more than 90% control of tame mustard, but only 56% when applied preplant incorporated at 5 oz/A. Green foxtail only 56% when applied preplant incorporated at 5 oz/A. Green foxtail only 56% when sethoxydim tended to be less when sethoxydim was with aciflurofen than AC 222,293. The low level of weed control with aciflurofen than AC 222,293. The low level of weed control with experiments control of most weeds has been similar to that with trifluralin. Aciflurofen was similar to AC 222, 293 for Russian thistle control; more aciflurofen was similar to AC 222, 293 for tame mustard control; and less effective than acceptable accounts of the control of the con

Weed control in sunflower, Casselton 1988. Preplant incorporated (ppi) treatments were applied and twice incorporated with a field cultivator plus harrow 3 inch deep in mallow soil on May 3. 'Interstate' 3001 sunflower applied to 2- to 10-leaf sunflower, 6- to 10-inch tall wild mustard, 10 and 10 mph wind. Patreatment was on June 20 with 88 F, 40% RH, an 8 ft wide area the length of 10 by 25 ft plots with a bicycle wheel type postemergence treatments. The experiment was a randomized complete block untreated areas because of drought, July 5. Weed density was about 1 plant/sq ft for each species.

Treatment	Rate	Sufl inj	Fxtl	KOCZ	Wimu
EPTC (ppi)	(oz/A)	(%)	(%	contro	1)
Trifluralin(ppi) Ethalfluralin(ppi) Pendimethalin(ppi) Pendimethalin+AC 222,293(ppi) Trifluralin+Chloramben-W(ppi) EPTC+Chloramben-W(ppi) Trifluralin+RE40885(ppi) Alachlor(pe) Pendimethalin(ppi)/AC 222,293(p1) Pendimethalin(ppi)/AC 222,293(p2) Pendimethalin(ppi)/Aciflurofen(pli) Pendimethalin(ppi)/Aciflurofen(pli) Sethoxydim+AC 222,293+MS(p1) Sethoxydim+Aciflurofen+MS(p1)	40 12 12 0 20+5 12+32 40+32 12+8 40 20/3 20/3 20/3 20/2 20/3 3+3+0.25G 3+2+0.25G	0 0 0 0 0 0 0 0 0 3 6 3 1 30	94 98 99 96 94 97 95 49 94 97 96 99 75	73 91 92 84 78 90 92 89 20 87 86 71 87 30 63	33 10 21 15 70 88 84 79 15 94 84 85 87 96 99
SD 5% SD 1% OF REPS		95 5 7 4	8 10 13 4	15 15 20 4	24 20 27 4

# Summary

Aciflurofen and AC 222,293 applied with MS (methylated seed oil adjuvant, Sun-it) injured sunflower. Green foxtail control exceed 90% with all with aciflurofen + MS gave 92% green foxtail control. Either AC 222,293 was antogonistic to sethoxydim or aciflurofen enhanced green foxtail control. Wild mustard control exceeded 80% with all postemergence mustard control even though emulsion instability occurred when mixed with trifluralin. Trifluralin and ethalfluralin were the only herbicides to give more than 90% kochia control.

Postemergence AC-222293 in sunflower, Farqo 1988. "Interstate' 3001 sunflower was seeded on May 18 to an area where wild oats had been seeded on the previous day. The wild oats did not emerge because of seed dormancy or dry soil conditions. Treatments were applied to 6- to 8-leaf sunflower, or dry soil conditions. Treatments were applied to 6- to 8-leaf sunflower, or dry soil conditions. Treatments were applied to 6- to 8-leaf sunflower, or dry soil conditions. Treatments were applied to 6- to 8-leaf sunflower, or dry soil conditions. The second application of split (/) 94 F, 35% RH, and 10 mph wind. The second application of split (/) 94 F, 35% RH, and 10 mph wind. The second application of split (/) 94 F, 35% RH, and 10 mph wind. The second application of split (/) 94 F, 35% RH, and 10 mph wind. The second application of split (/) 94 F, 35% RH, and 10 mph wind. The second application of split (/) 95 F, 35% RH, and 10 mph wind. The second application of split (/) 95 F, 35% RH, and 10 mph wind. The second application of split (/) 95 F, 35% RH, and 10 mph wind. The second application of split (/) 95 F, 35% RH, and 10 mph wind. The second application of split (/) 95 F, 35% RH, and 10 mph wind. The second application of split (/) 95 F, 35% RH, and 10 mph wind. The second application of split (/) 95 F, 35% RH, and 10 mph wind. The second application of split (/) 95 F, 35% RH, and 10 mph wind. The second application of split (/) 95 F, 35% RH, and 10 mph wind. The second application of split (/) 95 F, 35% RH, and 10 mph wind. The second application of split (/) 95 F, 35% RH, and 10 mph wind. The second application of split (/) 95 F, 35% RH, and 10 mph wind. The second application of split (/) 95 F, 35% RH, and 10 mph wind. The second application of split (/) 95 F, 35% RH, and 10 mph wind. The second application of split (/) 95 F, 35% RH, and 10 mph wind. The second application of split (/) 95 F, 35% RH, and 10 mph wind. The second application of split (/) 95 F, 35% RH, and 10 mph wind. The second application of split

	Rate	Sufl	Grft	Rrpw
reatment	(07/0)	<u>inj</u> (%)	(% cor	ntrol)
AC 222,293 AC 222,293+Sethoxydim+BCH AC 222,293+Sethoxydim+BCH AC 222,293+Sethoxydim+BCH AC 222,293+Sethoxydim+BCH AC 222,293+Sethoxydim+BCH+28% AC 222,293+Sethoxydim+BCH AC 222,293+Sethoxydim+BCH AC 222,293+Sethoxydim+BCH AC 222,293+Sethoxydim+BCH AC 222,293+Sethoxydim+BCH AC 222,293+Sethoxydim+BCH AC 222,293 Sethoxydim+BCH AC 222,293 Sethoxydim+BCH Sethoxydim+BCH Untreated  C.V. % LSD 5% LSD 1% # OF REPS	1.5 2+1.5+0.25G 2+3+0.25G 3+1.5+0.25G 3+2.4+0.25G 3+2.4+0.25G+1G 4+3+0.25G 6+2.4+0.25G 6/2.4+0.25G 6/2.4+0.25G 6+2.4+0.25G 1.5+0.25G 2.4+0.25G	15 16 32 20 7 37 42 13 62 40 51 8 15 3 0	13 78 87 78 50 84 85 84 80 75 73 22 13 80 91 0	53 53 63 79 58 84 78 52 65 77 62 34 38 0 0

#### Summary

Injury to sunflower was quite severe from AC 222,293 at 6 oz/A when applied in combination with sethoxydim and BCH (adjuvant from BASF-Dash). Green foxtail control was simuliar with sethoxydim applied seperately or with AC 222,293, except for treatment No. 5 which varied greatly among replications indicating a possible application error. Thus, AC 222,293 did not appear to antognize foxtail control with sethoxydim. Redroot pigweed control was too variable for interpretation of differences among the values.

AC 222,293 on weed free sunflower, Casselton 1988. 'Interstate 3001' sunflower was seeded on May 18. Weeds were removed by hand pulling within the row and by wheel hoe in between rows shortly after emergence. 10-leaves on June 16, 10- to 16-leaves on July 12, buds on July 22, and a shelterbelt and larger than in the other replication one were next to date. Refer to weather data in the first section of this report for were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at experiment was a randomized complete block with four replications. The Evaluation was on September 2. Estimates were made as to malformed heads birds.

Treatment	Rate	Height reduction	Malformed	Yield
AC 222,293(2 to 6 lf) AC 222,293(4 to 10 lf)	(oz/A) 6	(inch)	<u>heads</u> (%)-	<u>loss</u>
AC 222,293(4 to 10 lf) AC 222,293(10 to 16 lf) AC 222,293(bud to flowers) AC 222,293(late flowering)	6 6 6	4 6 1	59 15 56	56 12 59
C.V. %	6	2	0 0 0	1 4 0
LSD 5% LSD 1% # OF REPS		70 3 5	75 24 34	80 27
		4	4	37 4

## Summary

Injury from AC 222,293 did not relate to sunflower stage at treatment. The injurious June 6 application were when the day before, of, and after treatment had maximum temperatures of 96 to 100 F. The non-injurious June 16 treatments had a maximum temperatures of: 78 F on June 15, 79 F on June 16, and 90 F on June 17. The injurious July 12 treatments had maximum temperatures of: 81 F on July 11, 88 F on July 12, and 90 F on July 13. The at the vegetative stages may have contributed to the injury. Relative humidity was simuliar at all treatments, between 40 and 60%.

Weed control in soybean, Carrington 1988 Preplant incorporated (ppi) treatments applied and rototiller incorporated on June 1, "McCall' soybean planted on June 3 and preemergence treatments applied on June 7. A 0.34 inch rain occurred on June 12. Postemergence (P) treatment were to 1- to 2-trifoliolate soybeans, 4-leaf green foxtail and 8 inch tall redroot pigweed on June 24 and pl treatments were applied on June 24. Treatments were applied to an 8 ft wide area the length of 8 by 25 ft plots with a bicycle wheel type plot sprayer delivering 17 gpa for the ppi and pe and 8.5 gpa for the post emergence (p & pl) treatments, at 35 psi. The experiment was a randomized complete block with four replications. Evaluation was on July 15. Weed density was more than 2 plants/sq ft for all species.

	Rate	Soybean injury	Grft	Rrpw
reatment	(oz/A)	(%)	(% co	ntrol)
Trifluralin(ppi) Ethalfluralin(ppi) Pendimethalin(ppi) Trifluralin+Metribuzin-F(ppi) Pendimethalin+Imazethapyr(ppi) Trifluralin+Imazethapyr(ppi) Ethalfluralin+Imazethapyr(ppi) Imazethapyr(ppi) Alachlor(pe) Metolachlor(pe) Fenoxaprop+PO/Acifluorfen+Bent(p/pl) Sethoxydim+BCH/Acifluorfen+Bent(p/pl) Fluazifop+PO/Acifluorfen+Bent+BCH(p) Fluazifop+Acifluorfen+Bent+PO(p) Fenoxaprop+Acifluorfen+Bent+PO(p) Fenoxaprop+Acifluorfen+Bent+PO(p) Sethoxydim+BCH/Imazethapyr(p/pl+) Sethoxydim+BCH/Acifluorfen+X-77(p/pl) Sethoxydim+BCH/Bent+PO(p/pl) Imazethapyr(p) Imazethapyr(p) Imazethapyr+PO(p) Imazethapyr+MS(p) Untreated	12 12 20 12+3 14+1 12+1 12+1 1 40 40 2+0.25G/2+8 3+0.25G/2+8 3+0.25G/2+8 3+0.25G/2+8 3+2+8+0.25G 3+2+8+0.25G 2+2+8+0.25G 3+0.25G/1 3+0.25G/1 3+0.25G/1 3+0.25G/3+0.06G 1 1+0.25G 1+0.25G 0	0 0 0 0 0 0 0 0 0 0 1 1 1 0 4 1 1 0 4 0 0	99 98 98 97 98 99 90 40 55 80 95 68 81 59 43 87 94 66 82 78	86 80 0
C.V. % LSD 5%		267 2 NS	13 14 19	1 10
LSD 1% # OF REPS		4		1

Fenoxaprop, None of the herbicides caused important injury to soybean. sethoxydim, and fluazifop control of green foxtail was antagonized when applications were in tank mixture with bentazon + acifluorfen. from BASF, Dash) did not overcome bentazon + acifluorfen antogonism. Dinitroanalines, preplant incorporated, gave 97% or more green foxtail and redroot pigweed control. None of the postemergence or preemergence (pe) treatments gave adequate redroot pigweed control. However, postemergence treatments were to large plants. Imazethapyr postemergence with adjuvants gave 80% or more control of redroot pigweed

Preemergence weed control in soybean, Casselton 1988. Preplant soil incorporated (ppi) treatments were applied and field cultivator plus harrow (pe) treatments applied on May 5. The soil was moderately moist and mallow; Treatments were applied with a bicycle wheel type plot sprayer delivering 17 experiment was a randomized complete block with four replications. The Evaluations was on July 5. Weed densities were foxtail (green and yellow) 1 /sq ft and kochia more than 2/sq ft. Plants all severely drought stressed.

	Trailes all	severely dr	ought s	tressed.
Treatment  Trifluralin(ppi)	Rate (oz/A)	Soybean injury (%)	Grft	<u>KOCZ</u> ontrol)-
Ethalfluralin(ppi) Pendimethalin(ppi) Trifluralin + Metribuzin-DF(ppi Trifluralin + Imazethapyr(ppi) Pendimethalin + Imazethapyr(ppi) Ethalfluralin + Imazethapyr(ppi) Clomazone + Chloramben(ppi) Alachlor + Metribuzin-DF(ppi) Metolachlor + Metribuzin-DF(ppi) Metolachlor + Imazethapyr(ppi) Imazethapyr(ppi) Chloramben(pe) Imazethapyr(pe) Untreated  C.V. % LSD 5% LSD 1%	16 + 1 14 + 1 15 + 1 8 + 24		95 98 95 95 98 96 99 89 79 78 92 66 83 56 0	92 93 89 93 99 95 92 74 58 97 96 89 96 0
# OF REPS		NS NS 4	13 17 4	12 16 4

## Summary

None of the herbicides caused injury to soybean. All dinitroanaline herbicide treatments gave 95% or more foxtail control. Treatments containing imazethapyr all gave 95% or more kochia control.

Postemergence weed control in soybean, Casselton 1988. 'McCall' soybean was seeded on May 5. Treatments were applied to 1- to 2-trifolialate soybeans, and 5- to 6-leaf green and yellow foxtail on June 1 to 3 inch tall kochia, and 5- to 6-leaf green and yellow foxtail on June 16 with 60 F, 35% RH, and no wind at treatment. The second treatment of 16 with 60 F, 35% RH, and no wind at treatment. Treatments wind. Plant size had not changed from the June 16 treatment. Treatments wind. Plant size had not changed from the June 16 treatment. Treatments wind applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 were applied with a bicycle whee

Treatment	Rate (oz/A)	(% co	<u>Fxtl</u> ontrol)-
Sethoxydim+MSC/Acifluorfen+Bentazon Sethoxydim+Acifluorfen+Bentazon+MSC Sethoxydim+Acifluorfen+Bentazon+BCH Fluazifop+PO/Acifluorfen+Bentazon+PO Fluazifop+Acifluorfen+Bentazon+MSC Fluazifop+Acifluorfen+Bentazon+AG-2 Fenoxaprop+PO Quizalofop+PO Fenoxaprop+Bentazon+PO Quizalofop+Bentazon+PO Sethoxydim+Lactofen+PO Imazethapyr Imazethapyr Imazethapyr Imazethapyr Imazethapyr+MSC Untreated  C.V. % LSD 5% LSD 1% # OF REPS	3+0.25G/2+8 3+2+8+0.25G 3+2+8+0.25G 3+0.25G/2+8 3+2+8+0.25G 3+2+8+0.25G 3+2+8+0.25G 2.4+0.25G 1.2+0.25G 2.4+16+0.25G 1.2+16+0.25G 3+3+0.25G 0	8 71 73 10 60 82 70 0 90 86 46 9 98 0	87 79 85 46 44 60 55 74 77 72 60 89 42 75 0

## Summary

Kochia control with aciflurofen + bentazon, bentazon, and imazethapyr was enhanced by adjuvants. Kochia control was 98% with imazethapyr at 1 oz/A plus MS (Methylated seed oil adjuvant, Sun-it) compared to 9% control with imazethapyr alone. Green and yellow foxtail control was too variable with the drought to determine differences in antagonism of grass control herbicides or enhancement from various adjuvants.

Herbicide combinations postemergence in soybean, Casselton 1988. 'McCall' soybean was seeded on May 5. Treatments were applied to one to two trifolialate soybean, 5- to 6-leaf green and yellow foxtail, and 1 to 3 inch tall kochia on June 20 with 80 F, 40% RH, and 5 to 10 mph wind. The second mph wind. Plants were similar at the second split as first application. Plants were extremely drought stressed when treated. Treatments were applied area the length of 10 by 30 ft plots. The experiment was a randomized complete block with four replications. Evaluation was on July 12. Soybean gallons/A; BCH = adjuvants from BASF, Dash; MS = methylated seed oil, Sunit; product from Rhone Poulenc.

			iu, Ki
<u>Treatment</u>			
Bent+Acif+Seth+BCH+28%N Bent+Acif+PO/Seth+BCH+28%N Bent+Acif+PO/Bent+Seth+BCH+28%N Bent+Acif+BCH/Bent+Seth+BCH+28%N Bent+Acif+MS/Bent+Seth+MS+28%N Bent+Acif+PO/Bent+Acif+Seth+BCH+28%N Bent+Acif+PO/Bent+Acif+Seth+BCH+28%N Bent+Acif+PO/Acif+Seth+BCH+28%N Bent+Acif+PO/Acif+Seth+BCH+28%N Bentazon+Acifluorfen-RP+PO/Feno+P Bentazon+Acifluorfen-RP+Feno+PO Bentazon+Acifluorfen-RP+Fluazifop Bentazon+Acifluorfen-RP+Fluazifop Bentazon+Acifluorfen-RP+Fluazifop Bentazon+Acifluorfen-RP+PO/Flua+P Bentazon+PO/Sethoxydim+BCH Bentazon+Acifluorfen-RP+PO/Flua+P Bentazon+Bentazon+Bentazon Bentazon+Bentazon Bentazon+Bentazon Bentazon+Bentazon Bentazon+Bentazon Bentazon Benta	8+2+0.25G/8+2.4+0.25G+1G 8+2+0.25G/8+2.4+0.25G+1G 8+2+0.25G/8+2.4+0.25G+1G 8+2+0.25G/8+1+2.4+0.25G+1G 8+2+0.25G/1+2.4+0.25G+1G 8+2+0.25G/2+2.4+0.25G+1G 8+2+0.25G/2.4+0.25G 8+2+2.4+0.25G 8+2+2.4+0.25G 8+2+3+0.25G/3+0.25G	KOC (% c 64 19 69 91 75 83 78 60 64 16 16 5 20 45 3 28 34 10 5 53 73 73 73 0	Z Fxt1 ontrol 79 70 56 73 66 64 70 63 62 62 60 70 56 65 71 49 51 40 52 48 61 50 0 25 20 27 4

# Summary

Foxtail density was sparce and infestation variable making detailed comparison among treatment impossible. Kochia was not adequately controlled by any of the herbicide treatments.

Bentazon with adjuvants, Casselton 1988. 'McCall' soybean was seeded to moderately dry soil on May 6. Treatments were applied to 1 to 3 trifolialate soybeans, 1 to 4 inch tall kochia on June 17 with 78 F, 50% trifolialate sky, and no wind. Herbicide treatments were applied with a RH, clear sky, and no wind. Herbicide treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft wide area the length of 10 by 25 ft plots. The experiment was a randomized wide area the length of 10 by 25 ft plots. Evaluations were on July 1 and 28.

Combiece piece			
		July 1	July28
		Sobe KO	CZ KOCZ
	Rate	(% CO	ntrol)
Treatment	(oz/A)	22-(78 66	
	(02)	0 1	9 20
	8		3 18
Bentazon	16		31 64
Bentazon	8+0.12G		64 33 82
Rentazon+MSC	8+0.25G		
Bentazon+MSC	8+0.12G		
Bentazon+PO			
Bentazon+PO	8+0.25G		
Bentazon+AG-2	8+0.12G		85 76
Bentazon+AG-2	8+0.25G	0 3	48 48
Bentazon+X-77	8+0.25%	3	50 39
Bentazon+LI-700	8+0.25%	9	55 44
Bentazon+WK	8+0.25%	4 5	38 14
Bentazon+NaHCO3	8+16	5	77 64
Bentazon+NaHCO3+X-77	8+16+0.25%		
Bentazon-Manoocin		9	85 82
WEAN MEAN		0	13 14
HIGH MEAN		2	57 51
LOW MEAN		188	25 29
EXP MEAN		NS	21 21
C.V. %		NS	28 29
LSD 5%		4	4 4
LSD 1%			
# OF REPS			

# Summary

None of the treatments caused important injury to soybeans. Plants were extremely drought stressed at treatment. MSC (methylated seed oil from Agsco) and AG-2 in combination with bentazon at 8 oz/A gave greater kochia control then bentazon applied alone at 16 oz/A or at 8 oz/A with X-77, WK, PO, or LI-700. Sodium bicarbonate did not influence kochia control with bentazon, but increased control when with X-77 in the spray mixture.

Imazethapyr with adjuvants, Casselton 1988. 'McCall' soybean was seeded to moderately dry soil on May 6. Treatments were applied to 1 to 3 trifoliolate soybeans, 1 to 4 inch tall kochia on June 17 with 78 F, 50% RH, clear sky, and no wind. Herbicide treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft randomized complete block with four replications. Evaluations were on July 1 and 28.

Treatment	Rate	6.1	July		July 28
	(oz/A)	Sobe		Grft	KACZ
Imazethapyr Imazethapyr+X-77 Imazethapyr+MSC Imazethapyr+MSC Imazethapyr+PO Imazethapyr+PO Imazethapyr+AG-2 Imazethapyr+AG-2 Untreated Imazethapyr Imazethapyr Imazethapyr+MSC Imazethapyr+MSC Imazethapyr+PO Imazethapyr+PO Imazethapyr+PO Imazethapyr+AG-2 Imazethapyr+AG-2 Untreated HIGH MEAN	0.25 0.25+0.25% 0.25+0.12G 0.25+0.25G 0.25+0.12G 0.25+0.12G 0.25+0.25G 0 0.5 0.5+0.25% 0.5+0.12G 0.5+0.25G 0.5+0.25G 0.5+0.25G 0.5+0.25G 0.5+0.25G	0 1 0 0 1 0 0 3 0 1 5 0 0 1 0 1 0	19 87 93 98 88 92 95 95 3 33 87 97 98 93 93 97 98	15 53 71 86 73 78 70 75 0 36 45 62 65 55 77 84 88 0	1) 19 78 84 93 76 83 87 0 23 89 94 94 98 91 92 94 0
LOW MEAN EXP MEAN		5	98 0	88	94 0
C.V. % LSD 5% LSD 1% # OF REPS		266 NS NS	76 7 7 9	57 34 27 37	71 8 8

# Summary

None of the treatments caused important injury to soybeans. Green foxtail stand was sparse and variable so that differences among adjuvants with imazethapyr was not detectable, except that control was generally less when X-77 was the adjuvant. All adjuvants with imazethapyr enhanced kochia control with MSC (methylated seed oil from Agsco) and AG-2 causing the most enhancement.

Dash volume with sethoxydim plus bentazon, Prosper 1988. 'Wheaton' Hard Red Spring wheat, 'ND810104' oats and Siberian foxtail millet were seeded as bioassay species in adjacent 6 ft wide strips at Prosper, ND on May 23. Treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft wide area across the bioassay species. Plots were 10 ft wide to provide a 4 ft untreated area between treated areas to were 10 ft wide to provide a 4 ft untreated area between treated areas to aid evaluation. Treatments were applied to 5- to 6-leaf wheat, oats, and foxtail millet on June 27 with 95 F, 50% RH, and clear sky. The second foxtail millet on June 27 with 95 F, 50% RH, and partly cloudy sky. Evaluations were on July 8 and August 10.

Evaluations were on	July 8 and August 25			4		1	10
			July 8	3			10
		Fomi	Wht	Oats	Fomi	Wht	<u>Oats</u>
Treatment	Rate	10111		- (% co	ntrol)		
Treatment	(1b/A)						
Bent+COC/Seth+COC Bent+COC/Seth+Dash Bent+COC/Seth+MS Bent+Seth+COC Bent+Seth+Dash Bent+Seth+Dash Bent+Seth+Dash+28% Bent+Seth+Dash Bent+Seth+Dash Bent+Seth+MS Bent+Seth+MS Bent+Seth+MS Bent+Seth+MS C.V.% LSD 5% LSD 1%	1+0.25G/0.15+0.25G 1+0.25G/0.15+0.25G 1+0.25G/0.15+0.25G 1+0.15+0.25G 1+0.15+0.25G 1+0.15+0.25G 1+0.15+0.25G+1G 1+0.15+0.25G+1G 1+0.15+0.375G 1+0.15+0.375G	65 62 62 28 30 27 59 68 26 44 48 63 48 37		76 75 73 10 22 29 68 72 26 26 67 77 19 25	78 76 76 13 30 42 57 72 33 41 56 74 33 37	96 91 95 23 70 78 87 91 76 81 89 95 76 69	94 95 98 16 35 45 64 60 39 33 65 83 21 21
# OF REPS							

### Summary

Bentazon was antagonistic to grass control with sethoxydim regardless of adjuvant or volume of adjuvant. The inclusion of 28% N liquid fertilizer with Dash (adjuvant from BASF) or MS (Sun-it from Agsco) generally overcame the antagonism from bentazon.

Postemergence broadleaf weed control in dry bean, Casselton 1988. 'Upland' dry bean was seeded on May 5. Treatments were applied to first trifolialate dry bean, 3 to 8 inch tall kochia, and 2 to 6 inch tall wild mustard on June 21 with 95 F, 60 % RH, and 5 to 10 mph wind. Plants were extremelly drought as only a few were present. Treatments were applied for grass species control type plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft wide area the with four replication. Evaluation was on July 12. Injury to beans was not present or evaluated because the drought stress masked any injury.

<u>Treatment</u>		
Acifluorfen	Rate	KOCZ
Acifluorfen	(oz/A)	(control %
Acifluorfen	7	0
Acifluorfen	2 4	Ö
Acifluorfen+28%N	4	0
Acifluorfen+28%N	6	
Acifluorfen+28%N	1+1G	3 3 5 7
Acifluorfen+OC	2+1G	5
Acifluorfen+OC	1+0.25G	7
Sentazon+OC	2+0.25G	15
Sentazon+OC	8+0.25G	35
entazon+OC	12+0.25G	36
entazon+28%N	16+0.25G	
cifluorfen+Bentazon	12+1G	60
CITIU0rten+Rentazon	1+8	41
CITIUOTTEN+Rentazon. Do	1+12	19
- I I UUTTEN+Kentazon   DO	1+8+0.25G	18
- !! !UUrTen+Rentazon	1+12+0.25G	25
-ITIUOrten+Rentazon	2+8	43
ITIUOTten+Rentazon, 2000	2+12	15
III UUTTEN TRONT 3 7 0 m . DO	2+12+1G	13
111UUTTEN+Kentazon, DO	2+8+0.25G	55
11 140rten+Rentazon, DO /A . cz	2+12+0.25G	54
ifluorfen+Bentazon+PO/Acifluorfen+PO EP+MS	1+12+0.25G/1+0.25G	44
EP+MS EP+MS	1+12+0.25G/2+0.25G	41
treated	1+0.25G	54
	0	83
1. %	0	0
5%		
0 1%		40
F REPS		15
, ner s		20
		4

### Summary

None of the treatments gave adequate kochia control. The greatest kochia control was 83% with imazethapyr at 1 oz/A plus MS (methylated seed oil adjuvant, Sun-it).

Weed control with imazethapyr in dry bean, Casselton 1988. Preplant soil incorporated (ppi) treatments applied and twice field cultivator plus harrow incorporated 3 inches deep, 'Upland' dry bean seeded, and preemergence incorporated 3 inches deep, 'Upland' dry bean seeded, and preemergence treatments applied to a mallow moderately moist soil on May 5. to examine the second trifolialate beans, 1 to 3 inch Postemergence treatments (p) were to second trifolialate beans, 1 to 3 inch all kochia, and 5- to 6-leaf yellow foxtail on June 20 with 80 F, 40 % RH, tall kochia, and 5- to 6-leaf yellow foxtail on June 20 with 80 F, 40 % RH, tall kochia, and 5- to 6-leaf yellow foxtail on June 20 with 80 F, 40 % RH, tall kochia, and 5- to 6-leaf yellow foxtail on June 20 with 80 F, 40 % RH, tall kochia, and 5- to 6-leaf yellow foxtail on June 20 with 80 F, 40 % RH, tall kochia, and 5- to 6-leaf yellow foxtail on June 20 with 80 F, 40 % RH, tall kochia, and 5- to 6-leaf yellow foxtail on June 20 with 80 F, 40 % RH, tall kochia, and 5- to 6-leaf yellow foxtail on June 20 with 80 F, 40 % RH, tall kochia, and 5- to 6-leaf yellow foxtail on June 20 with 80 F, 40 % RH, tall kochia, and 5- to 6-leaf yellow foxtail on June 20 with 80 F, 40 % RH, tall kochia, and 5- to 6-leaf yellow foxtail on June 20 with 80 F, 40 % RH, tall kochia, and 5- to 6-leaf yellow foxtail on June 20 with 80 F, 40 % RH, tall kochia, and 5- to 6-leaf yellow foxtail on June 20 with 80 F, 40 % RH, tall kochia, and 5- to 6-leaf yellow foxtail on June 20 with 80 F, 40 % RH, tall kochia, and 5- to 6-leaf yellow foxtail on June 20 with 80 F, 40 % RH, tall kochia, and 5- to 6-leaf yellow foxtail on June 20 with 80 F, 40 % RH, tall kochia, and 5- to 6-leaf yellow foxtail on June 20 with 80 F, 40 % RH, tall kochia, and 5- to 6-leaf yellow foxtail on June 20 with 80 F, 40 % RH, tall kochia, and 5- to 6-leaf yellow foxtail on June 20 with 80 F, 40 % RH, tall kochia, and 5- to 6-leaf yellow foxtail on June 20 with 80 F, 40 % RH, tall kochia, and 5- to 6-leaf yellow foxta

ild mustard which was sparce and	Rate	Bean injury	Yeft	KOCZ	Wimu
<u> </u>	(oz/A)	(%)	(%	contr	01)-
EPTC+Chloramben(ppi) Pendimethalin(ppi) Imazethapyr(ppi) Imazethapyr(ppi) Imazethapyr+Pendimethalin(ppi) Imazethapyr+Pendimethalin(ppi) Trifluralin(ppi)/Bentazon(p) Pendimethalin(ppi)/Imazethapyr(p) Pendimethalin(ppi)/Bentazon(p) Imazethapyr(pe) Imazethapyr(pe) Imazethapyr(pe) Metolachlor+Imazethapyr(pe) Metolachlor(pe) Imazethapyr(p) Imazethapyr(p) Imazethapyr(p) Imazethapyr(p) Imazethapyr(p) Imazethapyr(p) Imazethapyr(p) Imazethapyr(p) Imazethapyr(p)	48+32 20 0.75 1 0.75+20 1+20 16/12 20/0.75 20/1 20/12 0.75 1 40+0.75 40+1 40 0.75 1 0.75+0.25G 0	0 1 0 1 4 0 0 0 0 5 0 0 6 4 0 8 11 5 14 0	97 98 49 70 98 99 94 98 96 29 23 72 71 70 35 31 63 0	98 63 84 93 68 99 81 86 68 73 73 67 64 83 0 18 21 93	80 25 65 58 28 94 97 90 82 45 53 75 9 83 90 98
Untreated  C.V. % LSD 5% LSD 1%		200 9 NS 4	19 19 25 4	28	27 36

## Summary

None of the herbicides caused important injury to dry beans. Yellow foxtail was controlled with herbicide treatments containing, EPTC, trifluralin, or pendimethalin. Imazethapyr alone did not give adequate (\$\psi^{75}) yellow foxtail control regardless of method of application. Kochia control was 90% or more with EPTC + chloramben, and imazethapyr at 1 oz/A alone or with pendimethylin preplant incorporated, and imazethapyr at 0.75 oz/A + MS pendimethylated seed oil adjuvant-Sun-it) postemergence. Wild mustard control (methylated seed oil adjuvant-Sun-it) postemergence. Wild mustard control (maxethapyr applied alone at 0.75 oz/A. MS with postemergence except imazethapyr applied alone at 0.75 oz/A. MS with postemergence imazethapyr enhanced or tended to enhance control of all weed species.

Weed control in corn, Casselton, 1988. An experiment was conducted to Bayaluate broad-spectrum weed control in corn. Preplant incorporated (PPI) treatments were applied and field cultivator Weed control in corn. Preplant incorporated (PPI) treatments were applied and field cultivator plus harrow incorporated twice in the opposite direction to a 3 inch depth on May 3, 1988. Inch depth on May 3, 1988. Inch depth on May 4. Soil was mellow with adequate soil moisture at seeding time. Early postemergence (EP) treatments were applied May 4. Soil was not a seeding time. mellow with adequate soil moisture at seeding time. Early postemergence (EP) treatments were applied to 1 to 3 leaf (2 to 6 inch) corn, 2 to 4 leaf green foxtail, and 1 to 3 inch kochia on June 1 with cloudy skies, 70 F, and 50% relative humidity. Postemergence (P) treatments were kochia on June 10 with sunny skies, 85 F, and 25% relative humidity. Each plot consisted of four rows of corn spaced 30 inches apart and 25 ft long. Treatments were applied with a bicycle wheel type plot sprayer delivering 17 gpa at 35 psi for soil applied treatments and 8.5 gpa at experimental design was a randomized complete block with four replications. Corn injury and green foxtail and kochia control were evaluated on June 20, with a second evaluation of kochia experimental design was a randomized complete block with four replications. Corn figury and green foxtail and kochia control were evaluated on June 20, with a second evaluation of kochia

<u>Treatment</u> <sup>a</sup>			June 20		July
	Rate	Corn	Green		<u>sury</u>
	(oz/A)	injury	foxtail	Kochia	Koch
EPTC&S+Cyanazine(PPI)	(/,//	(%)	(%	control	)
EPTC&S&E+Cyanazine(PPI)	64+32				
EPTC&S+Cyanazine+Atrazine(PPI)	64+32	0	81	74	84
Alachlor+Atrazine(PPI)	64+24+8	0	80	80	86
Alachlor+PPG-4000(PPI)	40+12	1	75	82	87
metolachior+Cvanazia-cos.	40+14.5	0	69	66	73
Alachlor+Cyanazine(PE)	40+32	0	42	72	85
Alachlor+Atrazine(PE)	40+32	1	45	25	53
Alachlor+PPG-4000(PE)	40+12	0	49	45	54
Propachlor+Cyanazine(PE)	40+14.5	1	80	66	64
Pendimethalia Com	64+32	0	60	53	46
Pendimethalin+Cyanazine(PE)	16+32	0	75	70	
Pendimethalin+Atrazine(PE)		1	51	59	68
endimethalin+PPG-4000(PE)	12+12	0	33	40	55
endimethal in+PPG-4000(EP)	24+14.5	0	43	55	61
ropachlor(PE)/Pyridate+Atrazine(EP)	24+14.5	0	28	15	55
	64/7+9.5	1	87		36
	64/7+9.5	5	92	95	96
yridate+Cyanazine(EP)	64/9.5+9.5	0	95	97	97
UDBCTLOT(PE\/Pomba-	14.5+20	11	78	98	98
opachlor(PE)/Bentazon+Atrazine+PO(EP) opachlor(PE)/DicambarAtrazine+PO(EP)	64/7+7+0.25G	Ö		94	92
ropachlor(PE)/Dicamba+Atrazine(EP)	64/8+8+0.25G	ő	89	98	98
	64/7+13	Ö	89	98	98
opachlor(PE)/Bromoxynil+Atrazine(EP) opachlor(PE)/Bromoxynil+Atrazine(EP)	64/3+6	4	86	98	98
	64/4+8	3	88	96	96
opachlor(PE)/Bromoxynil(EP)	64/4+4		90	98	98
Opachlor(PE)/Bromoxynil(EP)	64/4	11	86	98	97
Pacifical (PF )/Rromovim il . 2000.	64/6	3	87	95	94
opachlor(PE)/Clopyralid2,4-D-amine(P)	64/4+1G	7	85	94	91
ppachlor(PE)/2,4-D-ester(P)	64/1.5+8	2	81	93	95
ppachlor(PE)/Dicamba(P)	64/8	1	70	43	53
Phazine(EP)	64/4	1	72	91	90
nazine+SO(EP)	32	0	82	75	92
Dazino+Atana	19+0.25G	9	85	54	61
nazine+Atrazine+SO(EP)	15+5+0.25G	15		95	
. *	13+3+0.25G	9		95	92
5%				,,	95
		157	22	18	40
= dichlormid safener; E = dietholate ext ith 17% emulsifier; SO = soybean oil with		4		10	12

lormid safener; E = dietholate extender; & = formulated mixture; PO = petroleum oil with 17% emulsifier; SO = soybean oil with 15% emulsifier; G in the rate column represents

#### Summary

herbicide efficacy, and evaluation of the plots were adversely influenced by Plant growth, herbicide efficacy, and evaluation of the plots were adversely influenced by extremely hot, dry conditions and possible trifluralin carryover from the previous season. None of the treatments caused greater than 15% corn injury. Preemergence herbicide treatments provided less control than expected probably due to inadequate moisture. Preemergence propachlor plus postemergence pryridate plus atrazine tended to provide the highest green foxtail control. Several treatments that included postemergence herbicides provided 90% or greater kochia control at both evaluation dates. Cyanazine at 19 oz/A plus vegetable oil gave more injury, but provided better weed control than cyanazine alone at 32 oz/A. 2.4-D ester gave more injury, but provided better weed control than cyanazine alone at 32 oz/A. 2,4-D ester gave more than 35% greater kochia control than the clopyralid plus 2,4-D amine prepackage mixture at

Weed control in corn, Carrington, 1988. An experiment was conducted to broad-spectrum weed control in corn with various herbicide treatments. Preplant incorporated (PPI) treatments were applied and roto-tiller incorporated, and 'Top Farm 1185' corn was seeded on June 5. Preemergence (PE) treatments were applied June 8. PPI and PE treatments were applied to dry soil with air temperatures of 80 F and above. A 0.50 inch rain occurred within 12 hours after the preemergence treatments. Postemergence (P) treatments were applied to 4 to 5 leaf corn, 2.5 leaf green foxtail, and 4 to 6 leaf redroot pigweed on June 20. Temperatures exceeded 90 F the day of and the day following postemergence treatments. Each plot consisted of three rows of corn spaced 30 inches apart and 25 ft long. Treatments were applied with a bicycle wheel type plot sprayer delivering 17 gpa at 35 psi for soil applied treatments and 8.5 gpa at 35 psi for postemergence treatments. The experimental design was a randomized complete block with four replications. Corn injury and weed control were evaluated on July 15.

orn injury and weed control were evaluate		Corn	Green foxtail	Redroot
a		injury	(% COI	ntrol)
<u>reatment<sup>a</sup></u>	(oz/A)	(%)	(/// 001	1010.7
	-1.00	0	97	97
ODI)	64+32	0	97	98
PTC&S+Cyanazine(PPI)	64+32	1	73	79
utylate&S+Cyanazine(PPI)	40+32	0	91	96
etolachlor+Cyanazine(PE)	40+32	0	90	87
lachlor+Cyanazine(PE)	64+32	1	66	50
ropachlor+Cyanazine(PE)	16+32		54	40
endimethalin+Cyanazine(PE)	12+12	6	93	96
	64/7+9.5	0	87	96
Propachlor(PE)/Pyridate+Atrazine(P)	64/9.5+9.5		48	75
Dyonach Inr(PE)/Pyridate inti az inti	14.5+20	25	85	95
	64/8+8+0.25G	0	81	97
Propachlor(PE)/Bentazon&Atrazine+PO(P)	64/7+13	0	83	85
	64/3+6	1	81	92
Description of the I / Brolliox villiance and the	64/6	1	83	86
	64/1.5+8	0	92	91
- Liewide // Innvidium	64/8	0	85	91
Duanach   Or ( PF ) / 2, 4-U-E3 ( C) ( ' )	64/4	0	46	16
Propachior(PE)/Dicamba(1)	32	8	55	56
Cyanazine(P)	19+0.25G	19	36	63
Cuanazino+(IIIP)	15+5+0.25G	16	84	90
Cyanazine&Atrazine+SO(P)	0.5+0.25G	23	89	97
NPX-V9360+PU(P)	1.0+0.25G	16	43	69
DPX-V9360+PO(P)	0.33+0.25%	10		
CGA-136872+X-77(P)	0.5+0.25%	23		A CONTRACTOR OF THE PARTY OF TH
CCA = 136872 + X - I/(P)	8+0.12+0.259	% 16	40	
Atrazine+DPX-M6316+SO(P)			17	1
		96		
C.V. %		9	1/	
LSD 5%  a S = dichlormid safener; & = formulate	d mixture; PO	= petr	roleum oi	I WICH

S = dichlormid safener; & = formulated mixture; emulsifier; SO = soybean oil with 15% emulsifier.

#### Summary

Treatments containing postemergence cyanazine, DPX-V9360, CGA-136872, and DPX-M6316 caused corn injury. However, corn injury from postemergence herbicides was difficult to differentiate from pretreatment weed competition. EPTC&S and butylate&S tended to provide better green foxtail control than the other treatments. DPX-V9360 gave 84 to 89% green foxtail control and CGA-136872 gave 43 to 49% green foxtail control at the rates evaluated. herbicide treatments gave good redroot pigweed control.

<u>Postemergence weed control in corn, Casselton, 1988.</u> An experime conducted to evaluate grass and broadleaf weed control with several applied herbicides in 'Interstate 201' corn seeded May 4. Early postemergence (EP) treatments were applied to 1 to 3 (2 to 6 inch) leaf corn, 2 to 4 leaf green foxtail, and 1 to 3 inch kochia on June 1 with cloudy skies, 40 F, and 50% relative humidity. Postemergence (P) treatments were applied to 3 to 5 leaf (6 to 12 inch) corn, 5 leaf to tillering green foxtail, and 2 to 6 inch kochia on June 10 with clear skies, 85 F, and 25% relative humidity. Each plot consisted of four rows of corn spaced 30 inches apart and 25 ft long. Treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi to the middle two rows of the four row plots. Corn and weeds were exhibiting drought symptoms at both dates of application. was a randomized complete block with four replications. control were evaluated June 20, July 1, and July 25. The experiment Corn injury and weed

June 20 July 1 <u>Treatment</u><sup>a</sup> July Corn Rate injury Grft KOCZ Grft KOCZ (oz/A)KOCZ ----(% control)--(%) DPX-V9360+P0(EP) 0.25+0.25G DPX-V9360+P0(EP) 0.5+0.25G DPX-V9360+P0(EP) DPX-V9360+P0(EP) 0.75+0.25G DPX-V9360+P0(P) 1.0+0.25G 0.25+0.25G DPX-V9360+P0(P) 0.5+0.25G DPX-V9360+P0(P) 0.75+0.25G DPX-V9360+P0(P) 1.0+0.25G CGA-136872+X-77(EP) CGA-136872+X-77(EP) CGA-136872+X-77(P) 0.33+0.25% 0.5+0.25% CGA-136872+X-77(P) 0.33+0.25% 0.5+0.25% Cyanazine+DPX-M6316+SO(EP) 24+0.25+0.25G Atrazine+DPX-M6316+SO(EP) 8+0.12+0.25G Cyanazine(EP) Cyanazine+SO(EP) 19+0.25G Cyanazine&Atrazine+SO(EP) 15+5+0.25G C.V. % LSD 5% 

PO = petroleum oil with 17% emulsifier, X-77 = nonionic surfactant; soybean oil with 15% emulsifier; G in the rate column represents gallons/acre; & = formulated mixture.

#### Summary

Plant growth and herbicide efficacy were influenced by extremely conditions and possible trifluralin carryover from the previous season. Treatments containing cyanazine and soybean oil caused 14 to 16% injury. None of the other treatments caused important injury. DPX-V9360 provided better green foxtail control than CGA-136872 at the rates evaluated. control was less with the second evaluation than the first evaluation with the triazine herbicides, but was similar for both evaluation dates with DPX-V9360 DPX-V9360 and CGA-136872 provided greater than 70% kochia control with all application rates and times at the late evaluations. M6316 plus atrazine or cyanazine gave 97% or better kochia control. Cyanazine at 19 oz/A plus soybean oil provide 40% greater kochia control than cyanazine

An experiment was conducted Weed control in grain sorghum, Carrington, 1988. to evaluate broad-spectrum weed control in grain sorghum with soil applied and postemergence treatments. Preplant incorporated (PPI) treatments were applied and roto-tiller incorporated prior to seeding of flurazole safener treated Dekalb 'DK 18' grain sorghum on June 7. Preemergence (PE) treatments were applied June 13. Precipitation totaling 0.34 inch was received between the preplant incorporated and preemergence treatments. Precipitation for 2 weeks after preemergence treatments was 1.29 inch. Postemergence treatments were applied to 3 to 4 leaf grain sorghum and seedling green foxtail and redroot pigweed on July 5. The maximum temperature on July 5 was 100 F and 0.30 inch of rain was received approximately 8 hours after the postemergence treatments All treatments were applied with a bicycle wheel type plot sprayer delivering 17 gpa at 35 psi for preemergence treatments and 8.5 gpa at 35 psi for postemergence treatments. The experiment was a randomized complete block design with four replications. Grain sorghum injury and weed control were evaluated July 15.

. a	Rate	Grain sorghum injury	Green foxtail	Redroot pigweed
Treatment <sup>a</sup>	(oz/A)	(%)	(% CC	ontrol)
Atrazine+Alachlor(PPI) Propazine+Alachlor(PPI) Cyanazine+Alachlor(PPI) Cyanazine+Metolachlor(PPI) Cyanazine&Atrazine(PPI) Atrazine+Alachlor(PE) Cyanazine+Alachlor(PE) Propachlor+Cyanazine(PE) BAS-51400(PE) BAS-51400+MS(P) Propachlor(PE)/2,4-D(P) Propachlor(PE)/Bromoxynil(P) Propachlor(PE)/Bromoxynil&Atrazine(P) Propachlor(PE)/Bentazon&Atrazine(P) C.V. % LSD 5%	12+32 12+32 32+32 32+32 32+12 12+32 32+32 48+24 16 8+0.25G 48/8 48/4 48/6 48/4+8 48/8+8	0 0 0 0 0 0 0 0 0 0 0 8 5 1	78 70 91 87 94 85 92 82 95 80 89 88 80 84 86	98 95 98 98 98 94 96 91 60 10 91 94 92 99 97

 $<sup>^{\</sup>rm a}$  & = formulated mixture; MS = methylated vegetable oil with 15% emulsifier.

## Summary

None of the treatments caused important grain sorghum injury. All treatments provided 70% or greater green foxtail control. Preemergence BAS-51400 gave 95% green foxtail control. All treatments except BAS-51400 provided greater than 90% redroot pigweed control. Preemergence BAS-51400 at 16 oz/A gave better weed control than postemergence BAS-51400 at 8 oz/A plus methylated vegetable oil.

Postemergence weed control in canola, Langdon 1988. 'Westor' canola was seeded on May 16. Treatments were applied to 4- to 5-leaf canola, emerging to 8-leaf ladysthumb, 4 inch wild mustard, and 1- to 2-leaf wild buckwheat on Treatments were applied with a bicycle wheel type plot sprayer delivering 7.8 gpa at 35 psi to 8 by 23 ft plots. The experiment was a randomized complete block with four replications. Evaluation was on July 20.

							415 20.
Treatment	Rate	Canola	Vol.	Ladys-	Test		Canola
DPX-A7881-22(2-41f) DPX-A7881-22 DPX-A7881-22 DPX-A7881-22 DPX-A7881-22 DPX-A7881-22+X-77 DPX-A7881-22+MS	0.17 0.33 0.50 0.67 0.17+0.2 0.33+0.2 0.17+0.2	11 0 13 3 25% 18 25% 5 4	wheat	thumb Yiel htrol)(lb/A 45 366.1 50 720.1 71 276.1 76 476.0 39 669.0 45 590.5	d weight (bu/lb) 5 49.0 5 50.0 6 49.0 0 50.0 0 29.0 5 51.0	Canola (%) 97 90 88 93 96 97	Canola yield (1b/A) 354.2 648.5 243.0 444.4 644.6 574.1
DPX-A7881-22+MS DPX-A7881-22+Seth+MS DPX-A7881-22+Seth+MS DPX-A7881-22+Flua+MS DPX-A7881-22+Flua+MS Sethoxydim+MS Fluazifop+MS AC222293 Benazolin IMEP	0.33+0.2 0.17+3+0. 0.33+3+0.	25G 6 25 30 25 40 25 16 25 18 0 8 54	0 97 97 99 99 99 13 0	72 513.0 70 410.0 50 514.5 62 118.0 59 446.0 68 489.5 21 529.5 5 601.0 34 57.5 49 692.0	49.5	99 94 95 90 95 98 94 94	505.4 386.0 488.8 106.2 423.7 477.8 502.0 578.0 31.8
Clopyralid Clopyralid BAS-514+MSF Untreated	1.5 3 8+0.25G 0	78 0 8 60 0	0 0 0 5 0	46 33.0 23 752.0 30 278.5 30 195.0 9 478.0	.0 48.0 47.5 .0 49.0	15 80 95 73	656.4 5.1 603.1 264.6 140.0 447.9
LOW MEAN EXP MEAN C.V. % LSD 5% LSD 1% # OF REPS		0 18 2 69 2 17 23 1	0	23 400.4	15.9 12.9	15 87 4 5	556.4 5.1 06.0 47.4 01.5 NS 2

## Summary

Canola had adequate tolerance with DPX-A7881, except when applied with methylated sunflower oil adjuvant (MS) in combination with sethoxydim. DPX-A7881 injury to canola was less than 19% when applied alone, with adjuvants, or adjuvant plus fluazifop. Clopyralid and benasolin did not injure canola. DPX-A7881 control of ladysthumb was enhanced by the MS adjuvant, but not by Yields were taken from only two replications and were variable. However, treatments which caused severe injury resulted in low canola yield.

Preemergence weed control in Canola, Langdon 1988. Preplant (PPI) treatments applied and rototiller incorporated 3 inch deep, 'Westor' canola seeded at 7 lb/A + 0.5 lb ai/A Furadan CRIO, and preemergence incorporated (PEI) treatment applied and harrow incorporated twice on May 16. A 0.15 inch rain occurred on applied and harrow incorporated twice on May 16. A 0.15 inch rain occurred on Treatments were applied with a bicycle wheel type plot sprayer May 19. Treatments were applied with a bicycle wheel type plot sprayer delivering 17 gpa at 35 psi to 8 by 25 ft plots. The experiment was a delivering 17 gpa at 35 psi to 8 by 25 ft plots. Evaluations were on July randomized complete block with four replications. Evaluations were on July 20. Harvest was from 64 sq ft area.

20. Halveso has								
	Rate (oz/A)	St.Red.	Inj (%)(%	Ladys- thumb % control	Total yield )(lb/A)		Canola (%)	Canola yield (lb/A)
Trifluralin PPI Ethalfluralin PP Pendimethalin PP Trifluralin PPI Ethalfluralin PP Pendimethalin PP Trifluralin PEI Ethalfluralin PE Pendimethalin PE Pendimethalin PE	12 I 13 I 20 I 16 I 18 I 24 I 12 II 13 II 20 II 20	0 16 5 5 13 5 0 5 3 0	0 6 3 4 8 4 0 9 1 0 8 0	83 89 77 87 86 83 66 68 28 19 29	745 630 876 863 850 758 1036 1020 1053 1034 702 881	48.9 48.8 48.6 49.3 47.9 49.4 47.1 48.4 46.5 44.5 47.1 43.6	84 86 85 88 86 76 79 70 68 79	630 542 747 747 742 645 799 819 720 706 549 447
Untreated HIGH MEAN LOW MEAN EXP MEAN C.V. % LSD 5% LSD 1% # OF REPS	Ü	16 0 4 154 10 NS 4	9 0 3 148 NS NS NS	89 0 59 20 18 24 4	1053 630 871 27 NS NS	49.4 43.6 47.5 3.9 2.7 3.6	12 14 19 4	819 447 674 NS NS 4
<u> </u>				1	o/ ca	nola is	a visua	l estima

Total yield is canola plus ladysthumb seed, % canola is a visual estimate based on hand separated samples of the total yield, and canola yield is the yield corrected for ladysthumb using the % canola values.

## Summary

Ladysthumb was greater with preplant incorporated herbicides than preemergence incorporated on surface applied. The higher rates of the dinitroanalines for acceptable ladysthumb control, but control was still less than 90%. The amount of ladysthumb control related to the percent canola in the harvested sample.

Weed control in safflower, Williston 1988. treatments were applied and twice field cultivator incorporated and 'Finch' Preplant soil incorporated safflower planted on May 19 with 55 F, 30% RH, cloudy sky, a 3 mph wind, and dry soil. Postemergence treatments were to 2- to 12-leaf stage safflower, 2to 4-leaf green foxtail (some tillering, 6 inches tall), 4 to 6 inch tall wild mustard and Russian thistle, and 6 to 7 inch tall redroot pigweed on June 22. Treatments were applied with a tractor-mounted plot sprayer delivering 17 gpa to an 8 ft width the length of 10 by 25 ft plots. The experiment was a randomized complete block design with four replications. Evaluation was on July 20. Injury ratings or yield were not taken because of a variable wild mustard and redroot pigweed moderate, and green foxtail sparce. Activator 90 and PO = petroleum oil.

Treatment			Rate	0.0				
			(OZ/A)	Grf	1140		u Rrpv	v Colq
Trifl			(oz/A)		(		tro1)	corq
Trifluralin		PPI	10				,	
Trifluralin		PPĪ	12	90	48	23	84	97
Ethafluralin		PPI	16	96	78	13	91	
Trifluralin+EPTC		PI	20	96	76	24	95	99
Pendimethalin		PI	12+48	98	69	86	95	99
RE 40885		PI	16	88	18	30	85	99
RE 40885		PI	6	73	50	49		95
RE 40885		PI	.8	76	76	47	73	97
RE 40885		_	12	85	79	69	71	97
RE 40885+Trifluralin		PI	16	89	88	61	79	97
RE 40885+Sethovydim. Do		PI	12+8	79	65		89	97
VE TOOOD+SETHOYVOIM DO		PI+P	12+3+0.25G	89	85	44	83	97
"		PI+P	8+3+0.25G	87.	78	81	79	95
NE 40000+Sethovydim, no	PI	I	8+.0.25G	85	58	55	81	95
UI ATINOS I DEACT I VOE ON ON	P		4+3+0.25G	94	19	70	95	97
UFA-1103 10+50thoxyd:	P		0.25+0.25%V	6		89	71	97
DPX-M6316+Sethoxydim+PO	P		0.125+3+0.25G	93	93	70	96	99
DPX-M6316+Activator 90	P		0.167+3+0.25G	89	95	80	98	99
Chlorosulfuron+Activator 90	P		0.167+0.25%	0	88	91	98	97
AC 222,293	Р		0.167+0.25%		96	77	98	99
Chloroculture	Р		6	16	97	99	99	99
Chlorosulfuron+Sethoxydim+PO Sethoxydim+PO	P		0.167+3+0.25G	14	45	89	20	38
Fluazifop+PO	Р		4+0.25G	68	82	73	92	97
Untreated	P		4+0.125G	93	0	0	0	0
oucreated			0	84	0	0	0	Ö
C.V. %			U	0	0	0	0	0
C.V. % LSD 5%								U
				13	30	55	15	13
				13	26	42	16	23
# OF REPS				17	24	56	21	31
				4	4	4	4	
			Summan					2

#### Summary

Green foxtail control with sethoxydim was reduced by chlorsulfuron in the spray, but not by DPX-M6316. The greatest Russian thistle and wild mustard control was with the sulfonylurea herbicides. RE-40885 at 8 oz/A or more gave 75% or more Russian thistle control. Russian thistle control was increased from 47 to 77% when trifluralin rate was increased from 12 to 16 oz/A. Common lambsquarter was adequately controlled by all herbicides except AC 222,293, sethoxydim, and fluazifop. Redroot pigweed was adequately controlled by the

Annual forage crops response to herbicides, Minot 1988. PPI treatments were applied to moist soil and cultiweed incorporated twice, forage crops were seeded and PE treatments applied to the surface June 16. Preemergence treatments were applied in 17 gpa at 35 psi with sunny skies and 70 F. treatments were applied in 8.5 gpa at 35 psi to 4.3 to 4.8 leaf Postemergence treatments were applied in 8.5 gpa at 35 psi to 4.3 to 4.8 leaf Postemergence treatments were applied in 8.5 gpa at 35 psi to 4.3 to 4.8 leaf Postemergence treatments were applied in 8.5 gpa at 35 psi to 4.3 to 4.8 leaf Postemergence treatments were applied in 8.5 gpa at 35 psi to 4.3 to 4.8 leaf Postemergence treatments were applied in 8.5 gpa at 35 psi to 4.3 to 4.8 leaf Postemergence treatments were applied in 8.5 gpa at 35 psi to 4.3 to 4.8 leaf Postemergence treatments were applied in 8.5 gpa at 35 psi to 4.3 to 4.8 leaf Postemergence treatments were applied in 8.5 gpa at 35 psi to 4.3 to 4.8 leaf Postemergence treatments were applied in 8.5 gpa at 35 psi to 4.3 to 4.8 leaf Postemergence treatments were applied in 8.5 gpa at 35 psi to 4.3 to 4.8 leaf Postemergence treatments were applied in 8.5 gpa at 35 psi to 4.3 to 4.8 leaf Postemergence treatments were applied in 8.5 gpa at 35 psi to 4.3 to 4.8 leaf Postemergence treatments were applied in 8.5 gpa at 35 psi to 4.3 to 4.8 leaf Postemergence treatments were applied in 8.5 gpa at 35 psi to 4.3 to 4.8 leaf Postemergence treatments were applied in 8.5 gpa at 35 psi to 4.3 to 4.8 leaf Postemergence treatments were applied in 8.5 gpa at 35 psi to 4.3 to 4.8 leaf Postemergence treatments were applied in 8.5 gpa at 35 psi to 4.3 to 4.8 leaf Postemergence treatments were applied in 8.5 gpa at 35 psi to 4.3 to 4.8 leaf Postemergence treatments were applied in 8.5 gpa at 35 psi to 4.3 to 4.8 leaf Postemergence treatments were applied in 8.5 gpa at 35 psi to 4.3 to 4.8 leaf Postemergence treatments were applied in 8.5 gpa at 35 psi to 4.3 to 4.8 leaf Postemergence treatments were applied in 8.5 gpa at 3

were evaluated bec	ause or						Fami	Suav	Hygr	Foso	Suso
Hambigido	Rate		Trit	Spel	Pemi	Prmi	Fomi inju				
Herbicide	(oz/A)	(%)				(76	1113 4	, , ,			25
Triallate PPI Trifluralin PPI Cyanazine-L PPI Propachlor PE Dicamba P 2,4-D dma P MCPA dma P Imazamethabenz P Diclofop P Brox&MCPA P Bromoxynil P DPX-M6316+R11 P	16 12 16 48 4 8 8 6 12 8 4 0.375	0 90 28 5 92 85 30 8 0 60 30	10 85 63 23 10 3 0 0 0	10 75 15 8 13 3 5 3 0 3	0 99 3 48 8 3 5 45 5 10 35	0 99 18 93 10 3 5 0 20 0 3 50	0 99 20 90 25 18 28 7 63 10 3	35 99 3 13 15 3 18 13 13 15 85	83	55 97 0 0 5 0 8 25 53 5 3 83	99 0 0 5 0 15 25 63 8 5
C.V. % LSD 5% # OF REPS F-TRT		25 24 2 25	18	16 2 16	16 2 3 3	13	10 2 34	27	27	22 2 3 24	20

a Rrpw = Redroot pigweed control; Trit = Triticale; Spel = Speltz; Pemi = Pearl millet; Prmi = Proso millet; Fomi = Foxtail millet; Sugr = Sudan grass; Hysd = Hybrid sudan; Foso = Forage sorghum; Suso = Sudan sorghum; b R-11 = Surfactant applied at 0.25% v/v.

#### Summary

Trifluralin, dicamba, 2,4-D, and DPX+M6316 controlled redroot pigweed 85% or greater. Trifluralin severly reduced stands of all crops tested. Cyaniazine reduced the stand of triticale 63%. Propachlor reduced the stand of Mil-Hy reduced the stand of milets, and Manta proso millet. Diclofop and DPX-100 pearl, Cerise proso millets, and Manta proso millet. Diclofop was most injurious M6316 were injurious to pearl and proso millets. Diclofop was most injurious to proso millet, hybrid sudan grass, forage sorghum and the sudan X sorghum to proso millet, hybrid sudan grass, forage sorghum to proso millet, hybrid sudan grass, forage sorghum and the sudan X sorghum cross. DPX-M6316 injured all the warm season crops tested.

Commercial petroleum oils with sethoxydim, Exp 1, Fargo 1988. 'Wheaton' Hard Red Spring wheat and 'ND810104' oats were seeded as bioassay species in Hard Red Spring wheat and 'ND810104' oats were seeded as bioassay species in adjacent 10 ft wide strips at Fargo, ND on April 28. Treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi provide a 4 ft untreated area between treated areas to aid evaluations. Treatments were applied to 4- to 5-leaf wheat and oats on June 2.

Treatment					
Treatment  Sethoxydim+PO1(Atplus411F) Sethoxydim+PO2(Herbimax) Sethoxydim+PO3(Coop) Sethoxydim+PO4(Wilbur) Sethoxydim+PO5(Add-It) Sethoxydim+PO6(Dash) Sethoxydim+PO7(Assist) Sethoxydim+PO8(Zach)  HIGH MEAN LOW MEAN EXP MEAN	Rate (oz/A) 1+0.25G 1+0.25G 1+0.25G 1+0.25G 1+0.25G 1+0.25G 1+0.25G 1+0.25G	38 33 23 48 35 29 38 30	40 31 24 41 52 34 53 36	33 43 40 53 18 28 23 46	1ne 24 20ats ) 41 38 34 56 31 24 30 51
EXP MEAN C.V. % LSD 5% LSD 1% # OF REPS		23 34 32 NS NS	24 39 44 NS NS	18 35 52 NS NS	24 38 50 NS NS

## Summary

All petroleum oils were similar as adjuvants with sethoxydim. The control obtained with the dry environment was too little to detect differences.

Commercial petroleum oils with sethoxydim, Exp 2, Farqo 1988. 'Wheaton' Hard Red Spring wheat 'ND810104' oats, and Siberian foxtail millet were seeded as bioassay species in adjacent 6 ft wide strips at Fargo, ND on May seeded as bioassay species were applied with a bicycle wheel type plot sprayer 31. Treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft wide area across the bioassay delivering 8.5 gpa at 35 psi to an 8 ft wide area across the bioassay species. Plots were 10 ft wide to provide a 4 ft untreated area between species. Plots were 10 ft wide to provide a 4 ft untreated area between treated areas to aid evaluation. Treatments were applied to 5- to 6-leaf treated areas to aid evaluation. Treatments were applied to 5- to 6-leaf wheat, oats, and foxtail millet on July 6. Evaluation was on August 1 and 24.

24.			ugust Wht	1 Oats	Au Fomi	gust Wht	24 0ats
Treatment	Rate (oz/A)	Fomi		-(% co	ntrol)		
Sethoxydim+PO1(Atplus411F) Sethoxydim+PO2(Herbimax) Sethoxydim+PO3(Coop) Sethoxydim+PO4(Wilbur) Sethoxydim+PO5(Add-It) Sethoxydim+PO6(Dash) Sethoxydim+PO7(Assist)	1+0.25G 1+0.25G 1+0.25G 1+0.25G 1+0.25G 1+0.25G 1+0.25G	75 77 75 78 73 79 73 76	92 94 95 94 91 96 92 97	84 88 86 89 79 87 78 88	80 83 80 81 76 86 77 83	99 98 99 99 99 99	93 98 96 99 90 98 89 98
Sethoxydim+PO8(Zach)  C.V. % LSD 5% # OF REPS	110.200	5 NS 4	3 NS 4	5 6 4	5 NS 4	1 NS 4	2 3 4

## Summary

All petroleum oils were similar with sethoxydim for control of grass species. Oats control tended to be less with assist and Add-it than other oils.

Commercial petroleum oils with fluazifop, Prosper 1988. 'Wheaton' Hard Red Spring wheat, 'ND810104' oats, and siberian foxtail millet were seeded as bioassay species in adjacent 6 ft wide strips at Prosper, ND on May 23. Treatments were applied with a bicycle wheel type plot sprayer delivering were 10 ft wide to provide a 4 ft untreaten area between treated areas to aid evaluation. Treatments were applied to 5- to 6-leaf wheat and oats and August 5.

Treatment	Rate	Fomi	July			Augus	t 5
Flua+PO1(Atplus411F)	(oz/A)		<u>Wht</u>	Oats -(% cor	Fxm trol	i Wht )	Oats
Flua+PO2(Herbimax) Flua+PO3(Coop) Flua+PO4(Wilbur) Flua+PO5(Add-It) Flua+PO6(Dash) Flua+PO7(Assist) Flua+PO8(Zach)  HIGH MEAN	0.75+0.25G 0.75+0.25G 0.75+0.25G 0.75+0.25G 0.75+0.25G 0.75+0.25G 0.75+0.25G 0.75+0.25G	23 22 17 26 12 27 25 20	78 79 76 76 77 77 78 77	72 72 72 75 75 76 71	28 25 21 28 11 25 24 25	96 97 96 96 96 93 98	90 95 86 90 91 93 89
LOW MEAN EXP MEAN C.V. % LSD 5% LSD 1% # OF REPS		27 12 21 43 NS NS 4	79 76 77 4 NS NS 4	76 71 74 8 NS NS	28 11 23 19 6 9	98 93 96 2 NS NS	96 86 91 5 NS NS

Summary

All petroleum oil were similar as adjuvants with fluazifop.

Commercial petroleum oils with DPX-Y6202, Fargo 1988. 'Wheaton' Hard Red Spring wheat, 'ND810104' oats, and Siberian foxtail millet were seeded as bioassay species in adjacent 6 ft wide strips at Fargo, ND on May 31. Treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft wide area across the bioassay species. Plots were 10 ft wide to provide a 4 ft untreated area between treated areas to were 10 ft wide to provide a 4 ft untreated area between treated and oats and aid evaluation. Treatments were applied to 5- to 6-leaf wheat and oats and aid evaluation. Treatments were applied to 5- to 6-leaf wheat and oats and jointing foxtail millet on July 6 with 82 F, and 50% RH, 1.5h after a light rain. Evaluations were on August 1 and 24.

rain. Evaluations were on	August 1 une	Δ	uqust	1	Au		24
	Date	Fomi	Wht	Oats	Fomi		<u>Oats</u>
Treatment	Rate (oz/A)			-(% CO	ntrol)		
DPX-Y6202+P01(Atplus411F) DPX-Y6202+P02(Herbimax) DPX-Y6202+P03(Coop) DPX-Y6202+P04(Wilbur) DPX-Y6202+P05(Add-It) DPX-Y6202+P06(Dash) DPX-Y6202+P07(Assist) DPX-Y6202+P08(Zach) C.V. %	0.4+0.25G 0.4+0.25G 0.4+0.25G 0.4+0.25G 0.4+0.25G 0.4+0.25G 0.4+0.25G 0.4+0.25G	67 71 72 69 71 74 65 67	99 99 99 99 99 98 99 0 NS	68 78 72 77 76 81 64 72	35 37 36 43 43 59 36 38	97 99 99 99 99 99 99 89 99	55 77 76 81 78 95 65 79 15 17
LSD 5% # OF REPS		4	4	4			

## Summary

Assist and Add-it tended to be less effective than the other adjuvants for oats control with DPX-Y6202.

Sethoxydim formulations with adjuvants, Exp 1, Prosper 1988. 'Wheaton' Hard Red Spring wheat, 'ND810104' oats, and Siberian foxtail millet were seeded as bioassay species in adjacent 6 ft wide strips at Prosper, ND on May 23. Treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 and 35 points are a species the bioassay species. Plats were 10 gpa at 35 psi to an 8 ft wide area across the bioassay species. Plots were 10 ft wide to provide a 4 ft untreated area between treated areas to aid evaluation. Treatments were applied to 4- to 6-leaf wheat, oats, and foxtail millet on June 21 with 87 F. Evaluations were on July 8 and August 4.

Treatment	Rate <sup>a</sup>	=	July	/ 8		August	1
	(1b/A)	For	ni Whi		Fon	ni Wh+	Oats
Sethoxydim+OC L				(% con	trol)		
Sethoxydim+BCH BAS-0562-16H BAS-0562-16H+BCH	0.1+0.25G 0.1+0.25G 0.1+0.5G 0.125+0.25G 0.125+0.25G 0.125+0.5G 0.15+0.25G 0.15+0.5G 0.1 0.1+0.25G 0.125 0.125 0.125 0.125+0.25G 0.15+0.25G 0.15+0.25G 0.15+0.25G	78 3 82 81 82 85 84 88 90 82 85 82 85 83 87 89		73 4 83 78 86 85 83 90 91 79 83 83 90 81 86 85 89 94	93 07 97 95 96 97 98 99 94 97 99 99 99		77 0 98 99 99 99 99 99 99 99 99 99 99
DW MEAN (P MEAN V. % ID 5%		90 3 79 6	94 0 82 5	96 4 82 5	99 0 90 3	99 0 91	99 0 91
D 1%		6	6		3	3 4	4
OF REPS		8	8	6 8	4	6	5 7
G in the rate colu		4	4	4	4	O	4

a G in the rate column is for gallon/A.

#### Summary

BAS-0562 without an adjuvant generally gave similar grass species control as sethoxydim with BCH (adjuvant from BASF, Dash). BCH only enhanced BAS-0562 at

b Sethoxydim probably was not a part of the treatment.

Sethoxydim formulation with adjuvants, Exp 2, Fargo 1988. 'McCall' soybeans were seeded to an area with a natural infestation of yellow foxtail on May 19. Treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft wide area the length of 10 by 25 ft plots. Treatments were applied to first trifoliolate soybeans and by 25 ft plots. Treatments were applied to first trifoliolate soybeans and 3- to 4-leaf foxtail on lune 22. Evaluation was on July 14 3- to 4-leaf foxtail on June 22. Evaluation was on July 14.

- to 4-leaf foxtail		Soybean	Grft
	Rate	Injury	% control)
reatment	(1b/A)	(%)	% Concror,
	(10/14)		94
	0.1+0.25G	4	97
ethoxydim+OC	0.1+0.25G	4	89
ethoxydim+BCH	0.1+0.25G	4	98
ethoxydim+BCH	0.125+0.25G	3	99
ethoxydim+UC	0.125+0.25G	4	97
Sethoxydim+BCH	0.125+0.5G	4	97
Sethoxydim+BCH	0.15+0.25G	4	99
Sethoxydim+UC	0.15+0.25G	3	99
Sethoxydim+BCH	0.15+0.5G	5	95
Sethoxydim+BCH	0.1	4	95
RAS-0562-16H	0.1+0.25G	4	96
RAS-0562-16H+0C	0.1+0.25G	3 3 4	96
BAS-0562-16H+BCH	0.125	3	98
RAS-0562-16H	0.125 0.125+0.25G		98
RAS-0562-16H+0C	0.125+0.25G	4	98
BAS-0562-16H+BCH	0.125+0.254	4	99
RAS-0562-16H	0.15+0.25G	3	96
RAS-0562-16H+UL	0.15+0.125G	4	99
BAS-0562-16H+BCH	0.15+0.25G	3	95
BAS-0562-16H+BCH	0.15+0.25G	4	95
Fluazifop+PO	0.15+0.254		99
		5 3	89
HIGH MEAN		3	97
LOW MEAN		.4 32	4 .
EXP MEAN		NS NS	5
C.V. %		NS NS	NS
LSD 5%		NS 4	4
LSD 1%		4	
# OF REPS			

All treatment None of the treatments caused important injury to soybean. similarly gave 89% or more control of yellow foxtail.

Sethoxydim time-of-day application, Exp 1, Fargo 1988. 'Wheaton' Hard Red Spring wheat and 'ND810104' oats were seeded as bioassay species in adjacent 10 ft wide strips at Fargo, ND on April 28. Treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft wide area across the bioassay species. Plots were 10 ft wide to provide a 4 ft untreated area between treated areas to aid evaluations. Treatments were applied to 5- to 6-leaf wheat and oats on June 7 with clear sky and 87 F at 3.00 am, 98 F at 2:00 pm, and 85 F at 8:00 pm applications. Evaluation was on June 22.

		PP: Teactons.	Evaluation was o
Treatment		Digital Control of the Control of th	
		Rate	111-
		(oz/A)	Wheat Oats
Sethoxydim+PO	(8 am)		(% control)
Sethoxydim+Msc	(8 am)	1+0.25G	77 40
Sethoxydim+BCH	(8 am)	1+0.25G	
Sethoxydim+PO	(8 am)	1+0.25G	87 80 87 79
Sethoxydim+AG2	(8 am)	1+0.25G 1+0.25G	34 11
Sethoxydim+PO	(2 pm)	1+0.25G 1+0.25G	78 49
Sethoxydim+MSC Sethoxydim+BCH	(2 pm)	1+0.25G	82 53
Sethoxydim+PO	(2 pm)	1+0.25G	89 87
Sethoxydim+AG2	(2 pm)	1+0.25G	93 86
Sethoxydim+PO	(2 pm)	1+0.25G	80 53
Sethoxydim+MSC	(8 pm)	1+0.25G	87 80
Sethoxydim+BCH	(8 pm) (8 pm)	1+0.25G	88 76
Sethoxydim+PO	(8 pm) (8 pm)	1+0.25G	91 91
Sethoxydim+AG2	(8 pm)	1+0.25G	92 90 87 75
	( • pm)	1+0.25G	
HIGH MEAN			89 83
LOW MEAN EXP MEAN			93 91
C.V. %			34 11
LSD 5%			83 69
LSD 1%			13 14
# OF REPS			15 13
- OF INELS			21 18
			4 4

## Summary

Sethoxydim control of wheat and oats generally increased from 8:00 am to 8:00 pm applications when PO (petroleum oil 11N plus 77% Atplus), SOTM (once refined soybean oil with 15% T-Mulz vo) or AG2 were the adjuvants. Time of application did not influence control when MSC (methylated seed oil ) or BCH (adjuvant from BASF, Dash) were the adjuvants. Oats control, averaged over time of sehoxydim application was 56% with PO, 86% with MSC, 83% with BCH, and

Sethoxydim time-of-day application, Exp 2, Prosper 1988. 'Wheaton' Hard Red Spring wheat, 'ND810104' oats, and Siberian foxtail millet were seeded as bioassay species in adjacent 6 ft wide strips at Prosper, ND on May 23. Treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 Treatments were applied with a cross the bioassay species. Plots were 10 gpa at 35 psi to an 8 ft wide area across the bioassay species. Plots were 10 ft wide to provide a 4 ft untreated area between treated areas to aid ft wide to provide a 4 ft untreated area between treated areas to aid a clear sky and 75 F at 8:00 am, partly cloudy sky and 85 F at 2:00 pm, and a clear sky and 75 F at 8:00 pm applications. Evaluation was on July 19 partly cloudy sky and 68 F at 8:00 pm applications. Evaluation was on July 19 and August 10.

nd August 10.			July	19		August	10
	Rate	Fomi	Wht	Nats	Fomintrol)	Wht	<u> </u>
reatment	(oz/A)			(% 00			0.7
	1+0.25G	65	75	70	74 75	97 98	87 95
eth+PO (8 am) eth+MSC (8 am)	1+0.25G	64	76 78	78 81	76	99	96
eth+MSC (8 am) eth+BCH (8 am)	1+0.25G	68 57	64	56	68	86	84
Seth+PO (8 am)	1+0.25G	58	74	70	73	97	83
Seth+AG2 (8 am)	1+0.25G 1+0.25G	60	73	72	75	97 98	87 97
Seth+PO (2 pm)	1+0.25G	58	76	77	74 75	99	98
Seth+MSC (2 pm) Seth+BCH (2 pm)	1+0.25G	64	77 51	78 60	67	82	80
Seth+BCH (2 pm) Seth+PO (2 pm)	1+0.25G	46 50	68	72	71	96	88
Seth+AG2 (2 pm)	1+0.25G 1+0.25G	63	74	72	76	99 98	90 96
Seth+PO (8 pm)	1+0.25G	73	78	78	76 77	99	98
Seth+MSC (8 pm) Seth+BCH (8 pm)	1+0.25G	68	78 73	79 65	75	92	78
Seth+BCH (8 pm) Seth+PO (8 pm)	1+0.25G	62 66	77	76	75	99	91
Seth+AG2 (8 pm)	1+0.25G	00				00	98
		73	78	81	77	99 82	78
HIGH MEAN		46	51	56 72	67 74	96	90
LOW MEAN EXP MEAN		61	73 6	8		4	6
C.V. %		9		8	3	5	8
LSD 5%		11	6 8 4	11	4	7 4	10
LSD 1% # OF REPS		4	4	4	4	4	

## Summary

Time-of-day that sethoxydim was applied did not influence grass species control regardless of adjuvants. Grass control was greatest with MSC (methylated seed oil) and BCH (adjuvant from BASF, Dash) adjuvants.

Sethoxydim mixing method, Exp 1, Fargo 1988. 'Wheaton' Hard Red Spring wheat and 'ND 810104' oats were seeded as bioassay species in adjacent 10 ft wide strips at Fargo, ND on April 28. Treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft wide area across the bioassay species. Plots were 10 ft wide to provide a ft untreated area between treated areas to aid evaluations. Treatment were applied to 4- to 5-leaf wheat and oats on June 3 with 85 F.

	Rate	Wheat	Oats
Sethoxydim+BCH (at trt)	(oz/A)	(% cor	ntrol)
Sethoxydim+BCH (Form. 24H) Sethoxydim+BCH (Form.+H20 12H) Sethoxydim+BCH (Form.+H20 24H) Sethoxydim+MS (at trt) Sethoxydim+MS (Form. 24H) Sethoxydim+MS (Form.+H20 12H) Sethoxydim+MS (Form.+H20 24H) Sethoxydim+MSC (at trt) Sethoxydim+MSC (Form. 24H) Sethoxydim+MSC (Form. 24H) Sethoxydim+MSC (Form. 24H) Sethoxydim+MSC (Form.+H20 12H) Sethoxydim+MSC (Form.+H20 24H) HIGH MEAN LOW MEAN LOW MEAN EXP MEAN C.V. % LSD 5% LSD 1% # OF REPS	1+0.25G 1+0.25G 1+0.25G 1+0.25G 1+0.25G 1+0.25G 1+0.25G 1+0.25G 1+0.25G 1+0.25G 1+0.25G	54 36 56 38 76 45 50 46 60 58 58 51 76 36 52 27 20 NS 4	69 33 71 31 83 63 67 54 73 72 64 69 83 31 62 29 26 35 4

## Summary

Sethoxydim mixed with BCH (adjuvant from BASF, Dash) as formulants or formulants plus water for 24h before application reduced grass control compared to sethoxydim mixed and applied immediately or after 12h. With water for 12h or more before application reduced control compared to application immediatly after mixing. Pre-mixing sethoxydim with MSC did control than the other adjuvants with sethoxydim.

Sethoxydim mixing method, Exp.2, Prosper 1988. 'Wheaton' Hard Red Spring wheat, 'ND810104' oats, and Siberian foxtail millet were seeded as bioassay species in adjacent 6 ft wide strips at Prosper, ND on May 23. Treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 with an 8 ft wide area across the bioassay species. Plots were 10 ft wide psi to an 8 ft untreated area between treated areas to aid evaluation. to provide a 4 ft untreated area between treated areas to aid evaluation. Treatments were applied to 5- to 6-leaf wheat and oats on June 28 with 85 F. Evaluation was on July 18 and August 10.

Treatment	Rate (oz/A)		ly 18 Wht (	oats cont	Aug Fomi crol)	wst 10 Wht C	<u>lats</u>
Seth+BCH (at trt) Seth+BCH (Form. 24H) Seth+BCH (Form.+H20 12H) Seth+BCH (Form.+H20 24H) Seth+MS (at trt) Seth+MS (Form. 24H) Seth+MS (Form.+H20 12H) Seth+MS (Form.+H20 12H) Seth+MSC (at trt) Seth+MSC (Form. 24H) Seth+MSC (Form. 24H) Seth+MSC (Form.+H20 12H) Seth+MSC (Form.+H20 12H) Seth+MSC (Form.+H20 24H)  HIGH MEAN LOW MEAN EXP MEAN C.V. % LSD 5% LSD 1% # OF REPS	1+0.25G 1+0.25G 1+0.25G 1+0.25G 1+0.25G 1+0.25G 1+0.25G 1+0.25G 1+0.25G 1+0.25G 1+0.25G	64 62 63 68 60 65 65 58 67 67 63 67 68 58 64 5	77 77 79 78 76 76 77 70 77 79 77 76 3 3 3 5 4	79 79 76 78 73 76 75 75 74 75 78 79 73 76 3 NS NS	75 74 76 76 74 75 74 76 75 77 77 74 75 3 NS NS	99 99 98 98 98 98 98 98 98 98 98 98 1 1 NS 4	92 92 94 96 87 88 91 89 89 89 87 91 4 5 NS 4

## Summary

Sethoxydim with all adjuvants gave similar grass species control regardless if mixed as formulation or with water for 24h before application.

Sethoxydim formulations and bentazon antagonism, Prosper 1988. 'Wheaton' Hard Red Spring wheat, 'ND810104' oats, and siberian foxtail millet were seeded as bioassay species in adjacent 6 ft wide strips at Prosper, ND on delivering 8.5 gpa at 35 psi to an 8 ft wide area across the bioassay species. Plots were 10 ft wide to provide a 4 ft untreated area between treated areas to aid evaluation. Treatments were applied to 4- to 6-leaf wheat, oats, and foxtail millet on June 21 with 87 F. Evaluation was July

		Lvarua	CION W	las July
Treatment				
	Rate			
	(1b/A)	Fom	i Wht	Oats
Coth	(10)11)	(	% cont	rol)
Sethoxydim+PO	0.1.0.0=0	1	CONT	101)
Sethoxydim+PO	0.1+0.25G	00	0.1	
Seth+Bent+PO+UAN	0.15+0.25G	80	81	77
Seth+Bent+PO+UAN	0.1+0.75+0.25G+1G	86	88	90
Sothowiding	0.15+0.75+0.25G+1G	73	63	50
Sethoxydim+Dash	0.1+0.5G	76	78	78
Sethoxydim+Dash	0.170.56	82	85	
Seth+Bent+Dash+UAN	0.15+0.5G	93		85
Seth+Bent+Dash+UAN	0.1+0.75+0.5G+1G		92	91
BAS-562-16H	0.15+0.75+0.5G+1G	79	79	75
BAS-562-16H	0.1	84	86	80
BAS FED 1611 P	0.15	80	76	73
BAS-562-16H+Bent+UAN	0.1+0.75+1G	84	-87	90
BAS-562-16H+Bent+UAN	0.15.0.75.16	74	71	
DAS-562-16H+Dach	0.15+0.75+1G	82		54
BAS-562-16H+Dach	0.1+0.25G		82	79
BAS562+Bent+Dash+UAN	0.15+0.25G	91	97	97
BAS562+Bent+Dash+UAN	0.1+0.75+0.256+16	88	91	93
SHOSOZ+Bent+Dash+UAN	0.15+0.75+0.25G+1G	77	76	77
HTOU MEAN	20.70.23416	81	85	83
HIGH MEAN			00	03
LOW MEAN		93	0.7	
EXP MEAN			97	97
C.V. %		73	63	50
LSD 5%		82	82	79
LSD 1%		5	6	11
L3U 1%		6	7	12
# OF REPS		8	9	
and delice the same and the same and		4		16
		4	4	4
	Plantin Single William St. St. Co.			EE CHEE

## Summary

Sethoxydim + PO or + PO and DASH and BAS-562-16H alone or + DASH were similarly antagonized by bentazon + UAN. Sethoxydim + PO and BAS-562-16H alone gave similar grass species control. PO = petroleum oil with 17% Atplus 300 F, Dash = an adjuvant from BASF, and UAN = a urea + ammonium nitrate fertilizer.

Sethoxydim plus adjuvants, Fargo 1988. 'Wheaton' Hard Red Spring wheat and 'ND810104' oats were seeded as bioassay species in adjacent 10 ft wide strips at Fargo, ND on April 28. Treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft wide area wheel type plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft wide area cross the bioassay species. Plots were 10 ft wide to provide a 4 ft across the bioassay species. Plots were 10 ft wide to provide a 4 ft across the bioassay species area to aid evaluations. Treatments were untreated area between treated areas to aid evaluations. Treatments were applied to 4- to 5-leaf wheat and oats on June 3 with 85 F and clear sky. Evaluation was on June 24 and July 19.

Evaluation was on J	une 21 die	Wht	Oats	Wht	Oats
Twoatment.	Rate		- (% CO	ntrol)	
Sethoxydim+PO Sethoxydim+PO Sethoxydim+PO Sethoxydim+MSC Sethoxydim+MSC Sethoxydim+MSC Sethoxydim+AG-2 Sethoxydim+AG-2 Sethoxydim+AG-2 Sethoxydim+AG-2 Sethoxydim+AG-2	(oz/A)  0.5+0.25G 1+0.25G 1.5+0.25G 0.5+0.25G 1+0.25G 0.5+0.25G 1+0.25G 1+0.25G 0.5+0.25G 0.5+0.25G	11 44 80 28 71 91 30 61 77 44	23 55 78 40 78 93 38 77 82 46 84	5 19 45 9 26 68 6 30 46 16 51	0 16 38 9 25 63 3 35 8 56 85
Sethoxydim+BCH Sethoxydim+BCH HIGH MEAN LOW MEAN EXP MEAN C.V. % LSD 5% LSD 1% # OF REPS	1+0.25G 1.5+0.25G	91 91 11 59 21 18 24 4	91 93 23 65 19 18 24 4	80 80 5 33 20 10 13 4	85 0 31 25 11 15 4

## Summary

Control of wheat and oats with sethoxydim with MSC or BCH was similar at the first evaluation, but greater with BCH at the second evaluation. MSC and BCH both were more effective as adjuvants than PO or AG-2.

Fluazifop plus adjuvants, Fargo 1988. 'Wheaton' Hard Red Spring wheat, 'ND810104' oats, and Siberian foxtail millet were seeded as bioassay were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at wide to provide a 4 ft area between treated areas to aid evaluation. 8 with 85 F and partly cloudy sky. Evaluation were on August 2.

Treatment   Rate	Rate   Wht   Fomi   Oats   Oz/A   Fomi   Oats   Oz/A			magast Z.	
Rate	Rate	Treatment			
Coz/A    Whit Fomi Oats   Fomi Oats   Fomi Oats	(oz/A)       Wht Fomi Oats         Fluazifop+P0         Fluazifop+P0       0.25+0.25G       6       19       8         Fluazifop+P0       1+0.25G       35       86       68       19         Fluazifop+MSC       0.25+0.25G       35       86       68       6       16       11       75       53       86       68       68       69       Fluazifop+AG-2       1+0.25G       31       18       8       15       40       15       40       15       40       15       40       15       40       15       40       15       40       15       40       15       40       15       40       15       40       15       40       15       40       15       40       15		Rate		
Fluazifop+PO	Fluazifop+PO Fluazifop+PO Fluazifop+PO Fluazifop+PO Fluazifop+MSC Fluazifop+MSC Fluazifop+MSC Fluazifop+MSC Fluazifop+MSC Fluazifop+MSC Fluazifop+AG-2 Fluazifop+AG-2 Fluazifop+AG-2 Fluazifop+BCH Flu		(oz/A)		Oats
100 100	# OF REPS 8 13 11 17 14	Fluazifop+PO Fluazifop+PO Fluazifop+PO Fluazifop+MSC Fluazifop+MSC Fluazifop+AG-2 Fluazifop+AG-2 Fluazifop+BC-2 Fluazifop+BCH Fluazifop+BCH Fluazifop+BCH Fluazifop+BCH COW MEAN COW MEAN EXP MEAN COW MEAN COW MEAN	(oz/A)  0.25+0.25G  0.5+0.25G  1+0.25G  0.25+0.25G  0.5+0.25G  0.25+0.25G  0.5+0.25G  0.5+0.25G  0.5+0.25G  0.5+0.25G	(% cont) 6 19 9 65 35 86 6 16 11 75 36 85 3 18 15 40 40 82 0 21 11 69 46 85 46 86 0 16 18 55	8 19 68 11 53 69 8 15 68 6 30 65 69 6 35

## Summary

Grass species control with fluazifop was similar with all adjuvants. However, fluazifop at  $0.5\ \text{oz/A}$  tended to give greater control when with MSC than the other adjuvants.

Clethodim plus adjuvants, Farqo 1988. 'Wheaton'Hard Red Spring wheat, "ND810104' oats, and Siberian foxtail millet were seeded as bioassay species in adjacent 6 ft wide strips at Prosper, ND on May 23. Treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi applied a 4 ft untreated area between treated areas to aid evaluation. Treatments were applied to 4- to 6-leaf wheat, oats, and foxtail millet on June 22 with 83 F and partly cloudy sky. Evaluation was on July 7 and August 8.

august 0.					^	uguet	8
	Rate	Fomi	July 7 Wht	Nate	Fomi ntrol)	ugust Wht	Oats
Treatment Clethodim+PO Clethodim+PO	(oz/A) 0.25+0.25G 0.5+0.25G	71 79 83	64 77 86	54 80 90	66 81 85	67 79 98	38 92 99
Clethodim+PO Clethodim+MSC Clethodim+MSC Clethodim+AG-2 Clethodim+AG-2 Clethodim+AG-2 Clethodim+BCH Clethodim+BCH Clethodim+BCH Clethodim+BCH Clethodim+BCH	0.75+0.25G 0.25+0.25G 0.5+0.25G 0.75+0.25G 0.25+0.25G 0.75+0.25G 0.25+0.25G 0.5+0.25G 0.75+0.25G 0.75+0.25G 0.75+0.3%	77 81 85 72 78 80 77 82 82 75	78 84 89 74 79 87 77 90 90 78	78 87 95 74 80 88 76 92 91 80 88 92	64 77 85 61 68 77 70 86 83 67 81 83	67 96 99 67 79 99 88 99 77 98	95 99 99 60 97 99 93 99 91 99
Clethodim+MS Clethodim+MS HIGH MEAN LOW MEAN	0.75+0.3%	83 85 71 79	90 64 82	95 54 83	86 61 75	99 67 87	99 38 90
EXP MEAN C.V. % LSD 5% LSD 1% # OF REPS		4 4 5 4	5 6	8 10 13 4	7 8 11 4	7 8 11 4	6 8 11 4

#### Summary

MSC, BCH, and MS similarly enhanced grass species control with clethodim. PO and AG-2 were generally less effective than the above adjuvants with clethodim at 0.5 oz/A.

DPX-Y6202 plus adjuvants, Exp 1, Fargo 1988. 'Wheaton' Hard Red Spring wheat, 'ND810104' oats, and Siberian foxtail millet were seeded as bioassay species in adjacent 6 ft wide strips at Fargo, ND on May 31. Treatments psi to an 8 ft wide area across the bioassay species. Plots were 10 ft wide provide a 4 ft untreated area between treated areas to aid evaluation. Treatments were applied to jointing wheat, oats, and foxtail millet on July August 22.

Treatment   Rate   Fomi   Wht   Oats   Fomi   Wht   Oats   Oats							•	Lo and
Coz/A   Coz/A   Control   Color   Control   Color   Control   Color   Control   Color   Control   Color   Control   Color   Control	<u>Treatment</u>	Pato		July	29		August	- 00
DPX-Y6202-38+P0 DPX-Y6202-38+P0 DPX-Y6202-38+P0 DPX-Y6202-38+P0 DPX-Y6202-38+MSC DPX-Y6202-38+MSC DPX-Y6202-38+MSC DPX-Y6202-38+MSC DPX-Y6202-38+MSC DPX-Y6202-38+MSC DPX-Y6202-38+MSC DPX-Y6202-38+BC DPX-Y6202-38+BC DPX-Y6202-38+BC-2 DPX-Y6202-38+BC-2 DPX-Y6202-38+BCH DPX-Y6202-38+SOC-TM DPX-Y			Fom	i Wht		Fomi	August	
DPX-Y6202-38+P0 DPX-Y6202-38+P0 DPX-Y6202-38+P0 DPX-Y6202-38+BCC DPX-Y6202-38+MSC DPX-Y6202-38+MSC DPX-Y6202-38+MSC DPX-Y6202-38+MSC DPX-Y6202-38+MSC DPX-Y6202-38+MSC DPX-Y6202-38+AG-2 DPX-Y6202-38+AG-2 DPX-Y6202-38+AG-2 DPX-Y6202-38+BCH DPX-Y6202-38+SOC-TM DPX-Y6202-38-SOC-TM DPX-Y		(02/A)				ntvol	wnt	<u> </u>
LSD 5% LSD 1%  8 12 6 8 11 12  # OF REPS  11 15 9 11 14 17  4 4 4 4 4 4	DPX-Y6202-38+P0 DPX-Y6202-38+P0 DPX-Y6202-38+MSC DPX-Y6202-38+MSC DPX-Y6202-38+MSC DPX-Y6202-38+AG-2 DPX-Y6202-38+AG-2 DPX-Y6202-38+BCH DPX-Y6202-38+BCH DPX-Y6202-38+BCH DPX-Y6202-38+SOC-TM DPX-Y6202-38+SOC-TM DPX-Y6202-38+SOC-TM DPX-Y6202-38+SOC-TM DPX-Y6202-38+SOC-TM DPX-Y6202-38+SOC-TM DPX-Y6202-38+SOC-TM DPX-Y6202-38+SOC-TM DPX-Y6202-38+SOC-TM C.V. % LSD 5% LSD 5% LSD 1%	0.12+0.25G 0.38+0.25G 0.75+0.25G 0.12+0.25G 0.38+0.25G 0.75+0.25G 0.12+0.25G 0.75+0.25G 0.38+0.25G 0.38+0.25G 0.75+0.25G 0.75+0.25G 0.75+0.25G	70 84 18 64 79 28 65 78 25 74 89 5 21 59 89 5 53 10 8	99 99 81 99 99 63 99 78 99 17 31 84 99 17 82 10 12	9 34 73 7 30 76 13 27 75 6 63 86 8 8 20 86 6 35 13 6 9	21 35 67 11 32 45 6 30 36 9 41 74 3 8 28 74 3 30 19 8 11	96 99 99 99 99 99 58 99 99 15 70 96 99 15 87 91 14	24 52 74 16 41 82 16 41 69 11 68 83 5 7 14 83 5 40 22 12

#### Summary

PO (petroleum oil with 17% Atplus 300F), MSC (methylated seed oil from Agsco), and BCH (adjuvant from BASF, DASH) all were similar as adjuvants with DPX-Y6202. AG-2 was less effective then the above as an adjuvant only VO) was ineffective as an adjuvant with DPX-Y6202.

DPX-Y202 plus adjuvants, Exp 2, Fargo 1988. 'Wheaton' Hard Red Spring wheat, 'ND810104' oats, and Siberian foxtail millet were seeded as bioassay species in adjacent 6 ft wide strips at Fargo, ND on May 31. Treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 were applied area across the bioassay species. Plots were 10 ft wide psi to an 8 ft wide area across the bioassay species. Plots were 10 ft wide psi to an 8 ft untreated area between treated areas to aid evaluation. to provide a 4 ft untreated area between treated areas to aid evaluation. Treatments were applied to jointing to boot wheat, oats, and foxtail millet Treatments were applied to jointing to boot wheat, oats, and foxtail millet august 8 and 29.

August 8 and 29.		Δι	igust 8	8	Auc		9
reatment	Rate	Fomi	Wht	Oats	Fomi trol)-	Wht	<u>Oats</u>
DPX-Y6202-38+PO DPX-Y6202-38+PO DPX-Y6202-38+PO DPX-Y6202-38+MSC DPX-Y6202-38+MSC DPX-Y6202-38+MSC DPX-Y6202-38+AG-2 DPX-Y6202-38+AG-2 DPX-Y6202-38+BCH DPX-Y6202-38+BCH DPX-Y6202-38+BCH DPX-Y6202-38+BCH DPX-Y6202-38+SOC-TM HIGH MEAN LOW MEAN EXP MEAN C.V. % LSD 5% LSD 1% # OF REPS	(oz/A)  0.12+0.25G 0.38+0.25G 0.75+0.25G 0.12+0.25G 0.75+0.25G 0.12+0.25G 0.12+0.25G 0.12+0.25G 0.75+0.25G 0.12+0.25G 0.38+0.25G 0.75+0.25G 0.75+0.25G 0.75+0.25G 0.75+0.25G	9 58 70 3 44 65 4 29 63 20 55 69 0 3 10	8 54 70 8 48 70 6 34 36 16 71 7 10 26 71 6 36 20 10	5 50 74 6 46 71 4 10 43 13 71 77 0 5 13 77 0 32 22 10 13 4	10 39 55 8 33 49 5 24 42 63 3 1 5 63 1 26 16 8 4	23 64 64 22 49 66 18 43 53 25 64 63 14 26 41 66 14 42 20 12 16 4	14 37 62 11 32 56 9 19 43 16 50 64 6 11 18

## Summary

Results from experiment 2 were in close agreement with those from experiment 1, but the level of control was less than in experiment 1. The lower control in experiment 2 may have reflected the more mature plants or the lower temperatures at treatment. PO, MSC, and BCH were all similar as adjuvants with DPX-Y6202. AG-2 tended to be less effective then the above adjuvants. SOC-TM was ineffective as an adjuvant with DPX-Y6202.

Fenoxaprop plus adjuvants, Prosper 1988. 'Wheaton' Hard Red Spring wheat, 'ND810104' oats, and Siberian foxtail millet were seeded as bioassay species in adjacent 6 ft wide strips at Prosper, ND on May 23. Treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at wide to provide a 4 ft untreated area between treated areas to aid evaluation. Treatments were applied to 5- to 6-leaf wheat, oats, and foxtail millet on July 5 with 103 F, 30% RH, and partly cloudy sky.

Treatment	Rate	Fomi	Augus		Ford	Augus	
Fenoxaprop+PO Fenoxaprop+PO Fenoxaprop+PO Fenoxaprop+MSC Fenoxaprop+MSC Fenoxaprop+AG-2 Fenoxaprop+AG-2 Fenoxaprop+AG-2 Fenoxaprop+AG-2	Rate (oz/A) 0.5+0.25G 1+0.25G 1.5+0.25G 0.5+0.25G 1.5+0.25G 0.5+0.25G 1+0.25G 1.5+0.25G	Fomi 36 65 69 45 53 64 41 59	9 12 11 10 13 10 6 10	0ats -(% co 5 18 23 12 18 21 5	Fomi ntrol 23 42 45 28 43 49 22 44	Wht.	11 27 36 14 33 29 24 29
Fenoxaprop+BCH Fenoxaprop+BCH Fenoxaprop+BCH C.V. % LSD 5% LSD 1%	0.5+0.25G 1+0.25G 1.5+0.25G	67 57 63 70 10 8	13 12 11 19 41 NS	35 9 23 29 34 9	46 37 38 50 11 6	46 38 45 55	44 29 32 36
# OF REPS		11 4	NS 4	12	9	12 16 4	9 11 4

## Summary

All adjuvants were similar with fenoxaprop for the control of wheat, oats and foxtail millet. The rates probably were to low for the large plant. The high temperature at treatment apparently did not enhance phytotoxicity of fenoxaprop.

Haloxyfop plus adjuvants, Fargo 1988. 'Wheaton' Hard Red Spring wheat and 'ND810104' oats were seeded as bioassay species in adjacent 10 ft wide strips at Fargo, ND on April 28. Treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft wide area wheel type plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft wide area across the bioassay species. Plots were 10 ft wide to provide a 4 ft untreated area between treated areas to aid evaluations. Treatments were applied to 5- to 6-leaf wheat, oats, and foxtail millet on June 10 with 89 F, 50% RH and clear sky. Evaluation were on June 24 and July 19.

F, 50% KH allu Cicui			7.7.10			
			June		July	
Treatment	Rate (oz/A)	Rate	Wht 	<u>0ats</u> (% co	Wht ntrol)	<u> </u>
Haloxyfop+PO Haloxyfop+PO Haloxyfop+PO Haloxyfop+MSC Haloxyfop+MSC Haloxyfop+AG-2 Haloxyfop+AG-2 Haloxyfop+AG-2 Haloxyfop+BCH Haloxyfop+BCH Haloxyfop+BCH C.V. % LSD 5% LSD 1% # OF REPS	0.25+0.25G 0.50+0.25G 0.75+0.25G 0.25+0.25G 0.50+0.25G 0.25+0.25G 0.50+0.25G 0.75+0.25G 0.25+0.25G 0.75+0.25G 0.75+0.25G 0.75+0.25G		39 75 86 61 84 84 64 81 86 65 70 86	50 78 89 75 85 91 75 88 90 69 78 92 11 12 16 4	14 76 90 64 92 95 60 89 95 51 90 93	11 82 97 77 98 98 71 98 99 60 92 99

## Summary

MSC (methylated seed oil adjuvant from Agsco) and AG-2 were similar and more effective as adjuvants then PO (petrolium oil 11N with 17% Atplus 300F) with haloxyfop for grass species control. BCH (adjuvant from BASF-DASH) was intermediate to PO and MSC or AG-2.

Sethoxydim with NaHCO3 or Na2CO3, Fargo 1988. wheat and 'ND810104' oats were seeded as bioassay species in adjacent 10 ft wide strips at Fargo, ND on April 28. Treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft wide area across the bioassay species. Plots were 10 ft wide to provide a 4 ft untreated area between treated areas to aid evaluations. Treatments were applied to 5- to 6-leaf wheat and oats on June 6 with 98 F, 28% RH, and clear sky. Evaluations were on June 24 and July 20.

<u>Ireatment</u> Salt	Rate		Jun	e 24	Jul	y 20
(ppm)	(oz/A)		Wht	Oats	Wht	Oat
Seth+PO				- (% c	ontrol	)
Seth+PO+NaHCO3 (500)	1.5+0.25G		70			
Setu+PO+NaHCO3 (1000)	1.5+0.25G+0.5		75	74	23 1	4
Seth+PO+NaHCO2 idooc	1.5+0.25G+1		74	56	30 1	8
Seth+PO+NaHCO3 (1000)	1.5+0.25G+4		62	33	17	6
Setn+PO+Na2CO3 /FOOL	1.5+0.25G+16		34	28	10	3
Setn+PO+Na2CO3 /1000	1.5+0.25G+0.5			13	8	3
Seth+PO+Na2CO3 (4000)	1.5+0.25G+1				21 1:	2
Seth+PO+Na2CO3 (16000)	1.5+0.25G+4		1	44	15	7
Seth+PO ( )	1.5+0.25G+16			8	6	3
Seth+PO+NaHCO3 (500)	3+0.25G	9	0	0	4 ]	-
Seth+PO+NaHCO2 /1000	3+0.25G+0.5	8			73 61	
Setu+PO+NaHCO3 (4000)	3+0.25G+1				31	
Seth+PO+NaHCO3 (16000)	3+0.25G+4	8.			0 24	
Setn+PO+Na2CO3 (500)	3+0.25G+16	76	_		3 15	
Seth+PO+Na2CO3 /1000	3+0.25G+0.5	68		_		
Seth+PO+Na2CO3 (4000)	3+0.25G+1	83				
Setn+PO+Na2CO3 (16000)	3+0.25G+4	80				
petn+PO+28%N	3+0.25G+16	65			0 9	
eth+PO+NaHCO3 (FOO)	1.5+0.25G+0.06G	33 77			0 9	
eth+PO+NaHCO3 (1000) 2000	1.5+0.25G+0.5+0 nsc	80	67		15	
eth+PO+NaHCO2 (4000)	1.5+U.25G+1+0 06C	81	76			
eth+PO+NaHCO3 /16000)	1.5+U.25G+4+0 06C	85	74			
etn+PO+Na2CO3 (500) 2000	1.5+0.25G+16+0 nec	38	83			
eth+PO+Na2CO3 /1000)	1.5+U.25G+0 5+0 06C	84	34	-		
eth+P0+Na2C03 (4000)	1.3+U.25G+1+0 06C	80	80	51	40	
th+PO+Na2CO3 (16000) -0000	1.5+U.25G+4+0 nsc	65	78	41	33	
tn+P0+28%N	1.5+U.25G+16+0 nec	0	43	17	6	
th+PO+NaHCO3 (500)+28%N	3+U.25G+0 25G	94	0	0	0	
th+PO+NaHCO3 (1000) -0000	3+U.25G+U.5+D 25C	95	94 95	92	92	
Th+PO+NaHCO3 /40001 20001	3+U.25G+1+D 25C	93	94	95	95	
th+PO+NaHCO3 (16000)	3+0.25G+4+0.25G	94	93	93	94	
th+PO+Na2CO3 (500)+28%N	3+U.25G+16+0 25C	95	96	93	93	
th+PO+Na2CO3 (1000) .0000	3+0.25G+0.5+0.25G	96		96	97	
n+P0+Na2C03 (4000) .0000	3+U.25G+1+0 25C	96	96 05	97	97	
th+PO+Na2CO3 (16000)+28%N	3+0.25G+4+0.25G	95	95	98	98	
	3+0.25G+16+0.25G	80	94	97	97	
. %		00	78	33	17	
5%		8	10	0.0		
F REPS		8	12	22	18	
		4	10	13	9	

Antagonism of sethoxydim was greater by Na2CO3 than NaHCO3 in the spray mixture. Antagonism began with the lowest amount of carbonates, 500 ppm, and increased as concentration increased. Generally 0.066 gpa 28%N overcame 4000 ppm NaHCO3 and 1000 ppm Na2CO3 antagonisms of sethoxydim at 1.5 oz/A.

28%N at 0.25G overcame antagonism of sethoxydim at 3 oz/A caused by 16000 ppm NaHCO3 and 4000 ppm Na2CO3.

DPX-R9674 with adjuvants, Fargo 1988. 'Wheaton' Hard Red Spring wheat, Fl sunflower, 'McCall' soybean and redroot pigweed were seeded in 6 ft wide adjacent strips as bioassay species on May 18. Treatments were applied to 6-adjacent redroot pigweed and sunflower, 4- to 6-leaf wheat, and 3 to 8-leaf redroot pigweed and sunflower, 4- to 6-leaf wheat, and 3 trifolialate soybeans on June 15 with 72 F and partly cloudy sky. Application trifolialate soybeans on June 15 with 72 F and partly cloudy sky. Application was with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi to an was with a bicycle wheel type plots across the various species. The experiment 8 ft wide area of 10 ft wide plots across the various species. Evaluations were on June 28 and July 21.

June 28 and July	21.						July	21	
			June	28	KOCZ	Soy	Wht	Vosf	Rrpw
	Rate	Rrpw	Vosf	Wht	% con	trol)			
Treatment  DPX-R9674+X-77 DPX-R9674+MSC DPX-R9674+MSC DPX-R9674+P0 DPX-R9674+AG-2 DPX-R9674+AG-2 Untreated DPX-R9674+MSC DPX-R9674+MSC DPX-R9674+MSC DPX-R9674+P0 DPX-R9674+P0 DPX-R9674+AG-2 Untreated DPX-R9674+AG-2 Untreated DPX-R9674+AG-7  HIGH MEAN LOW MEAN EXP MEAN C.V. % LSD 5% LSD 1% # OF REPS	(oz/A)  0.06+0.25% 0.06+0.12G 0.06+0.25G 0.06+0.12G 0.06+0.12G 0.06+0.25G 0 0.12+0.25G 0.12+0.12G 0.12+0.12G 0.12+0.12G 0.12+0.12G 0.12+0.25G 0.12+0.25G 0.12+0.25G 0.25+0.25%	82 88 86 76 80 87 88 0 87 89 90 88 86 89 92 0 94 0 54 7	6	8 11 8 4 10 6 13 0 11 11 14 13 15 11 13 0 11	91 95 95 87 90 94 95 0 96 98 97 93 94 97 95 0 99 99	4 5 5 2 6 5 3 0 5 3 6 5 3 5 5 0 3 6 0 3 9 6 7 7	4 110 3 7 3 9	57 1 8 7 7	0 56 3 7

## Summary

DPX-49674 at 0.12 oz/A gave similar control of a species regardless of adjuvants. In general species control with DPX-R9674 was or tended to be greater with MSC or AG-2 than X-77 or PO adjuvants.

2,4-D with adjuvants, Fargo 1988. 'McCall' soybean and redroot pigweed were seeded in 6 ft wide 'Wheaton' adjacent strips as bioassay species on May 18. Treatments were applied to 2 to 3 trifoliolate soybean, 6- to 8-leaf wheat on June 8 with 72 F and partly cloudy sky. Application was with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft wide area of 10 ft wide plots across the various species. The experiment was a randomized complete block with four replications. Evaluations were on June 28 and July 21.

	Westerna W	ere on Jun	e 28	and Ju	uly 2	21.	, ,	TOCK	With	four
<u>Treatment</u>	Rate		D	Jul	y 28				July 2	1
	(oz/A)		Rrpw	KOCZ		sf	Wht	Wht	Vosf	
2,4-Ddea 2,4-Ddea+MSC 2,4-Ddea+MSC 2,4-Ddea+PO 2,4-Ddea+PO 2,4-Ddea+AG-2 2,4-Ddea+AG-2 2,4-Ddea+X-77 2,4-Dbee 2,4-Dbee+MSC 2,4-Dbee+MSC 2,4-Dbee+PO 2,4-Dbee+AG-2 Brox-RP+BSC Brox-RP+BSC Brox-RP+BSC Brox-RP+AG-2 Brox-RP+AG-2 Brox-RP+AG-2 Brox-RP+AG-2 Brox-RP+AG-2 Brox-RP+AG-2 Brox-RP+AG-1 Brox-RP+AG-2 Brox-RP+AG-2 Brox-RP+AG-1 Brox-RP+AG-2 Brox-RP+AG-1	(oz/A)  3 6 3+0.12G 3+0.25G 3+0.12G 3+0.25G 3+0.12G 3+0.25% 3 6 3+0.12G 3+0.25G 3+0.12G 3+0.25G 3+0.12G 3+0.25G 3+0.12G 3+0.25G 3+0.12G 3+0.25G	3 3 5 3	48 85 79 86 82 84 83 88 75 89 96 86 65 89 88 89 88 45 45 45 45 45 45 45 45 45 45 45 45 45	31 72 69 87 73 75 86 73 78 91 78 85 55 77 86 89 73 41 69 41 63 68 68 65		(% c) 0 5 7 1	5 5 5 0 6 9 8 3 9 1 3 1	Wht 1) 4 6 7 10 8 2 2 7 1 4 9 9 9 9 9 9 9 9 9 9 9 9 9	Vosf  88 91 98 99 97 98 97 98 97 98 97 98 97 98 97 98 97 98 97 98 97 98 97 98 98 99 97 98 98 99 97 98 98 99 99 99 99 99 99 99 99 99 99 99	8rpw 91 95 94 92 93 88 96 92 96 99 98 88 88 88 88 88 88 88 88 88 88 88
		4		1	4	4	4	NS 4	17 4	

#### Summary

Species control with 2,4-D butoxyethyl ester (bee) was not enhanced by adjuvants. In general control of the species by 2,4-D diethanalomine (dea) was enhanced by all adjuvants and enhancement tended to be greater with 0.25G (1 qt/A) than 0.12G (1 pt/A) adjuvant. MSC and PO adjuvants at 0.25G with bromoxynil at 3 oz/A gave similar species control to bromoxynil alone at 6 oz/A. X-77 surfactant at 0.25% or oil adjuvants at 0.12G did not enhance

Sethoxydim with adjuvants and antagonism, Fargo 1988. 'Wheaton' Hard Red Spring wheat and 'ND810104' oats were seeded as bioassay species in adjacent 10 ft wide strips at Fargo, ND on April 28. Treatments were applied with a 10 ft wide type plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft wide bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft wide area across the bioassay species. Plots were 10 ft wide to provide a 4 ft area across the bioassay species. Plots were 10 ft wide to provide a 4 ft area between treated areas to aid evaluations. Treatments were untreated area between treated areas to aid evaluations. Treatments were applied to 5- to 6-leaf wheat and oats on June 7 with 98 F, 28% RH, and clear Sky. Evaluation was on June 24.

lear Sky. Evaluation was on C	- · à	Wheat	Oats
	Rate-	(% co	ntrol)
reatment	(oz/A)		
		64	63
	1+0.25G	77	79
ethoxydim+P0	1+0.25G+1G	0	0
athavydim+PU+60%	1+0.25G+16	8	9
- Thought mittletit	1+0.25G+16	8 4	6
**+PONNY I MTP()+Nd[ICO]	1+0.25G+16	15	22
- 1 L 0 V V A 1 M 1 P U + N A C C C C	1+0.25G+16+1G		78
LI	1+0.25G+16+1G	76	79
- II - WAIMIDITINATION	1+0.25G+16+1G	76	83
Sethoxydim+PO+Na2CO3+28%	1+0.256+10+14	82	
Sethoxydimino	1+0.25G	85	84
Sethoxydim+MS	1+0.25G+1G	13	18
Sethoxydim+MS+28%	1+0.25G+16	48	50
Sethoxydim+MS+Bent Sethoxydim+MS+Bent	1+0.25G+16	24	29
Cathorydiminithalicos	1+0.25G+16	78	75
C-TLOVIAIMIMIMITHIACUUS	1+0.25G+16+1G	81	84
a LL - MARMEM TENER LT COM	1+0.25G+16+1G	83	84
o TF ***********************************	1+0.25G+16+1G	79	83
Sethoxydim+M5+Nazcosizon	1+0.25G	59	65
Cathavyd1M+BLD	1+0.25G+1G	14	19
Cathavydim+BUT+20%	1+0.25G+16		19
Cathovydim+KI.H+DUIL	1+0.25G+16	18	2
Cathorydim+KLH+NdHCU3	1+0.25G+16	3	49
C TL SWINTER HTNACOUS	1+0.25G+16+1G	55	
a IIdimLRI H+DPIILTEU/	1+0.256+10+16	73	76
a II dam LRI H+NdIN,USI LOV	1+0.25G+16+1G	77	8:
Sethoxydim+BCH+Na2CO3+28%	1+0.25G+16+1G		
Sethoxydim+poiithaza		21	1
		14	1
C.V. %		19	1
LSD 5%		4	
LSD 1%			
# OF REPS			

a G in the rate column is for gallon per acre.

## Summary

Sethoxydim with PO was enhanced by 28%N, but 28%N antagonized grass control when sethoxydim was with BCH. Bentazon, NaHCO3, and Na2CO3 were highly antagonistic to grass control with sethoxydim regardless of oil adjuvant. The 28%N at 1 gpa only partly overcame bentazon antagonism of sethoxydim, However, MS with 28%N overcame more of the regardless of oil adjuvants. However, MS with 28%N. The antagonism bentazon antogonism of sethoxydim than other oils with 28%N. The antagonism of sethoxydim caused by NaHCO3 and Na2CO3 was overcome by 28%N with all of the oil adjuvants.

Fluazifop with adjuvants and antagonism, Prosper 1988. 'Wheaton' Hard Red Spring wheat, 'ND810104' oats, and Siberian foxtail millet were seeded as bioassay species in adjacent 6 ft wide strips at Prosper, ND on May 23. Treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft wide area across the bioassay species. Plots were 10 tuntreated area between treated areas to aid evaluation. Treatments were applied to 4- to 6-leaf wheat, oats, and foxtail 14 and August 9.

Treatment	Rate	Fom		14		August	9
	(oz/A)	FUII	i Wht	Oats	Fomi	lilh+	Oat
Fluazifop+PO				-(% cor	itrol)		
Fluazifop+PO+28% Fluazifop+PO+Bent Fluazifop+PO+NaHCO3 Fluazifop+PO+Bent+28% Fluazifop+PO+Bent+28% Fluazifop+PO+NaHCO3+28% Fluazifop+PO+Na2CO3+28% Fluazifop+MS Fluazifop+MS Fluazifop+MS+Bent Fluazifop+MS+NaHCO3 Fluazifop+MS+NaHCO3 Fluazifop+MS+NaHCO3+28% Fluazifop+MS+NaCO3+28% Fluazifop+BCH+SHACO3+28% Fluazifop+BCH+SHACO3+28% Fluazifop+BCH+Bent Fluazifop+BCH+Bent Fluazifop+BCH+NaHCO3 Fluazifop+BCH+NaHCO3 Fluazifop+BCH+NaHCO3 Fluazifop+BCH+NaCO3 Fluazifop+BCH+NaCO3+28% FluaZifop+BCH-NaCO3+28% FluaZifop+BCH	1+0.25G 1+0.25G+1G 1+0.25G+16 1+0.25G+16 1+0.25G+16+1G 1+0.25G+16+1G 1+0.25G+16+1G 1+0.25G+16 1+0.25G+16 1+0.25G+16 1+0.25G+16 1+0.25G+16+1G 1+0.25G+16+1G 1+0.25G+16+1G 1+0.25G+16 1+0.25G+16 1+0.25G+16 1+0.25G+16 1+0.25G+16 1+0.25G+16 1+0.25G+16	45 48 34 43 9 37 40 11 37 38 50 51 12 52 47 6 31 32	10	67 74 53 70 73 61 80 80 70 70 55 55 79 57 77 65 84 77 81 81 81 84 56 81 76	30 26 5 26 26 26 28 5 16 26 8 23 24 28 36 7 26 33 0 25 24 28 31 26 31 26 31 31 31 31 31 31 31 31 31 31 31 31 31	94 96 72 95 97 91 95 98 78 91 98 81 97 90 96 99 83 96 99 83 94 80 3	76 79 78 79 78 79 78 89 74 91 79 89 79 79 79 79 79 79 79 79 79

## Summary

PO (petroleum oil 11N with 17% Atplus 300F), MS (methylated seed oil, Sun-it), and BCH (adjuvant from BASF, Dash) with fluazifop all gave similar grass grass control. 28%N with fluazifop and the oil adjuvants did not influence Na2CO3 alone with the oil adjuvants or with 28%N did not influence grass control with fluazifop. Bentazon generally antagonized grass control from fluazifop. However, the antagonism was less evident for wheat and oats bentazon partly overcame the antagonism from bentazon, which was dependant on the species and the oil adjuvant.

Adjuvants with fluazifop, Fargo 1988. 'Wheaton' Hard Red Spring wheat, 1980 104' oats, and Siberian foxtail millet were seeded as bioassay species in adjacent 6 ft wide strips at Fargo, ND on May 31. Treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi applied area across the bioassay species. Plots were 10 ft wide to an 8 ft wide area across the bioassay species. Plots were 10 ft wide to provide a 4 ft untreated area between treated areas to aid evaluation. Treatments were applied early jointing wheat, oats, and foxtail millet on July 7 with 103 F, 30% RH and partly cloudy sky. Evaluations were on July 29 and August 23.

29 and August 23.			uly 2	9	Au		23
Treatment	Rate (oz/A)	Fomi	Wht	Oats	Fomi trol)-	Wht	<u> 0ats</u>
Fluazifop+PO Fluazifop+BCH Fluazifop+MSC Fluazifop+Inhance Fluazifop+LI-700 Fluazifop+X-77 Fluazifop+AG-2 Fluazifop+SCI-40 Fluazifop+SOTM  C.V. % LSD 5% LSD 1% # OF REPS	1+0.25G 1+0.25G 1+0.25G 1+0.25G 1+0.25G 1+0.25G 1+0.25G 1+0.25G 1+0.25G	10 11 8 5 9 6 13 1 9 62 NS NS	92 91 92 90 86 83 91 91 89 4 5 NS 4	69 62 77 60 73 39 77 38 44 18 15 21 4	16 11 9 6 10 8 13 5 2 46 6 8 4	99 99 99 98 98 99 99 98 NS NS 4	53 72 78 48 51 26 82 31 31 31 24 33 4

## Summary

The control of foxtail millet was too low and of wheat too high for determination of differences among adjuvants. Fluazifop gave greater oats control when applied with MSC (methylated seed oil from Agsco), AG-2, and BCH (adjuvant from BASF) than with any other adjuvant at the final evaluation (August 23).

Adjuvants with sethoxydim, Farqo 1988. 'Wheaton' Hard Red Spring wheat and 'ND810104' oats were seeded as bioassay species in adjacent 10 ft wide strips at Fargo, ND on April 28. Treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft wide area across the bioassay species. Plots were 10 ft wide to provide a 4 ft applied to 5- to 6-leaf wheat and oats on June 20 with 84 F, 50% RH, and clear sky. Evaluation was on June 28.

- a amoli c	Rate		
Ireatment  Sethoxydim+PO Sethoxydim+BCH Sethoxydim+MSC Sethoxydim+LI-700 Sethoxydim+X-77 Sethoxydim+AG-2 Sethoxydim+SCI-40 Sethoxydim+SOTM  C.V. % LSD 5%	Rate (oz/A) 1+0.25G 1+0.25G 1+0.25G 1+0.25G 1+0.25% 1+0.25G 1+0.25G 1+0.25G	78 84 81 66 56 58 77 57	0ats ontrol 77 83 80 64 44 53 76 57 67
LSD 1% # OF REPS		8	11 11
		11	14

Wheat and oats control with sethoxydim was greater with BCH (adjuvant from BASF, Dash) and MSC (methylated seed oil from Agsco) adjuvant than the other adjuvants. PO (petroleum oil 11N with 17% Atplus 300F) and AG-2 were similar, but more effective than SOTM (soybean oil with 5.5% T-Mulz VO), L1-700, X-77, or SCI-40 as adjuvants with sethoxydim.

Glyphosate with adjuvants, Prosper 1988. - 102Wheaton' Hard Red Spring wheat, "ND810104' oats, and Siberian foxtail millet were seeded as bioassay species in adjacent 6 ft wide strips at Prosper, ND on May 23. Treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft wide area across the bioassay species. Plots were 10 ft wide to provide a 4 ft untreated area between treated areas to aid evaluation. Treatments were applied 5- to 6-leaf wheat, oats and foxtail millet on June 30 with 80 F, 60% RH, and partly cloudy sky. Distilled water was the carrier. Evaluations were on July 8 and August 4.

valuations were on July 8			July 8			August 4 Fomi Wht Oat		
	D-1-a	Fomi	Wht	Oats	Fomi		Uals	
reatment	Rate			(% cont	ro1)-			
reacment	(oz/A)					92	96	
		93	86	81	99		99	
lyphosate(DH20)	3	92	88	86	97	97	99	
1bacatallH/UlTA-//	3+0.25%	90	87	84	97	98	99	
Typhosate(DH20)+LI-700	3+0.25%	91	92	88	97	99	99	
Typhosate (DH20)+28%	3+1G	91	90	84	99	98	99	
Glyphosate(DH20)+AMS	3+24	79	85	82	97	96		
Glyphosate(DH2O)+AG-2	3+0.25G	85	79	77	97	96	99	
Typhosate (DH20)+MSC	3+0.25G	82	83	81	96	96	98	
Glyphosate(DH20)+MSC	3+0.25G	80	80	79	95	93	98	
Glyphosate(DH20)+AG-2-C1	3+0.25G		88	86	96	98	99	
Glyphosate(DH20)+AG-2-C2 Glyphosate(DH20)+MSC-C1	3+0.25G	87	69	75	92	83	90	
CTURROCATOLUNIOUTINOU VA	3+0.25G	62	59	57	97	20	68	
civnhosate(UHZU) Thise of	3+16	63		56	96	47	76	
CJURNACATETNAHLUS	3+16+0.25%	61	69	65	93	63	66	
orbocototNanLUSTA //	3+16+0.25%	70	70	91	98			
Clumbacate+Nancote1	3+16+1G	92	95	88	98			
Clumbacata+Nancustcon	3+16+24	90	90	71	94			
CIURPACATATINAHUUSTAIIS	3+16+0.25G	76	71		96			
Clumbacate+Nancotau -	3+16+0.25G	74		66	95			
AT LACATALNAMI.UATIIAU		83			90			
AT LACATAINAMI.UATAU L		79			93	400	•	
Olimbocatornamination o		68						
OI bocotoINAHI.USTINO		79			88		3 7	
cl vnhosate+Nahcus+nsc oz		7:	72		8	_	7 8	
Clushacat Q+N3/LU3	3+16 3+16+0.25%	6!	72			-	54 7	
C1.mbocata+N3/LUSTA*//		7	1 73				99 9	
Clumbacate+NaZCUOTLI / V	3+16+0.25%	9	6 96				97 9	
Clyphosate+NaZUU3+ZU/		9	0 91	3 86			_	
Clumbacate+NaZLUSTANS	3+16+24	8	0 7	6 72	•			
Clunhosate+NaZUU3+Au-2	3+16+0.25G		4 6					
C1.unhacatatN3/LU3tN3/	3+16+0.25G		1 6	4 58				
01boca+01N3/1.U3+AU-6	C1 3+16+0.25G			5 5			0 2	
CTunhacatatNA/LUSTAU-L	02 01.20			3 5	-		54	
Ciunhacata+Na/LUSTISC V	- 050			9 6	5	94	62	
Glyphosate+Na2CO3+MSC-C	2 3+16+0.25G							
Gryphosacethazoostilos			12	10 1	5	4	20	
0 11 0/					5	5	20	
C.V. %				4	4	4	4	
LSD 5% # OF REPS			4					


or Na2CO3 in the spray carrier reduced grass species control from Adjuvants with glyphosate in distilled water (DH20) did not influence grass control at the August 4 evaluation. At the July 8 evaluation none of the adjuvants enhanced grass control, but AG-2 and MSC containing treatments tended to reduce grass control compared to that with glyphosate Ammonium sulfate (AMS) and 28%N in the spray overcame the antagonism for control of all grass species from glyphosate caused by NaHCO3 and Na2CO3.

Paraquat with adjuvants, Prosper 1988. 'Wheaton' Hard Red Spring wheat, 'ND810104' oats, and Siberian foxtail millet were seeded as bioassay species applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi to an 8 ft wide area across the bioassay species. Plots were 10 ft wide to provide a 4 ft untreated area between treated areas to aid evaluation. I with 73 F. and cloudy sky. Evaluation was on July 13 and August 9.

<u>Treatment</u>	Rat	P			y 13		Διιο	Tuct C
	(oz/	A)	Fc	omi W	nt Oa	ts F	omi W	lust 9 Iht Oa
Paraquat	( - 7 )				(%	cont	rol)	Iht Oa
Paraquat+X-77	5		0					
Paraquat+LI-700	5+0.25	5%		4 75			31	3 4
Paraquat+28%	5+0.25	%	8				16 3	
Paraquat+AMS	5+1G		8		94		33 56	
Paraquat+AG-2	5+24		8		71	2	20	
Paraquat+MSC	5+0.25	G	80		75		28 9	_
Paraguating	5+0.25	Ğ	76		96		6 41	_
Paraquat+AG-2-C1	5+0.25	G	75		84		1 18	
Paraquat+AG-2-C2	5+0.250		76		94		8 32	
Paraquat+MSC-C1	5+0.250		73		89		3 30	
Paraquat+MSC-C2	5+0.250		76		90		9 26	• •
Paraquat+NaHCO3	5+16		77	80	91		_	74
Paraquat+NaHCO3+X-77		E0/	81	78	76	38	_	95
araquat+NaHCO3111 700	5+16+0.2	3% E0/	78	80	80	19	_	13
ar aquat+NaH(1)3+28%	5+16+1G	3%	79	81	82	22		45
araquat+NaHCO3+VWc	5+16+24		79	77	79	28		14
araquat+NaHCO3+AC-2	5+16+0.2		81	76	77	36	•	13
al adnat + Nahlus I wec	5+16+0.2	06 -0	74	76	76	15		6
araquat+NaHCO3+AC 2 C.		0 G	74	77	75			4
an adrianthyahilistve o co		) G	77	76	74	11	7	8
ar adrig (+N9HCU3TMcC C1		96	70	73	71	12		11
ar adrighthancourt co	5+16+0.25	G	72	76	69	6	6	4
vi aquat+Na2CO3	5+16+0.25	G	73	74	63	6	6	4 3 9
raquat+Na2cn3_v 77	5+16	•	78	80	80	10	8	9
Indquat+Na2CO3+11 700	5+16+0.25	%	70	81	84	21	37	45
1 addgf+N32LU3T30%	5+16+0.25	<b>%</b> .	79	81	82	6	20	46
raquat+Na2CO3+AMc	5+16+1G		78	78	80	23	34	43
raquat+Na2CO3+AC 2	5+16+24		81	78	77	28	11	6
Ladnat+Na2CU3 Mcc	5+16+0.250	i	75	80	76	26	6	9
raquat+Na2CO3_AC 2 C1	5+16+0.250		69	78	75	16	9	5
addd + Nazi lisi ve a ca	5+16+0.256		77	79	72	9	25	5
addarthys/1037Mcc C1	5+16+0.25G		71	80		13	14	6
	5+16+0.25G		71	76	68	12	11	6 3
	5+16+0.25G		76	77	66	10	5	0
. %			, ,	11	71	16	18	9
5%			5	2	7	1 4 1		
F REPS			6	3	7	44	51	43
			4	4	7	10	13	18
			7	4	4	4	4	4

# Summary

NaHCO3 or Na2CO3 in the paraquat spray did not influence grass species control. Grass species control with paraquat was not greatly influenced by adjuvants.



Fluroxypyr alone and with auxin herbicides for leafy spurge control. Lym, Rodney G., and Calvin G. Messersmith. Fluroxypyr is a picolinic acid herbicide similar to picloram but with less soil residual and a different weed control spectrum. The purpose of this experiment was to evaluate fluroxypyr for leafy spurge control when applied once, when applied with auxin herbicides, and when applied in a repetitive treatment program.

The experiment was established and original herbicide treatments were applied in a dense stand of leafy spurge near Dickinson, ND, on July 14, 1986. Previous research had indicated the optimum application time for leafy spurge control with fluroxypyr was post seed-set rather than during true flower as picloram. The herbicides were applied using a tractor-mounted sprayer delivering 8.5 gpa at 35 psi. The retreatments were applied as a split-block treatment with three replications. The original whole plots were 15 by 56 ft, and the retreatment subplots were 10 by 15 ft. Retreatments were applied in mid-July 1987 and 1988. Evaluations were based on visible percent stand

Fluroxypyr at 0.5 and 1 1b/A provided an average of 90 and 41% leafy spurge control 2 and 11 months after treatment (MAT), respectively (data not shown). Control was similar when fluroxypyr at 0.25 or 0.5 1b/A was applied

No treatment provided satisfactory leafy spurge control in June 1988, 11 months following the first retreatment (Table). Picloram at 1 lb/A provided the best leafy spurge control of the original treatments (44%) when averaged over retreatments. Picloram at 0.5 lb/A or fluroxypyr plus picloram at 0.5 + 0.25 1b/A were the best retreatments, but control averaged 41 and 40%, respectively, when averaged over original treatments.

Control increased to an average of 95% with all retreatments except picloram alone at 0.25 and 0.5 lb/A following the second application in July 1988 (Table 1). Leafy spurge was under severe drought stress and generally was 12 inches or less tall at the time of retreatment. Picloram alone appeared to be absorbed poorly as many plants showed little visible injury 2 MAT. Although fluroxypyr alone or fluroxypyr plus dicamba, picloram, or 2,4-D generally provided similar or less leafy spurge control than picloram or picloram plus 2,4-D in 1987, fluroxypyr alone was much better than picloram alone under dry conditions in 1988. Fluroxypyr at 0.5 lb/A averaged 95% control as a retreatment compared to 50 and 70% with picloram at 0.25 or 0.5 1b/A, respectively. Fluroxypyr may be useful in a retreatment program, especially in areas where picloram cannot be used or in late-season treatments during dry conditions. (Published with approval of the Agric. Exp. Stn., North Dakota State Univ., Fargo 58105).

Table. Leafy spurge control with fluroxypyr alone and in combination with auxin herbicides (Lym and Messersmith).

				Retr	eatment/r	ate (lb/A			
	all the Steel			ı	luro.	Fluro.	PIC.T	0	
			Pic. F		. Dic	+ Pic.	2,4-D	Con-	Maan
riginal		Fluro.	0.05	5 0	25+0.25	0.5+0.25	0.25+1	trol	Mean
reatment	Rate	0.5	0.25		(% contro	June 198	8)		
reatment	(lb/A)				(10				
			Tank i	20	43	34	10	3	25
Emilia vissi ila	0.5	30	28	29	34	46	23	9	30
Luroxypyr	1	26	28	42		49	25	2	29
Fluroxypyr	0.25 + 0.25	25	17	54	33	42	16	3	26
Fluroxypyr + pictoram	0.5 + 0.25		32	39	35	37	42	0	27
Fluroxypyr + pictoram	0.5 + 1	42	12	34	22	53	39	15	30
Fluroxypyr + 2,4-D	0.25 + 0.25	22	25	40	17	41	27	13	25
Fluroxypyr + dicamba	0.25 + 1	20	17	42	16		65	30	44
Picloram + 2,4-D	1	40	35	43	45	47	18	0	22
Picloram		29	9	45	32	12	29	10	
Control		27	23	41	31	40	27		
LSD (0.05)	wiic	ic pro-				whole pl			
dune take and			, , , , , , , , , , , , , , , , , , , ,				96	8	73
	0.5	98	63	76		77	97	32	
Fluroxypyr	9	95	50	72	92	98	91	3	
Fluroxypyr				64	97	97	98		
Fluroxypyr + piclora		25 90		69	96	97	94	100	
Fluroxypyr + pictors	0.5 + 1	9	4 33	65	93	98	97		
Fluroxypyr + 2,4-D	0.5 4 1			7	99	98			0 70
	a 0.25 + 0 0.25 + 1		8 68	3 70	6 95	79	96		1 7
Fluroxypyr + dicamb	0 75 4 1		5 40		6 96	95	92	•	0 7
Fluroxypyr + dicamb					98	99	98	,	,
Fluroxypyr + dicamb	1			7 7	3 40		_		0
Fluroxypyr + dicamb		9	9 3		7 96	93	9	6 1	0
Fluroxypyr + dicamb Picloram + 2,4-D Picloram	1	9	9 3° 95 5	0 7	0 %	93 ; whole		4744	

<sup>&</sup>lt;sup>a</sup>Evaluated prior to second retreatment in July, 1988.

Evaluation of sulfometuron applied in mid-summer and fall for leafy spurge control. Lym, Rodney G., and Calvin G. Messersmith. Previous research at North Dakota State University has shown that sulfometuron delays, and sometimes stops, bud growth on leafy spurge roots. A herbicide that spurge reestablishes by growth should improve long-term control, since leafy control. The purpose of these experiments was to evaluate sulfometuron alone and in combination with auxin herbicides applied from mid-July to mid-September for leafy spurge control.

All herbicides were applied with a tractor-mounted sprayer delivering 8.5 gpa at 35 psi. All plots were 10 by 30 ft in a randomized complete block design. The sulfometuron experiment establishment dates in 1986 and leafy mature growth stages were: July 22 and August 27 near Chaffee, ND, at the City, ND, well branched and in the fall regrowth stage; and September 3 near Valley Dickinson, ND, in the fall regrowth stage with most leaves chlorotic or bright was applied 12 months after the original treatment of picloram at 4 oz/A to the back one-third of each plot at Chaffee and Dickinson and at 8 oz/A at compared to the control.

Sulfometuron plus auxin herbicide treatments applied in July near Chaffee provided 82 to 100% top growth control 1 month after treatment (MAT) (Table 1). Sulfometuron alone did not provide satisfactory leafy spurge control. When evaluated in May 1987, grass injury tended to increase as the picloram or dicamba compared to sulfometuron alone. When evaluated in August 1987, control was similar whether sulfometuron was applied alone or with an auxin herbicide prior to the picloram retreatment (62%). Control decreased rapidly and no treatment provided satisfactory leafy spurge control in 1988.

Leafy spurge control tended to be better when sulfometuron plus an auxin herbicide was applied in August or September (Table 2) compared to July control tended to be higher as the sulfometuron rate increased up to 2 oz/A. The dicamba and 2,4-D rate had little effect on control over the ranges increased. Long-term control was much higher at Valley City compared to the other two locations. The best treatment for long-term control at Valley City compared to 32% control with picloram at 2 + 16 oz/A which averaged 80% 22 MAT picloram at 4 or 8 oz/A increased leafy spurge control at Chaffee and Valley Original sulfometuron treatment rate. (Published with approval of the Agric. Stn., North Dakota State Univ., Fargo 58105)

Table 1. Leafy spurge control by sulfometuron plus auxin herbicides applied in July at Chaffee, ND (Lym

and Messersmith).		Aug 86 Con-	Con-	87 Grass	Aug Con- Re	87	May 8	treat-	Aug 8 Con- Re trol m	treat
eatment	Rate (oz/A)	trol	trol			(%)		4/	0	10
ulfometuron + picloram ulfometuron + dicamba ulfometuron + 2,4-D sulfometuron + picloram sulfometuron + dicamba sulfometuron + dicamba Sulfometuron + dicamba	0.5 + 8 0.5 + 16 1 + 8 1 + 8 1 + 16 2 + 32 2 + 128	31	40 5 18 60 47 97 96 18	11 0 3 20 11 30 49 10	15 7 8 16 14 60 59 7 8	52 54 53 54 76 66 69 66 72	6 10 10 10 4 53 26 6 0	16 16 43 27 28 65 37 41 33 26	7 1 6 0 38 11 1 3	6 19 13 6 35 15 9 19
Sulfometuron Control LSD(0.05)	0	13 0 15	0	0 21	0 22	48 NS	0 NS	NS	NS	24

a Picloram at 4 oz/A applied as a split-block treatment to the back one-third of each plot on June 29, 1987.

Table 2. Sulfometuron plus auxin herbicides applied in August or September for leafy spurge control (Lym and Messersmith).

		_	M- 07		Evaluation	on date		
Treatment		Con	May 87		Aug 87		une 88	
Treatment	Rate			Con	- Grass	Con-		Sept 88
	(OZ/A)	tro	linjury	tro	l injury	trol		Retreat.
Chass						-(%)	ment	ment
Chaffee						(7)		
Sulfometuron + picloran	0.5 + 8							
Suttometuron + dicamba	0.5 + 16	89	35	15				
Sulfometuron + 2 (-n		68	8	16	• •	5	78	11
Sultometuron + niclorem		35	83	1	• •	13	72	10
Sucrometuron + dicamba		95	46	32	••	0	44	11
Sulfometuron + niclorem	1 + 16	81	36	17	• •	8	67	16
Sulfometuron + dicamba		94	56		• •	5	78	11
Fosamine	2 + 128	95	53	70	• •	29	68	
Fosamine	64	43	15	56		8	78	12
Control	96	56		9		3	78	16
COLLLOG	• •		13	20		6		16
		0	0	0			70	12
LSD (0.05)					••	0	63	10
		29	19	28		1.0		
Dickinson						NS	NS	NS
Sulfometuron + 2,4-D	0.5							
Sulfometuron + picloram	0.5 + 16	55	61	23				
Sulfometuron + 2,4-D	0.5 + 12	97	71		33	0	3	
Sulfometuron + 2,4-D	2 + 16	75	73	67	26	1	25	• 0
Sul fomotion	2 + 32	78	70	26	33	1	16	
Sulfometuron + picloram	2 + 8	95		29	33	4	14	• •
Sulfometuron + pictoram	2 + 12		89	83	60	11		
Sulfometuron + picloram	2 + 16	99	94	90	80	8	14	
	- 10	99	98	93	91		36	
LSD (0.05)						20	39	
		20	29	22	24			
Malley City					E4	NS	NS	
ulfometuron + 2 4-n	0.5							
ulfometuron + 2,4-D	0.5 + 16	41	0	44				
ulfometuron + picloram	0.5 + 32	57	0	11	0	6	96	20
ulfometuron + picloram	0.5 + 8	96	7	9	0	1	91	
ul fometusen	0.5 + 12	98	3	39	0	3	98	19
ulfometuron + picloram	0.5 + 16	99		68	0	15	99	43
ulfometuron + 2,4-D	1 + 16	90	4	81	0	16	99	36
ulfometuron + 2,4-D	1 + 32		5	26	0	5		51
ulfometuron + picloram	1 + 8	93	6	41	0	8	94	29
it i ometuron + nichana	1 + 12	99	8	85	0		99	34
rometuron + niclorem		99	6	88	Ö	36	97	37
it tometuron + 2 4-h	1 + 16	99	8	86	0	34	96	53
Ittometuron + 2 4-n	2 + 16	97	34	68		45	99	43
lfometuron + picloram	2 + 32	99	29	73	4	10	99	57
lfometuron + picloram	2 + 8	99	49		14	13	98	
lfometuron + pictoram	2 + 12	99	41	97	20	52	100	52
cloram	2 + 16	99	77	95	0		100	68
ntrol	16	99		98	20	80	99	75
ici ot				63	0	32	97	65
150 (0.05)		• •	• •	• •	••	0		25
LSD (0.05)		12					98	29
		12	22	22	20	22	7	

Picloram at 4 oz/A applied as a split-block treatment to the back one-third of each plot in Aug 1987 at Chaffee and Dickinson and at 8 oz/A at Valley City.

Sulfometuron applied alone and with auxin herbicides as an annual treatment for leafy spurge control. Lym, Rodney G., and Calvin G. Messersmith. Sulfometuron is an analog of chlorsulfuron but with slightly less soil residual and a different weed control spectrum. Sulfometuron currently is used for grass suppression along roadsides and also has currently is used for grass suppression along roadsides and also has controlled some broadleaf weeds including leafy spurge. The purpose of these controlled some broadleaf weeds including leafy spurge. The purpose of these experiments was to evaluate sulfometuron alone and in combination with auxin herbicides for leafy spurge control.

The first experiment was established in cropland severely infested with leafy spurge near Hunter, ND. Spring and fall treatments were applied on June 27 and September 4, 1985, respectively. Leafy spurge was 26 to 36 June 27 and September 4, 1985, respectively. Leafy spurge was 26 to 36 June 27 and September 4, 1985, respectively. Leafy spurge was 26 to 36 June 27 and September 4, 1985, respectively. Leafy spurge was 26 to 36 June 27 and September 5. The summer dormancy had begun when treatments were applied in September. The summer dormancy had begun when treatments were applied in September. The summer dormancy had begun when treatments were applied in September. The summer dormance applied with a tractor-mounted sprayer delivering 8.5 gpa at herbicides were applied with a tractor-mounted sprayer delivering 8.5 gpa at herbicides were applied with a tractor-mounted sprayer delivering 8.5 gpa at herbicides were applied with a tractor-mounted sprayer delivering 8.5 gpa at herbicides were applied in August 1987, as a split-block picloram at 4 oz/A was applied in August 1986 and 1987, as a split-block picloram at 4 oz/A was applied to the front one pretreatment to picloram. Picloram at 8 oz/A was applied to the front one pretreatment to picloram. Picloram at 8 oz/A was applied to the front one stand reduction as compared to the control.

Leafy spurge growth stopped following application of sulfometuron alone, regardless of application date. Plants treated with sulfometuron alone in June had slightly chlorotic leaves during the growing season but root bud elongation was inhibited when evaluated in August. Leafy spurge top growth was killed when treated with sulfometuron plus an auxin herbicide, and root bud growth was inhibited. Leafy spurge root buds were white and short on bud growth was inhibited. Leafy spurge root buds were white and short on plants treated with sulfometuron, compared to the pink elongated buds on plants treated plants. Sulfometuron plus an auxin herbicide provided better leafy untreated plants. Sulfometuron alone, and long-term control was better when spurge control than sulfometuron alone, and long-term control was better when sulfometuron was mixed with picloram than with 2,4-D or dicamba (Table 1).

Leafy spurge control increased following retreatment with picloram at 4 oz/A, sometimes to 100% control, (Table 1). In August 1987, control averaged 81 and 67% when picloram was applied to plants originally treated with 81 sulfometuron in the spring and fall, respectively. Control increased following the picloram retreatment as the sulfometuron rate increased following spring but not fall treatments. Control averaged 82 and 75% for following spring but not fall treatments. Control averaged 82 and 75% for following and fall treatments, respectively, in August 1988 12 months the spring and fall treatment of picloram at 4 oz/A. Control was similar following the second retreatment of picloram at 4 oz/A. Control was similar among sulfometuron treatments applied in 1985 and retreated with picloram in 1986 and 1987 and picloram retreatments alone.

An experiment to evaluate leafy spurge control from sulfometuron plus auxin herbicides applied annually spring or fall was established on very mature leafy spurge stands near Chaffee and Dickinson, ND. Treatments were applied as described previously in June or late-August to early September in 1986, 1987, and 1988. Leafy spurge was in the true flower or fall regrowth stages when the original treatments were applied but was in the vegetative stages when the treatments were reapplied the following growing seasons.

Leafy spurge control averaged 36 and 68% at Chaffee and Dickinson, respectively, following three spring applications of sulfometuron plus an auxin herbicide (Table 2). Sulfometuron at 1 or 2 oz/A alone resulted in control after three applications averaged over both locations. Sulfometuron and there was little grass injury.

Leafy spurge control was higher when annual sulfometuron treatments were applied in the fall compared to the spring (Table 2). Control in the spring sulfometuron plus dicamba two applications of sulfometuron plus picloram or by fall at Dickinson and Chaffee, respectively. Sulfometuron alone or applied with 2,4-D generally provided poor leafy spurge control. Grass applications and averaged 87% The grass began to recover at Chaffee and Dickinson because severe drought conditions prevented grass reestablishment.

Sulfometuron applied with picloram or dicamba in the fall has provided better leafy spurge control than either auxin herbicide applied alone at similar rates. Grass was injured severely when sulfometuron was fall applied but did recover by the following growing season. Sulfometuron in combination with an auxin herbicide could be used in a leafy spurge management program to provide initial control but probably should not be applied annually because for grass injury. (Published with approval of the Agric. Exp. Stn., North Dakota State Univ., Fargo 58105)

Table 1. Leafy spurge control with sulfometuron applied either alone or with various auxin herbicides near Hunter,

ND (Lym and Messersm								E	valuation	on date			August	88
											May 88		Picloram	retreat
								A. m. sc P	1987	P	icloram r	0	- 4	8 oz/A
		Aug	May	Aug		1ay 19	987 a	August Single R	8	Single 4	oz/A 8	OZ/A	4 0Z/A	0 02,
oplication date/		4005	1086	1986	sin	gle R	etreat	Single R	201600					
			1700					(	& contro	,()				
1 eacherre	(oz/A)									0	98	97	76	49
une 27, 1985				0		0	87	5	63		97	93	80	71
	1	0	6			12	88	17	85	20	98	87	73	54
ulfometuron	1.5	0	63	25		3	87	10	82	3	94	92	77	54
ulfometuron	2	0	36	6			84	24	64	18		95	78	63
iul fometuron	1 + 10	5 95	76			8	98	55	86	33	99	94	88	71
Sulfometuron + 2,4-D	1 + 3	2 96	85	41	•	35		67	94	33	100		94	54
Sulfometuron + dicamba	1 + 8		96	, 5	9	51	100	5	93	5	99	77	89	61
Sulfometuron + picloram	2 + 0		60	) 2	4	0	98	0	55	4	97	93	07	
Sulfometuron + metsulfuron		(	. (	)	0	0	63	U						NS
Control	. •								31	25	NS	NS	NS	NJ
		2	5 2	2 2	26	25	31	20	31	gayasu.				
LSD (0.05)		2	, -											
200 (000)										0	99		54	0.0
September 4, 1985				, de	0	0	54	0	40		99		76	0.0
September 4, 1,200	0.5	,		16	0	23	77	21	56	1	99	0.0	77	0.0
sulfometuron	1			95	7	3	92	8	72	0			82	0 (
sulfometuron	9 +	16	• •	99	17		91	13	73	8	100	0.0	83	d •
Sulfometuron + 2,4-D	1 +	32		97	23	15	83	38	83	5	100	9 0	85	
Sulfometuron + dicamba	1 +			99	74	33		91	62	8	99		81	
sulfometuron + picloram	0.5 +		. •	95	24	21	87	48	84	2	100	, 0	64	
sulfometuron + 2,4-D	0.5			97	51	19	83	94	71	6	100	0 6		
sulfometuron + dicamba	0.5			99	40	17	86		62	0	99	•		
sulfameturon + picloram		+ 0.5		88	13	0	83		49	3	100			
Sulfometuron + metsulfuro				44	6	4	. 76			(		•	. 7	0
DPX-L5300		1	0 4	0	0	0	7	3 0	38					
Control		0 0	0.0	U						N:	e NS			IS
GOILET & C				26	30	34	6 2	9 32	NS.	N	3 10			
LSD (0.05)				26	20									

a Picloram at 4 oz/A applied as a split-block to the back one-third of each plot in August 1986 and 1987. b Picloram at 8 oz/A applied as a split-block to the front one-third of each plot in August 1987.

Table 2. Leafy spurge control with sulfometuron applied annually either alone and in combination with dicamba, picloram, and 2,4-D applied in the spring and fall at two locations in North Dakota (Lym and Messersmith).

				affee			Dickir	
Treatment and		May	1988	Aug	1988	June		
application date	Rate	Control	Grass injury		Grass injury		Grass	<u>Sept 1988</u>
	(OZ/A)					Control	injury	Control
Chaine					(%)			
Spring								
Sulfometuron	1	1	6					
Sulfometuron	2	4	43	0	0	0	25	32
Sulfometuron + picloram	1 + 8	39	11	3	0	5	21	17
Sulfometuron + dicamba	1 + 32	33		39	9	64	25	79
Sulfometuron + 2,4-D	1 + 16	15	26	28	14	27	16	67
		13	10	42	15	5	16	59
all								39
ulfometuron								
ulfometuron	1	27	70	0	5	49	88	4.
ulfometuron + picloram	2	69	96	0	12	55		15
11 60-04		99	93	73	51	98	92	9
111-	1 + 32	97	85	48	28	98	100	82
icloram	1 + 16	71	91	13	17		95	94
o cor call	32	99	41	78	11	66	56	27
						99	44	94
1.00								
LSD (0.05)		18	33	25	0.7			
				25	23	18	25	28

a Applied annually in 1986, 1987, and 1988.

Grass injury was not evaluated at Dickinson in Sept. 1988 due to severe drought conditions.

Leafy spurge control under trees and along waterways. Lym, Rodney G., Leafy spurge is difficult to control with herbicides near trees or open water such as ponds, ditches, and rivers because of potential damage to desirable vegetation or of water contamination. However, these areas provide a source of seed for infestation of nearby and downstream areas if no control measures are initiated. The purpose of these experiments was to evaluate several herbicides both for leafy spurge control and for potential to damage desirable vegetation.

Three experiments for leafy spurge control under trees were established in a shelter belt located in a waterfowl rest area near Valley City, ND. The plots were located in a dense stand of leafy spurge growing under mature ash and elm trees that had been planted 5 ft apart in 12-ft rows. The herbicides were applied either with a hand-held single-nozzle sprayer delivering 40 gpa or with the controlled droplet applicator (CDA) which applied approximately 4 gpa. The hand-held sprayer treatments were applied as a premeasured amount of herbicide: water per plot to assure the correct rate and three passes were made across each plot to assure adequate coverage. The CDA treatments covered each plot only once. The experiment starting dates and leafy spurge stage at treatment were: June 26, 1986, flowering and beginning seed set; September 3, 1986, post-seed set and chlorotic leaves; and June 16, 1987, yellow bract to flowering growth stage. Plots were 12 by 24 ft arranged in a randomized complete block design with four replications. Evaluations were based on visible percent stand reduction as compared to the control.

Initial leafy spurge control was poor when glyphosate was applied alone, regardless of rate or treatment date (Table 1). Control improved to over 90% 12 months after treatment (MAT) following a June but not September application. Grass injury was nearly 100% with all glyphosate treatments. Leafy spurge control declined to 50% or less by June 1988 but very little grass had reestablished.

Sulfometuron alone did not control leafy spurge, however, control was improved consistently when sulfometuron was applied with glyphosate regardless of rate or treatment date (Table 1). Leafy spurge control averaged 97% 12 MAT with sulfometuron plus glyphosate at 1 or 2 + 17 oz/A but in general declined rapidly the second year after treatment. Leafy spurge control with sulfometuron plus 2,4-D declined rapidly following the 12 month evaluation. Picloram, applied with the CDA at a picloram:water concentration of 1:7 (v/v), provided nearly 100% leafy spurge control with no grass injury. Control averaged 75% in August 1988 following application in June 1986 or 1987 but only 57% when applied in September. Several ash trees had some leaf curling but no visible permanent damage from this treatment.

The experiment to evaluate leafy spurge control with herbicides that can be used near water was established on June 27, 1986, along a ditchbank in Fargo. The experimental design and application methods were similar to the tree experiment. All plots were treated with 2,4-D at 1 lb/A in June 1987 to control leafy spurge seedlings.

Amitrole at 4 lb/A provided 91 and 95% leafy spurge control 12 and 15 MAT, respectively, but the treatment caused 64% grass injury (Table 2). Increasing the amitrole rate to 8 1b/A increased grass injury but not leafy spurge control which averaged 78% 27 MAT. Unfortunately, amitrole no longer is cleared for use near water. Fosamine at 8 lb/A provided 90 and 74% leafy spurge control 12 and 27 MAT, respectively. No other fosamine treatment provided satisfactory control, and evaluations varied considerably from plot to plot indicating this herbicide may provide inconsistent control. (Published with approval of the Agric. Exp. Stn., North Dakota State Univ., Fargo 58105)

Table 1. Leafy spurge control under trees (Lym and Messersmith).

Application date		Aug 86	May	87	Aug	87				
and treatment	Rate			Grass		Grass	_ June	88	Aug	88
	(oz/A)	Control	Control	injury	Contro	linjur	<u>Control</u>	Grass		Gras
June 26, 1986					••••••	(%)			Control	injur
Glyphosate										
Glyphosate	8.5	9	92	88	79					
Sulfometuron	17	41	96	98	94	• •	46	70	33	71
Sulfometuron	0.5	15	0	0	29	••	53	89	54	91
Sulfometuron	1	9	0	0		• •	4	0	26	
Sul fometuron + al	2	9	28	15	19	• •	0	0	14	0
Sulfometuron + glyphosate	0.5 + 8.5	13	98		19	• •	4	0	12	0
A GIVENER OF A GIVENER		13	96	98	90		58	63		10
Sulfometuron + glyphosate	2 + 8.5	24	99	99	95	• •	75	%	50	68
Pictoram (CDA)	1:78	99		96	85	• •	71	70	81	95
		,,	95	0	85		76		66	94
LSD (0.05)		19					,,,	0	79	0
		19	8	14	23		28			
September 3, 1986							20	31	27	24
Glyphosate	17									
Sulfometuron + glyphosate		• •	65	99	54					
Sulfometuron + 2,4-D	2 + 17	0 0	99	99	89	• •	22	98	10	94
ricloram (CDA)	2 + 17	• •	69	66	51	0 0	63	99	55	75
(357)	1:7ª		86	9		• •	6	29	1	
LSD (0.05)				,	66		67	0	57	25
(0.05)			26	17					91	0
une 16, 1987				17	31	• •	29	21	25	
									25	40
yphosate	8.5									
yphosate	17		••	• •	13	98	36	89	4.0	
Ilfometuron + glyphosate	0.5 + 8 5		• •	• •	30	98	76		18	99
TOMELUTON + glyphosate	1 + 8 5		•	• •	9	83	21	94	36 1	00
lfometuron + glyphosate	2 + 8.5	•	•		12	86		60	9	88
lfometuron + 2,4-D		•		. 3	36		51		31	96
lfometuron + 2,4-D	1 + 17 .		•		75			87	11	84
cloram (CDA)	2 + 17				9			40		23
	1:7ª				6			14		1
LSD (0.05)				. ,	0	0	80	0	94	0
				1:	2	25	18 2	20 1	6 2	3

a Solution concentration pictoram (Tordon 22K):water.

Table 2. L	eafy spu	rge contro	ol along (	di tchbanl	(s, fargo	, ND .		Aug 88	3
		Aug 86	May	878	Aug 87	June	Grass	- Augusta	Grass
				Grass	Control	Control	injury	Control	injury
Treatment	Rate	Control	Control	injury	(%				
	(lb/A)						8	71	11
	2	99	69	23	80	63	36	75	26
Amitrole	4	100	91	64	95	82 78	40	78	30
Amitrole Amitrole	8	100	87	81	96 59	54	5	39	4
Fosamine	2	5	14	3	55	59	3	60	3
Fosamine	4	19	58	10 57	82	74	18	73	13
Fosamine	8	40	90	51			23	27	NS
Len	(0.05)	19	17	42	28	27			
LSU	10.00								

<sup>&</sup>lt;sup>8</sup> 2,4-D at 1 lb/A applied in June 1987 to control seedlings.

Leafy spurge control with various herbicide combinations applied alone or with spray additives. Lym, Rodney G., and Calvin G. Messersmith. The relatively high cost of herbicides limits their use on pasture and rangeland for leafy spurge control, especially when applied at rates that are in excess that picloram plus 2,4-D at 0.25 + 1 lb/A applied annually is more cost-effective than high rates of herbicides applied once. Picloram or 2,4-D control. The purpose of this research was to evaluate various herbicide combinations applied alone or with spray additives for leafy spurge control.

Three experiments were established in a dense leafy spurge infestation growing along railroad tracks in West Fargo. The first two were established on June 9, 1987, when leafy spurge was in the true flower growth stage, and the third on September 1, during fall regrowth. A fourth experiment was established on June 16, 1987, near Valley City with leafy spurge in the seed set growth stage. A retreatment of picloram at 8 oz/A was applied in June 1988 as a split-block treatment to the back one-third of each plot at Valley 8.5 gpa at 35 psi. All plots were 10 by 30 ft in a randomized complete block to the control.

Picloram plus 2,4-D or picloram applied with methylated sunflower oil, with NH<sub>4</sub>SO<sub>4</sub> (liquid or granular formulations), or at pH 4.7 buffered with Though not evaluated in this experiment, leafy spurge control (Table 1). herbicide treatments applied without additives in adjacent experiments provided better initial and long-term control. Visual observations indicated plants recovered quickly and regrowth from the lower portion of the stems sulfometuron also was lower than expected and may be due to application during control if applied when relative humidity is high.

Sulfometuron applied with picloram, fluroxypyr, or various formulations of 2,4-D generally provided leafy spurge control similar to the same herbicides applied alone (Table 2). Subsequent research has shown sulfometuron increases leafy spurge control when applied with these herbicides in the fall but not in the spring, when this experiment was established. Leafy spurge control with sulfometuron plus fluroxypyr or picloram decreased as the fluroxypyr rate increased, but increased with increasing rates of picloram.

In August 1987, sulfometuron at 1 to 2 oz/A applied with picloram at 8 oz/A or 2,4-D at 16 oz/A provided an average of 94% leafy spurge control but only 61% when applied with dicamba at 16 oz/A (Table 3). The average control at Valley City was much higher than with similar treatments applied at West Fargo (Table 2). The increase could be due to application at a later growth during treatment. It was 90 F with 50% relative humidity at Valley City The soil pH and organic matter was similar at the two locations.

Dicamba plus sulfometuron at 16 + 2 oz/A and all sulfometuron plus picloram treatments provided good leafy spurge control 12 months after treatment (Table 3). Control declined rapidly, thereafter, and no treatment provided satisfactory control by August 1988 even with a picloram retreatment. In general, no herbicide combination treatment or additive provided longer-term or more cost-effective leafy spurge control than picloram plus 2,4-D. (Published with the approval of the Agric. Exp. Stn., North Dakota State Univ., Fargo, 58105)

Table 1. Leafy spurge control with various herbicides and additives spring or fall applied at West Fargo, ND (Lym and Messersmith).

fall applied at West Fargo, NI	(Lym and me		
Tall applies	1007 Amplic	ation/eval	uation date
	1987 Applic	actony of	September 1
	June 9		May 88
no+o	Aug 87	May 88	May oo
Rate (A)		(%)	,
Treatment (oz/A)			Tax ma star
	27	6	22
$a + b + MSW^a + 4 + 16 + 16$	16 27	8	23
Picloram + 2,4-D + $MSWa$ 4 + 16 + 8 + 16 +	16 15		• •
Picloram + 2,4-D + MSWa	16 33	0	34
Picloram + MSWa 8 +	16 25	6	
PICTOTAIII MSWa	10	18	• •
Picloram + MSW <sup>a</sup> Picloram + MSW <sup>a</sup> 8 + 1 +	16 4/	14	
Picloram + Surrolle at 5 64		40	
Fosamme	13	13	• •
Farming Ct . 1	6		• •
Framino + cilitometurui		0	36
Focamine + SUITOMELUION 20 . 1		13	
Fosamine + Sulfometaron 32 + 1 Dicamba + MSW 4 + 16 +	O	• •	15
Dicamba + MSW 4 + 16 +	0.10	e 0	21
Picloram + 2,4-0 + NH4504b 8 + 16 +	+ 0.13		4
Picloram + $2,4-0$ + $10,40$ 4 + $16$	+ 40	• •	23
Picloram + $2,4-0$ + NH4504c 8 + 16	+ 40	0 0	11
Picloram + 2,4-D + NH4504 acid 4 +	16	0 0	10
Dicloram + 2.4-D + citric actual 4	16		10
Pictoram + 2,4-D + citric acid 8 +	10		of Figure 1980 S. Fr
$\begin{array}{llllllllllllllllllllllllllllllllllll$	20	15	NS
	20		The last state of the state of
LSD (0.05)	sponto 4K-315	o omulcifi	er.
	conto 4K-315	R GIIII 12111	

a MSW = Methylated sunflower oil plus sponto 4K-3158 emulsifier.

<sup>8-0-0-9</sup> liquid fertilizer equivalent to N and S at 0.2 lb/A respectively.

Equivalent to 2.5 lb N/A. Spray solution buffered to pH 4.7.

Table 2. Leafy spurge control with sulfometuron plus various auxin herbicides at West Fargo, ND (Lym and Messersmith).

<u>Treatment</u>		Fvalu	24:
	Rate	9 Aug 8	ation date <sup>a</sup>
Picloram + sulfometuron	(oz/A)	- nay o	7 24 May 88
Picloram + sulfometuron			(%)
Picloram + sulfometuron	4 + 1		
Picloram + sulfometuron	8 + 1	6	13
	12 + 1	9	5
	16 + 1	14	17
· · · · · · · · · · · · · · · · · · ·	4 + 1.5	49	44
	8 + 1.5	0	11
	12 + 1.5	22	23
· · · · · · · · · · · · · · · · · · ·	16 . 1 . 5	54	37
	16 + 1.5 4 + 2	65	
		7	48
	8 + 2	33	0
	12 + 2	29	3
2,4-D alkanolamine + sulfometuron 2,4-D alkanolamine + sulfometuron	16 + 2	60	61
2.4-D alkanolamine + sulfometuron	8 + 1	7	74
2,4-D alkanolamine + sulfometuron 2,4-D alkanolamine + sulfometuron	16 + 1	6	4
2,4-D alkanolamine + sulfometuron 2,4-D alkanolamine + sulfometuron	32 + 1	6 9	0
2,4-D alkanolamine + sulfometuron 2,4-D alkanolamine + sulfometuron	8 + 1.5		0
,4-D alkanolamine + sulfometuron ,4-D alkanolamine + sulfometuron	16 + 1.5	14	0
,4-D alkanolamine + sulfometuron ,4-D alkanolamine + sulfometuron	34 + 1.5	20	3
,4-D alkanolamine + sulfometuron ,4-D alkanolamine + sulfometuron	8 + 2	16	0 3 0
,4-D alkanolamine + sulfameturon	16 + 2	17	3
,4-D mixed amine a surfometuron	32 + 2	16	3 0
,4-D ester + 2 4 DD	16	14	7
,4-D alkanolamine + sulfometuron ,4-D mixed amine + sulfometuron ,4-D mixed amine + sulfometuron ,4-D ester + 2,4-DP + dicamba + sulfometuron	16 + 1	6	ó
Cloram	2 . 2		U
cloram	2 + 2 + 0.5	13	•
4-D alkanolamina	8	9	0
uroxypyr + sulfometuron	16	59	4
uroxynyn	16	6	35
"' OVIDAL & CILLAD 1	4 + 1	78	5
TONTUY! + CILL+Ome +.	8 + 1	38	48
- oxypyr	16 + 1		13
LSD (0.05)	16	49	5
		32 · · · · · · · · · · · · · · · · · · ·	13

Mixed amine salts of 2,4-D (2:1 dimethylamine:diethanolamine)-EH 736. 2,4-D isooctyl ester:2,4-DP butoxyethanol ester:dicamba (4:4:1)-EH 680.

Table 3. Leafy spurge control with sulfometuron applied with picloram, 2,4-D, or dicamba at Valley City, ND (Lym and Messersmith).

able 3. Leafy spurge control or dicamba at Valle	y City, ND (L	yill and he	Evaluation	n date	
			Evaluacion	Allu O	Retreat-
	D. La	Aug 87	June 88	Original	ment
reatment	Rate (oz/A)		(%)		25
Picloram + sulfometuron	8 + 1	98 99	88 87	19 26 15	39 26
Picloram + Sulfometuron	8 + 1.5 8 + 2	92	95	8	32
2,4-D alkanoramine	16 + 1	89	15	12	29
2-4-D alkanolamine	16 + 1.5	89	26	11	22
2,4-D alkanolamine	16 + 2 16 + 1	98 54	56 34	3 1 15	35 40 36
Dicamba + sulfometuron	16 + 1.5 16 + 2	45 84	11 80	15 0	36 28
Dicamba + sulfometuron Untreated	10 + 5	0 28	0 25	NS	NS
LSD (0.05)				ana_third	of each

a Picloram at 8 oz/A applied on June 2, 1988, to the back one-third of each plot.

Picloram and 2.4-D combination treatments for long-term leafy spurge management. Lym, Rodney G., and Calvin G. Messersmith. Picloram is an effective herbicide for leafy spurge control, especially when applied at rates from 1 to 2 1b/A. However, the high cost of picloram at 2 1b/A makes it uneconomical for treating large acreages of pasture and rangeland. Research by North Dakota State University has shown that picloram at 0.25 to 0.5 lb/A applied annually will give satisfactory leafy spurge control after 3 to 5 years. The purpose of this experiment is to establish the number of annual applications of picloram needed to provide 90 to 100% control of leafy spurge and to investigate possible synergism between picloram and 2,4-D.

The experiment was established at three locations in North Dakota, beginning at Dickinson on August 25, 1981, at Sheldon on September 1, 1981, and at Valley City on June 11, 1982. The soil at Dickinson was a loamy fine sand with pH 6.6 and 3.6% organic matter, at Sheldon was a fine sandy loam with pH 7.7 and 2.1% organic matter, and at Valley City was a loam with pH 6.7 and 9.4% organic matter. Dickinson, located in western North Dakota, generally receives much less precipitation than the other two sites located in eastern North Dakota. All treatments were applied annually, except 2,4-D alone was applied biannually (both spring and fall). Picloram treatments were applied in late August 1981 and in June of 1982 through 1988. The Sheldon location was discontinued following the fall evaluations in 1985. Thus, the Dickinson site has received eight picloram and picloram plus 2,4-D treatments and 15 2,4-D treatments, while the Valley City site has received seven and 14 treatments, respectively. The plots were 10 by 30 ft and each treatment was replicated four times in a randomized complete block design at all sites. Evaluations were based on percent stand reduction as compared to the control.

Picloram at 0.25, 0.38 and 0.5 lb/A provided 42, 67 and 69% leafy spurge control, respectively, 84 months after the original treatment (Table). Control had increased slightly compared to the previous year reversing a trend towards decreased control the last 2 years. 2,4-D at 1 to 2 lb/A alone provided an average of 59% control of leafy spurge after biannual applications for 7 years.

Leafy spurge control 84 months after treatment (MAT) increased by an average of 41, 24, and 26% when 2,4-D at 1 to 2 lb/A was applied with picloram at 0.25, 0.38, or 0.5 lb/A, respectively, when compared to the same picloram rate applied alone (Table). This increase (30%) was nearly twice as much in 1988 compared to the average (18%) increase the previous year. Visual observation indicated picloram alone was not absorbed well in 1988 as many plants showed little herbicide injury which was probably due to hot dry conditions during application. The greatest enhancement with 2,4-D plus picloram seems to be with 2,4-D at 1 lb/A and picloram at 0.28 lb/A. In general, leafy spurge control has been similar at all sites and does not seem to be influenced by soil types, pH, or organic matter.

All picloram at 0.38 or 0.5 lb/A plus 2,4-D treatments are near or have reached the target of 90% or better leafy spurge control. However, control declined an average of 10% between the 72 and 81 MAT evaluations for all picloram treatments. This indicates leafy spurge could reestablish quickly even after treatment for 7 years, although some of the reestablishment was due to invasion from border rows and control plots. An annual treatment would be needed to maintain control on a field scale, but perhaps more economical treatments than picloram would sustain the target control level. with approval of the Agric. Exp. Stn., North Dakota State Univ., Fargo 58105).

Table. Leafy spurge control from annual picloram or picloram plus 2,4-D treatments at two locations in North Dakota (Lym and Messersmith).

in Nort	th Dakota (Lym	and Mes	261.211	10117			<del> a</del> -		
און און און און	il Duite		W	onths	after '	treatme	ent"	01	84
				26	AR	00	72	81	
11-440	Rate	12	24		(% cont	ro1)			
Herbicide	(1b/A)				,,,			20	42
			AO	48	58	49	38	38	67
a. Janes	0.25	39	48	52	77	69	62	51	69
Picloram	0.38	65	62 71	81	86	77	71	60 29	59
Picloram	0.5	65	30	38	50	39	28	35	60
Picloram	1	22	24	26	45	49	35	37	57
2,4-D bian 2,4-D bian	1.5	22	30	26	54	54	26	70	83
2,4-D bian 2,4-D bian	2	19	66	63	85	73	87	65	84
Pic + 2,4-D	0.25 + 1	52 58	66	70	85	77	85 76	67	81
Pic + 2,4-D	0.25 + 1.5	57	62	66	83	76	82	72	89
Pic + 2,4-D	0.25 + 2	69	72	70	90	84	87	76	92
Pic + 2,4-D	0.38 + 1	68	74	76	93	84	90	81	93
Pic + 2,4-D	0.38 + 1.5	68	59	76	91	86	90	83	95
Pic + 2,4-D	0.38 + 2	71	75	84	94	87	93	89	95
Pic + 2,4-D	0.5 + 1	64	73	80	97	91	93	86	95
Pic + 2,4-D	0.5 + 1.5	76	75	81	95	91	33		
Pic + 2,4-D	0.5 + 2	, 0				14	13	14	13
		18	14	19	14	14			feat feet
LSD (0.05)		- 13 H							
					ant in	clude (	data fi	rom the	9

Mean values through 48 months after treatment include data from the Sheldon location which was discontinued after 1985.

Leafy spurge control with low rate annual picloram and 2,4-D combination treatments. Lym, Rodney G. and Calvin G. Messersmith. Previous research at 2,4-D for 3 to 5 years will give leafy spurge control similar to expensive gives 20 to 30% better leafy spurge control than picloram at 0.25 + 1 lb/A generally but the benefit of a herbicide combination declines as the picloram or 2,4-D 10% better control than picloram at 0.5 lb/A alone. The purpose of this treatments of picloram plus 2,4-D amine at relatively low application rates.

The experiment was established at four locations in North Dakota. Spring treatments were applied in June 1984 at Dickinson, Hunter, and Valley City, and the fall treatments were applied in September 1984 at Valley City and at the Sheyenne National Grasslands near McLeod. The soil was a loamy National Grasslands, and a loam at Valley City. Dickinson, located in western North Dakota, generally receives much less precipitation than the other two annually in June in 1984 through 1987. The fall treatments were applied in September 1984 and 1985, but were discontinued thereafter. The herbicides plots were 10 by 30 ft in a randomized complete block design with four replications except at Hunter which had 8 by 25 ft plots and three reduction as compared to the control.

The results from the Dickinson location were different than the other sites and will be discussed separately. Picloram at 0.12, 0.25, 0.38, and 0.5 lb/A provided 12, 24, 47 and 59% leafy spurge control, respectively, as a spring-applied treatment at Hunter and Valley City 48 months after treatment (MAT) (Table). Fall-applied treatments at Sheyenne and Valley City resulted in evaluation. Table). The addition of 2,4-D to picloram tended to increase leafy spurge control slightly for spring- but not for fall-applied treatments. The slight increase in control was similar regardless of 2,4-D rate. Spring-typically 0 to 3 inches tall when a killing frost occurred.

Leafy spurge control 48 months after spring-applied treatment averaged 36% with picloram alone and increased slightly to 40% when picloram was applied with 2,4-D (Table). This increase is much less than previously reported when 2,4-D at 1 to 2 lb/A was applied with picloram. Picloram at 0.5 lb/A plus 2,4-D averaged 73% leafy spurge control compared to 59% with picloram at 0.5 lb/A alone and was the best treatment evaluated.

Leafy spurge control averaged 26% 48 months after the initial treatment at Dickinson compared to 41% at Hunter and Valley City (Table). The reason similar experiment, begun in 1981 at Dickinson, resulted in annual increases for the first 36 months of the experiment and leafy spurge may be growing more vigorously than previously.

In general, leafy spurge control increased when 2,4-D was applied with picloram at 0.25 to 0.5 lb/A as a spring applied but not as a fall-applied picioram at 0.25 to 0.5 10/A as a spring applied but not as a latit-applied treatment. The 2,4-D application rate did not affect leafy spurge control with picloram. Picloram at 0.25 to 0.5 1b/A plus 2,4-D at 1 1b/A remains the most cost effective application rate for an annual leafy spurge control program. (Published with approval of the Agric. Exp. Stn., North Dakota State Univ., Fargo, 58105)

Table. Leafy spurge control in 1988 from annual pictoram or pictoram plus 2,4-D amine treatments spring or fall applied since 1984 at four locations in North Dakota (Lym and Messersmith).

1904 at .	rge control our locations					Amolica	tion ti	me/loca	tion/eva	lustion	Fa	11		
						Spring					11 11 70	City 19	86/1987	
						Valley	city_	Mean	Sheyenn	e-1987	Valley	417 20	May 20	Hear
		Hunt	er	Dickin	nson_	Aug 87	June 88	1988	May 30	Aug 24	June 3	AUG EU	May 20	
	Rate	Aug 87	May 88	Sept 87	June oo	AUG U.		-(% co	ntrol)					
atment	(1b/A)											0	1	0
	((2)				100	55	23	12	42	0	3	•	0	1
	0.12	3	0	2	13	62	35	24	67	0	25		2	6
loram	0.25	27	12	17	9		54	47	74	9	56	3	43	27
loram	0.38	67	39	64	29	81	64	59	89	16	92	38	17	4
clorem		79	53	76	25	82		12	72	0	32	8	200	6
cloram	0.5		3	3	13	57	21	5	62	8	12	0	0	1
cloram + 2,4-	0 0.12 + 0.		3	3	8	55	. 7	16	67	2	7	0	0	
clorem + 2,4-	0 0.12 + 0.	-		7	15	61	27		70	5	19	1	0	3
cloram + 2,4-	0 0.12 + 0.	•	20	40	11	70	28	28	64	0	18	1	0	9
ictoram + 2,4	0 0.25 + 0.			62	30	71	34	33		2	35	6	6	4
ictorm + 2,4	0.25 + 0.			30	28	73	23	21		15	56	11	14	13
iclor + 2,4	.0 0.25 + U	,5 2		45	15	81	49	47		6	48	3	4	
ictors + 2,4	-D 0.38 + V	.12 6		84	25	82	63	61		18	64	3	4	1
iclores + 2,4	-0 0.38 + 0	.25 8		52	37	88	59	45		15	75	8	8	1
iclor + 2,4	-0 0.38 + 0	.5 4	4 31	94	33	86	78	74		22		18	19	2
pictora + 2,4	-D 0.5 + 0	.12 9	2 72		46	83	78	7				15	7	1
pictor + 2,		.25	69		49	94	82	7		18		6	7	
Pictor . 2		).5	59		35	82	46	3	4 92	12	63			
Pictora + 2,			40 21	22	35	~						45	18	
Pictoran + 2,	4-A 0.22					19	25	9	2 28	) H:	31	15	10	
LSD (0.05)			31 26	3 26	. 24	19	23		1					

Average control at Hunter and Valley City 48 months following the original 1984 treatment date.

Average control 24 months following the original 1984 treatment date, fall treatments discontinued after 1985.

Leafy spurge control with picloram plus dicamba or various 2,4-D formulations applied annually. Lym, Rodney G., and Calvin G. Messersmith. Picloram remains the most effective herbicide for leafy spurge control. Previous research at North Dakota State University has shown picloram +2,4-D at 0.25+1 lb/A applied annually to be more cost effective than picloram at 1 to 2 lb/A applied once. The purpose of these experiments was to compare the effect of dicamba and/or various 2,4-D formulations applied with picloram for leafy spurge control.

The initial 2,4-D formulation experiments were established on the Sheyenne National Grasslands near McLeod, ND, on June 15, 1984, and near Hunter, ND, on May 30, 1985. The herbicides were applied using a tractor-mounted sprayer delivering 8.5 gpa at 35 psi. All plots were 10 by 30 ft in a randomized complete block design with four replications. Evaluations were based on visible percent stand reduction as compared to the control.

Picloram + 2,4-D mixed amine provided better leafy spurge control than picloram + 2,4-D alkanolamine (Table 1). Leafy spurge control from picloram + 2,4-D mixed amine at 0.25 + 1 lb/A was similar to control from picloram at 0.5 lb/A alone, but picloram + 2,4-D is approximately 30% less expensive. Similarly, leafy spurge control from picloram + dicamba was greater when applied with 2,4-D mixed amine than with the alkanolamine. Neither 2,4-D formulation alone controlled leafy spurge.

Picloram + dicamba + 2,4-D mixed amine provided 72% leafy spurge control 2 yr after application at Hunter (Table 1). This level of control was similar to that attained with picloram at 2 lb/A in North Dakota but is 70% less expensive. Therefore, similar experiments were begun in 1986 to evaluate this combination treatment further. Experiments were established on June 11 and 18, near Dickinson and Valley City, respectively, and on August 28 on the Sheyenne National Grasslands and September 3 and 15 near Valley City and Dickinson, respectively. Treatments were applied annually as previously described in the spring or fall through 1988.

Leafy spurge control was similar regardless of the 2,4-D formulation applied with picloram + dicamba in the spring (Table 2). Control was only 20% when averaged across all treatments and both locations 1 yr after the second application. This is much lower than would be expected with picloram + 2,4-D at 0.25 + 1 lb/A which averages 60% or more based on long-term observations.

Leafy spurge control with picloram at 0.5 lb/A averaged 32% 1 yr following the second fall application (Table 2). Control improved to 49% when applied with dicamba at 2 lb/A. Leafy spurge control with picloram + dicamba was not improved by adding 2,4-D regardless of the 2,4-D formulation.

In general, leafy spurge control was similar with all 2,4-D formulations in combination with picloram and dicamba. However, the 2,4-D mixed amine formulation occasionally did provide better leafy spurge control in a combination treatment than the alkanolamine or ester formulations. These experiments will be continued to evaluate the long-term effect of picloram combined with various 2,4-D formulations and dicamba on leafy spurge control. (Published with approval of the Agric. Exp. Stn., North Dakota State Univ., Fargo, 58105)

Table 1. Leafy spurge control with picloram applied with various formulations of 2,4-D (Lym and Messersmith).

Location/		Mo	nths af	ter tre	atment	
application date	Rate	3	12	15	24	27
Treatment	(1b/A)		(%			
	(10/11)					
Sheyenne, June 1984						
D 1 3	0.25	76	23	4	1	
Picloram	0.5	95	75	43	10	• •
Picloram + 2,4-D						
alkanolamine	0.25 + 1	78	14	6	3	
Picloram + 2,4-D						
mixed amine	0.25 + 1	94	72	23	21	• •
2,4-D mixed amine	4	47	7	13	0	
2,4-D alkanolamine	4	42	20	7	5	• •
2,10				9 6	3.0	
LSD (0.05)		15	25	15	12	
Hunter, June 1985						
Picloram + dicamba					7.0	60
+ 2,4-D mixed amine	0.25 + 1 + 2	99	98	89	72	60
Picloram + dicamba	a Falai S da med		-	0.5	25	10
+ 2,4-D alkanolamine	0.25 + 1 + 2	51	51	25	25	18
2,4-D mixed amine	4	6 5	3	0	0	0
2,4-D alkanolamine	4		0	0 15	0	7
Picloram + dicamba	0.25 + 1	53	38	13	U	,
		15	15	15	15	20
LSD (0.05)		10	10	10		

a Mixed amine salts of 2,4-D (2:1 dimethylamine:diethanolamine)-EH 736.

Table 2. Leafy spurge control with picloram applied with dicamba and various formulations of 2,4-D applied annually since 1986 for leafy spurge control (Lym and Messersmith).

Application date/		V-22	Locatio	on/1988	evalu	ation d	ate	
treatment	Rate		CILY	DICKI	nson	_Shey	enne	
or occupant	(lb/A)	June	Aug	June		June	Sept	Mean
	(10/1/			(%	contro	1)		
Spring								
2,4-D mixed amine +b								
dicamba+picloram,	0.25+1.25	24	63	10	79	• •	• •	17
dicamba+picloram 2,4-D mixed amine+	2+0.5+0.25	17	59	26	88	• •	• •	22
picloram+dicamba	1+0.5+0.12	15	52	12	76			
2,4-D alkanolamine+					, 0	• •	• •	13
dicamba+picloram Dicamba+picloram	2+1+0.25 1+0.25	25 32	54	18	84	• •	• •	22
a protor am	170.23	32	56	20	86	• •		, 26
LSD (0.05)		NS	NS	NS	NS			NS
Fall								
2,4-D mixed amine +b								
dicamba+picloram	2+1+0.25	84	31	66	8	73	38	26
2,4-D alkanolamine+ dicamba+picloram	2+1+0.25	89	20					
2,4-D mixed amine +	2+1+0.25	09	29	• •	0 0	73	54	42
dicamba+picloram 2,4-D ester + 2,4-DP	4+2+0.5	96	61	94	45	93	67	56
+picloram								
+dicamba 2+ 2,4-D_ester <sup>C</sup> + 2,4-DP	2+0.5+0.25	74	14	49	8	74	46	22
+picloram							,,,	22
	2+0.5+0.5	96	47	82	35	00	67	
4-D alkanolamine+	2.000.00	30	17	02	33	93	67	50
dicamba+picloram	4+2+0.5	94	50	81	39	90	61	50
licamba+picloram	2+0.5	95	41	93	51	83	56	49
icloram	0.5	95	46	57	15	65	33	32
LSD (0.05)		6	20	20	0.6			
(3,00)		0	20	32	36	17	27	11

a Mean 24 months after first treatment.

b Mixed amine salts of 2,4-D (2:1 dimethylamine:diethanolamine)-EH 736.

c 2,4-D isooctyl ester:2,4-DP butoxyethanol ester:dicamba (4:4:1)-EH 680.

Late-season leafy spurge control with sulfometuron and picloram. Lym, Rodney G. and Calvin G. Messersmith. Previous research has shown the best time to apply herbicides for leafy spurge control is the true flower growth stage in mid-June or during fall regrowth. Late-fall treatments generally provide good control of many perennial weeds, but leafy spurge shows more visible changes due to cold temperatures and changes in daylength than most other perennial weeds. Many landowners often are unable to treat all of the infestations present before a light frost occurs and/or the leaves lose chlorophyll and turn red. The purpose of this research was to determine if frost or changes in leafy physiology prior to herbicide application would prevent satisfactory leafy spurge control the following growing season.

The experiment was established in dense leafy spurge stands at West Fargo and Hunter, ND in 1987. Treatments were applied on four dates from September 1 to October 14, and leafy spurge ranged from vigorous fall regrowth to senescence (Table 1). Fall 1987 was warmer than average as there were no temperatures colder than 38 F in September, and the coldest overnight temperatures during the experiment were 31 and 24 F on October 2 and 7, respectively. The herbicides were applied with a tractor-mounted sprayer delivering 8.5 gpa at 35 psi. All plots were 10 by 30 ft in a randomized complete block design with four replications. Evaluations were based on a visual estimate of percent stand reduction as compared to the control.

Leafy spurge control was better at Hunter than West Fargo regardless of treatment on application date (Table 2). The environmental conditions were similar at the two locations, except leafy spurge at Hunter was growing more vigorously, probably due to higher soil moisture available than at West Fargo. Leafy spurge control at West Fargo was similar when herbicides were applied on September 1 and 15, and averaged 73% in May 1988. Leafy spurge leaves had turned red by September 15 but were firmly attached to the stem (Table 1). However, the plants were beginning to defoliate when the treatments were applied on October 1 and the stems were bare by October 14. Control declined accordingly and averaged only 49 and 30%, respectively. Leafy spurge control with picloram at 16 oz/A tended to be better than with sulfometuron plus picloram at 1 + 8 oz/A, but both treatments were better than picloram plus 2,4-D at 8 + 16 oz/A. The plots were burned during a grass fire in July 1988 and could not be reevaluated.

Leafy spurge leaves at Hunter had turned red by September 15, but did not defoliate during the experiment despite the temperature dropping to 24 F on October 7 (Table 1). Leafy spurge control was similar in May 1988 regardless of treatment date at Hunter except for picloram plus 2,4-D at 8 + 16 oz/A which averaged only 36% control when applied on October 1 (Table 2). The reason for the decline is not known, since this treatment was applied prior to frost and the October 15 treatment averaged 81% control. Sulfometuron plus picloram at 1 + 8 oz/A and picloram at 1 + 8 oz/A provided better long-term control than picloram plus 2,4-D at 1 + 8 oz/A when applied on September 15 or later.

Soil and air temperature (including below 32 F) and relative humidity did not affect leafy spurge control with these herbicides. Good control can be expected the following spring if fall treatments are applied when leafy spurge leaves are green or red but still firmly attached to the stem. (Published with approval of the Agric. Exp. Stn., North Dakota State Univ., Fargo, 58105).

Table 1. Growth and environmental conditions during late-season leafy spurge control treatments (Lym and Messersmith).

1987	Wes	t Fargo		2 10 10 10 10				W a		
treatment date	<u>Leafy</u> Stems	spurge Leaves	Air	Soil 1 inch	Relative humidity		Spurge Leaves		erature Soil	Relative
September 1			(	(F)	(%)		reaves		1 inch (F)	humidity (%)
achrempel 1	branched	green	68	70	51	branche	ed slightly red	67	71	53
September 15	red	red	81	74	45	yellow	red	79	72	
October 1	red	senes- cence	59	60	65	red	red	61	60	58 60
October 15	bare		50	48	60	red ,	red	48	48	60

Table 2. Leafy spurge control from late-season herbicide treatments (Lym and Messersmith).

Picloram + 2,4-D 8 + 16 92 47 91 18 36 0 81 + picloram 1 + 8 100 59 99 70	t 15
Picloram + 2,4-D 8 + 16 54 60 24 10  Sulfometuron	Au
Picloram + 2,4-D 8 + 16 54 60 24 10  Sulfometuron	
Sulfometuron + picloram 1 + 8 54 80 56 43 Picloram 16 77 94 67 38  LSD (0.05) May = 38  Hunter Picloram + 2,4-D 8 + 16 92 47 91 18 36 0 81 + picloram 1 + 8 100 59 99 70	
Picloram 16 77 80 56 43 LSD (0.05) May = 38  Hunter Picloram + 2,4-D 8 + 16 92 47 91 18 36 0 81 + picloram 1 + 8 100 59 99 70	
16 77 94 67 38  LSD (0.05) May = 38  Hunter  Pictoram + 2,4-D 8 + 16 92 47 91 18 36 0 81  + pictoram 1 + 8 100 59 90 70	
LSD (0.05) May = 38  Hunter  Picloram + 2,4-D 8 + 16 92 47 91 18 36 0 81  + picloram 1 + 8 100 59 90 70	
Hunter Picloram + 2,4-D 8 + 16 92 47 91 18 36 0 81 + picloram 1 + 8 100 59 90 70	
Picloram + 2,4-D 8 + 16 92 47 91 18 36 0 81 + picloram 1 + 8 100 59 99 79	
Sulfometuron 91 18 36 0 81 + picloram 1 + 8 100 59 99 79	
Sulfometuron 91 18 36 0 81 + picloram 1 + 8 100 59 99 79	
+ pictoram 1 + 8 100 59 00 70	23
100 39 00 %	
16 0/ 59 39 93	61
72 42 91 38 86	56
LSD (0.05) May = 26; Aug = 25	

Leafy spurge control with resulting forage production from several Lym, Rodney G., and Calvin G. Messersmith. An experiment to evaluate long-term leafy spurge control and forage production herbicide treatments. was established at two sites in North Dakota in 1983. The predominate grasses were bluegrass (Poa spp.) with occasional crested wheatgrass, smooth brome, big bluestem, or other native grasses. The treatments were selected based on previous research conducted at North Dakota State University and included 2,4-D at 2 1b/A, picloram + 2,4-D at 0.25 + 1 1b/A, picloram at 2 lb/A, and dicamba at 8 lb/A, and were applied in August 1983 or June 1984 as fall or spring treatments. The 2,4-D at 2 lb/A and picloram plus 2,4-D treatments were applied annually, while the picloram alone and dicamba treatments were reapplied when leafy spurge control declined to 70% or less. Sulfometuron + picloram at 0.08 + 0.5 lb/A were applied to previously untreated plots in June or August, 1988. The plots were 15 by 50 ft with four replications in a randomized complete block design at each site. Forage yields were obtained by harvesting a 4 by 25 ft section with a rotary mower in July 1984 and every year thereafter. Sub-samples were taken by hand along each harvested strip and separated into leafy spurge and forage so the weight of each component in the mowed sample could be calculated. The samples were oven dried and reported with 12% moisture content. Economic return was estimated by converting forage production to animal unit days (AUD) and then to pounds of beef at \$0.60/1b minus the cost of the herbicide and estimated application cost, i.e., 2,4-D = \$2.00/lb ae, dicamba = \$11.75/lb ae, picloram = \$40.00/lb ae, and application = \$2.05/A. The cost of treatments applied in fall 1988 is not subtracted from the net return.

Most treatments resulted in less economic gain at Dickinson than Valley City despite excellent leafy spurge control from several treatments. Dickinson generally receives 5 to 6 inches less precipitation annually than Valley City. Total forage production averaged after 5 yr across all treatments was 4301 lb/A at Dickinson and 9043 lb/A at Valley City (Table). Leafy spurge control from 2,4-D at 2 lb/A was not satisfactory from spring or fall applications at either site. However, 2,4-D provided short term control resulting in an economic gain of \$114/A and \$81/A at Valley City and of \$34/A and \$43/A at Dickinson as spring and fall applied treatments, respectively. Leafy spurge control with picloram + 2,4-D at 0.25 + 1 lb/A averaged over both locations was 78% in 1987 (Table) as a spring applied treatment. Much below average precipitation was received at both locations in 1988 but timely rains in June at both locations resulted in average forage production. Leafy spurge control was poor with picloram + 2,4-D at 0.25 + 1 lb/A fall applied, but forage production (averaged across locations) of 7142 lb/A was only slightly less than the spring average of 8282 1b/A.

Picloram at 2 lb/A provided good leafy spurge control and a slight economic gain at Valley City, but an average loss of \$114/A at Dickinson (Table). Dicamba generally gave good leafy spurge control as a fall but not as a spring applied treatment. All treatments have reduced leafy spurge production compared to the control. (Published with approval of the spurge Stn., North Dakota State Univ., Fargo.)

Table. Leafy spurge control, forage production and estimated net return from several herbicide treatments at two sites in North Dakota (Lym and

Original	treatme	ent Re-tr	eatmen	t			007			
da Herbicide		tim	e				987		Yielda	Total
HELD ICIDE	Rate		e Rat	e Yea	r Cost	100	ntrol	_ For	- Leafy	/ net
	(1b/A	()	(1b/	A)	(\$/A			age	spurc	ie returnb
					(4/A)	)	(%)	(1	b/A)	- (\$/A)
Spring 198	34	_Spring						Va	lley C	
2,4-D	2	2,4-D	20	84-8	00 00				incy c	rcy
Picloram	0.2	5 Picloran	1 0.25		30	16	28	962	2 372	3 114
+ 2,4-D Picloram	+1	+ 2,4-[	+10		8 70	58		100-		
Dicamba	2	Picloram	20	88	164	68				
Sulfomature	8	Dicamba	80	85-8	7 384			1140		1 7
Sulfometure + piclorar				00 0	, 204	58	93	1084	1 1986	5 (-221)
piciol di	m +0.5	• •	• • •	88	20		43	2347	7 1676	
Fall 1983		_ Fall						2047	7 1679	15
2,4-D	2	2,4-D	2C	84-88	2 20					
Picloram	0.25	Picloram	0.25	04-88	30	6	4	7427	6596	81
+ 2,4-D	+1	+ 2,4-D	+1C	84-88	3 70	C0	0.0			O1
Picloram	2	Picloram	2d	85	164	69	22	9722		76
Dicamba	8	Dicamba	8d	86		96	84	10988		0
100 /0		Control		00	192	86	69	10788		(-30)
LSD (0.05	)				•	0	0	9352	7706	( 00)
						18	13	2079	1665	
Spring 1984		Spring						Dick.	inson	
2,4-D	2	2,4-D	2C	04.07				DICK	1113011	
Picloram	0.25	Picloram	0.25	84-87	30	7	40	4268	529	34
+ 2,4-D	+1	+ 2,4-D	+1C	84-87	7.0				025	34
Picloram	2	Picloram	2d	86,87	70	49	89	6312	158	24
Dicamba	8	Dicamba	8d	85-87	246	60	99	6177	122	(-153)
ulfometuron			0-	05-8/	384	43	99	5063	352	(-308)
+ picloram	+0.5			88	20		9 9			, 555)
all 1983		5 40 1 1 1 1 1			20		11	1189	479	(-2)
4-D	2	Fall Park	Day The							THEFT
icloram		2,4-D	2C	84-87	30	10	11	4927	1200	A SHOTE SALE
+ 2,4-D	0.25 +1	Picloram	0.25				44	7361	1382	43
icloram	2	+ 2,4-D	+1C	84-87	70	21	22	4561	1200	
icamba	8	Picloram	2d	86	164	84	68	5971	1269	(-2)
- Gillou	0	Dicamba	8d	85,86	288	77	64	5604	47	(-74)
LSD (0.05)		Control				0	0	4117	109	(-204)
-00 (0.00)						17	20	797	2882 484	

a Total production of 1984 through 1988 harvest. b Total net return for 1984 through 1988. Fall 1988 treatment cost is not c Annual retreatment.

d Applied when control declines to less than 70%.

Leafy spurge control following a nine-year management program. Rodney  $\overline{G}$ , and Calvin G. Messersmith. An experiment to evaluate long-term leafy spurge management was established at four sites (Sheyenne National Grassland near McLeod, Sheldon and two near Valley City) in North Dakota in 1980. All sites were established in early June, except one site at Valley City which was established in September 1980. The herbicides applied in 1980 included 2,4-D as liquid and picloram as liquid (2S) and granular (2%G) formulations, and picloram applied using the roller and pipe-wick applicators. The conventional broadcast treatments were applied using a tractor-mounted sprayer delivering 8 gpa water at 35 psi. A granular applicator was used to apply the picloram 2%G treatments. Solution concentration in the roller was 0.25 lb/gal; this is the same solution concentration as picloram at 2 lb/A sprayed at 8.5 gpa. The solution concentration was increased for the pipe-wick applicator to picloram at 0.5 lb/gal, because the pipe-wick applied about half the total volume per acre as the roller applicator. The roller and pipe-wick applicator height was adjusted to treat the top one-half of the tallest leafy spurge stems. The additive in the roller and pipe-wick treatments was a 5% (v:v) oil concentrate (83% paraffin based petroleum oil plus 15% emulsifier). The plots were 15 by 150 ft and treatments were replicated twice at each site in a randomized complete block design. Each plot was divided into six 7.5 by 50 ft subplots and retreatments of 2,4-D, picloram 2S, dicamba or no treatment were applied in June 1981, except the fall Valley City site which was retreated in August 1981.

Original 1980 whole plot treatments were reapplied in 1982 with several of the treatments changed (see Table). A carpet applicator was substituted for the roller applicator. The carpet applicator was designed by Magnolia Spray Equipment Corp., Jackson, MS, and consists of a 1 by 8 ft carpet attached to a rectangular spray box. The herbicide solution was sprayed onto the backside of the carpet through nozzles inside the spray box. Excess solution was returned to the spray tank. The picloram solution on the carpet applicator was 0.25 lb/gal and 0.4 lb/gal for two and one pass applications, respectively. The granular picloram treatments were replaced by picloram applied with the pipe-wick or carpet applicator with two passes, the second pass in the opposite direction to the first. Dicamba at 8 lb/A spray applied replaced the picloram plus oil concentrate pipe-wick applied treatment. The whole plots were retreated in 1982 with the original treatment, except picloram at 2 lb/A was reapplied to the control subplot only since subplots receiving annual retreatments maintained satisfactory leafy spurge control. The experimental site at the Sheyenne National Grasslands was treated in the fall of 1982 to establish an equal number of spring and fall treatment sites. Subplot retreatments were applied again in 1983 through 1988. Evaluations are based on visual percent stand reduction as compared to the control.

Previously, leafy spurge control has been higher from spring than fall-applied treatments (data not shown). However, control averaged 33 and 46% from the spring- and fall-applied treatments, respectively, in 1988 (Table). Leafy spurge was drought stressed when the treatments were applied in 1988 from very dry and hot conditions which probably resulted in poor herbicide absorption.

Post-harvest treatments for field bindweed control. Lym, Rodney G. Previous research has shown that dicamba fall applied provides good field bindweed control the following growing season. Field bindweed is a problem weed in western North Dakota, especially where minimum till and strip-fallow farming are common practices. The purpose of these experiments was to evaluate several herbicides as single and combination treatments for late-

Two experiments were established on September 10, 1987, on a dense stand of field bindweed near the Ranch Headquarters of the Dickinson Experiment Station. The herbicides were applied in 6- to 8-inch corn stubble which had been harvested 7 days prior to treatment. The field bindweed was in the vegetative growth stage with 8 to 10 inch long stems and was growing vigorously. The herbicides were applied using a tractor-mounted sprayer delivering 8.5 gpa at 35 psi. All plots were 9 by 30 ft in a randomized complete block design with four replications. The weather was overcast, 68 F, 50% relative humidity with a soil temperature of 66 F at 4 inches. 2-row barley was seeded in April 1988, but due to severe heat and drought was not harvested. Field bindweed control and barley injury evaluations were based on a visual estimate of percent stand reduction as compared to the

The first experiment evaluated dicamba applied either alone or with various auxin herbicides or glyphosate and various picolinic acids for field bindweed control. Dicamba at 64 oz/A and glyphosate plus dicamba at 90 + 44oz/A provided 79 and 70% field bindweed control, respectively, 9 months after treatment (MAT) (Table). Barley injury averaged 86% with dicamba at 64 oz/A but declined to an average of 26 and 16% when applied at 32 oz/A alone or at 16 oz/A with 2,4-D, respectively. Glyphosate plus dicamba at 45 + 22 oz/A tended to provide better field bindweed control (64%) than dicamba alone at 32 oz/A (41%). Treatments that included picloram applied at 4 oz/A caused severe auxin injury to the barley after heading but did provide the best season-long field bindweed control (data not shown). The addition of triclopyr and/or 2,4-D to picloram or dicamba did not increase control compared to the herbicides applied alone at similar rates.

The second experiment compared fluroxypyr alone and with auxin herbicides for field bindweed control. Field bindweed control averaged only 20% 9 MAT regardless of the fluroxypyr application rate but improved to 41% when applied with 2,4-D (Table). Dicamba plus 2,4-D at 16 + 16 oz/A and glyphosate plus 2,4-D at 64 + 58 oz/A provided similar field bindweed control, averaging 59%. Dicamba at 32 oz/A provided 71% field bindweed control but also 71% barley

Dicamba at 32 to 64 oz/A, picloram at 2 to 4 oz/A and glyphosate plus dicamba at 45 + 22 oz/A provided fair field bindweed control 9 MAT, but only the glyphosate plus dicamba treatment did not injure barley severely. Severe drought occurred in 1988 with 70% below average precipitation received from September 1987 to August 1988. The dry conditions probably accentuated injury to the barley crop from these treatments. (Published with approval of the Agric. Exp. Stn., North Dakota State Univ., Fargo).

Table. Field bindweed control from several post-harvest herbicide treatments applied in September 1987 (Lym).

pplied in September 1987 (Lym).		Evaluation 3	June 1988
THE RESERVE OF THE PROPERTY OF		National Parket	Barley
	Rate	Control	injury
reatment	(oz/A)	0000000	(%)
2011			
experiment One	8 + 0.5 + 8	10	3 9
1 - nich + nich Ordin + Car	8 + 0.5 + 16	15	18
	8 + 4 + 8	34	13
	8 + 4 + 16	10	6
	16 + 0.5 + 8	35	13
	16 + 0.5 + 16	6	12
	16 + 4 + 8	19	19
	16 + 4 + 16	28	10
	12 + 2.2 + 12	18	11
- · · · · · · · · · · · · · · · · · · ·	17 + 2.2 + 12	40 35	13
- 1 1 L N1C   D1C	7 + 2.2 + 12	31	16
	12 + 4.4 + 12	31	
	12 + 0.12 + 12	18	5 8
	12 + 2.2 + 17	23	10
	12 + 2.2 + 7	20	4
Triclonyr + DICIOFall ' -, "	64 + 58	45	16
04 L 04 A / H-11-	16 + 16 + 0.5%	41	26
Dicamba + 2,4=0= ' ^ '	32 + 0.5%	79	86
Dicamba + X=//	64 + 0.5%		
Dicamba + X-77 Dicamba + X-77	EST Dita on the two	31	23
2,4-D mixed amine <sup>C</sup> + picloram	32 + 4 + 16	50	16
+ dicamba	32 + 4 + 16	51	13
2,4-Da + picloram + dicamba	4 + 16	49	53
Picloram + dicamba 2,4-D esterd + 2,4-DP + dicamba	160 + 160 + 4	non haselunts.	
2,4-D esterd + 2,4-DP + dicamba		49	66
2,4-D estern + 2,4 Di	160 + 160 + 40 + 4	64	14
+ picloram Glyphosate + dicamba	45 + 22	70	4
I	90 + 44	37	
Glyphosate + 2,4-Da + dicamba	64 + 58 + 8	35	1
LSD (0.05)			
Experiment Two	2 2 2 2 2 2	14	
Fluroxypyr	2 4	5	
Fluroxypyr	6	41	1
Flurnyvnvr	2 + 16	34	
$Fluroxypyr + 2,4-D^{\alpha}$	1 + 16	23	e dogs l'h
picloram + 2,4-0"	1 + 16	41	1
Picloram + dicamba	16 + 16	59	beaugah .
Dicamba + 2,4-Da	32	71	Sections
Dicamba	64 + 58	58	
Glyphosate + 2,4-0°	Ladhempson & Seems Puri	24	
LSD (0.03)			
a 2,4-D alkanolamine.			

a 2,4-D alkanolamine.
b 2,4-D isopropylamine.
c Mixed amine salts of 2,4-D (2:1 v/v diethylamine:diethanolamine)-EH736.
d 2,4-D isoctylester:2,4-DP butoxyethanol ester:dicamba (4:4:1 v/v/v)-EH680.

Triazine-tridiphane mixtures for postemergence grass control in corn, Fargo, 1988. Interstate 201 corn was planted 2 inches deep on May 5 at 22,000 seeds per acre. Silty clay soil was conventionally prepared. Herbicides were applied May 28 when relative humidity was 87%, air temperature was 83 F, corn was 2.5 to 3.5-leaf (mostly 3.5-leaf), wild oats were 1 to 2.5-leaf and 2 to 4-inches tall, and yellow foxtail was 1 to 2-leaf and 0.25 to 1-inch tall. Soil conditions were moist with no plant stress evident at time of spraying. Treatments were applied using a bicycle wheel sprayer delivering 8.5 gpa with 8001 nozzles and 40 psi. Estimates of percent corn injury and weed control were taken on June 6 during significant drought stress. Plot size was 10 by 25 ft and the experiment was a randomized complete block design

<u>Treatment</u> <sup>a</sup>	Rate	Corn		
	(1b/A)	injury		Yef
Cyanazina	(10/1/)		(%)	
Cyanazine	1.5			
Atrazine+Cyanazine	0 125.1 275	11	52	73
ALI'dZINe+(.vanazina	0.125+1.375	11	73	73
Atrazine+Cvanazino	0.25+1.25	11	83	75
Aurazine+Cvanazino	0.375+1.125	12	84	77
Atrazine+Cvanazino	0.5+1	11	83	
Tridiphane+Cyanazine	0.75+0.75	ii		75
Tridinhane+Atmazine	0.5+1.5	12	84	73
Tridiphane+Atrazine+Cyanazine	0.5+0.125+1.375	11	95	92
Tridiphane+Atrazine+Cyanazine	0.5+0.25+1.25		97	87
	0.5+0.375+1.125	13	98	92
	0.5+0.5+1	13	99	94
	0.5+0.5+1	12	98	92
	0.5+0.75+0.75	12	100	89
Ilulonane+Atrazina.c.	0.75+1.5	13	98	95
	0.75+0.125+1.375	14	100	94
	0.75+0.25+1.25	13	100	
ridiphane+Atrazine+Cyanazine	0.75+0.375+1.125	13		93
ridinhane+Atmazine	0.75+0.5+1	13	100	94
ridiphane+Atrazine+Cyanazine ontrol	0.75+0.75+0.75		99	90
3110101	0		100	95
V or		4	0	0
V. %		THE REAL PROPERTY.		
SD 5%		15	5	5
ll treatments were applied wit rate containing 15% emulsifier yanazine and atrazine were use	1 1	2	6	_

cyanazine and atrazine were used.

Summary. Tridiphane generally increased corn injury although only by 2 to 3%. Wild oat control by cyanazine alone was increased when part of the cyanazine was replaced with atrazine in a tank mixture, although maximum control by cyanazine-atrazine combinations in the absence of tridiphane was 85%. wild oats was achieved when cyanazine-atrazine combinations were mixed with tridi-Essentially complete control of Foxtail control without tridiphane was about 75%, regardless of the proportion of atrazine in the mixture. The addition of tridiphane to cyanazine-atrazine combinations increased control to about 90 to 95%.

Corn plantback following imazaquin and imazethapyr in no-till soybeans, Fargo, 1988.

Fall (F) treatments were applied October 29, 1986, early preplant (EPP) treatments were applied April 25, 1987, and preemergence (Pre) treatments were applied June 8, were applied April 25, 1987, and preemergence (Pre) treatments were applied June 8, 1987 using a bicycle wheel sprayer delivering 17 gpa with 8002 nozzles and 35 psi. McCall soybeans were seeded on June 3 using a Hiniker no-till planter. Soil type was a silty clay with pH of 8 and organic matter of 5%. Interstate 201 corn was planted on May 5, 1988 into undisturbed soybean stubble using a Hiniker no-till planted on May 5, 1988 into undisturbed soybean stubble using a Hiniker no-till planted on May 5, 1988 into undisturbed soybean stubble using a Hiniker no-till planted on May 5, 1988 into undisturbed soybean stubble using a Hiniker no-till planted on May 5, 1988 into undisturbed soybean stubble using a Hiniker no-till planted on May 5, 1988 into undisturbed soybean stubble using a Hiniker no-till planted on May 5, 1988 into undisturbed soybean stubble using a Hiniker no-till planted on May 5, 1988 into undisturbed soybean stubble using a Hiniker no-till planted on May 5, 1988 into undisturbed soybean stubble using a Hiniker no-till planted on May 5, 1988 into undisturbed soybean stubble using a Hiniker no-till planted on May 5, 1988 into undisturbed soybean stubble using a Hiniker no-till planted on May 5, 1988 into undisturbed soybean stubble using a Hiniker no-till planted on May 5, 1988 into undisturbed soybean stubble using a Hiniker no-till planted on May 5, 1988 into undisturbed soybean stubble using a Hiniker no-till planted on May 5, 1988 into undisturbed soybean stubble using a Hiniker no-till planted on May 5, 1988 into undisturbed soybean stubble using a Hiniker no-till planted on May 5, 1988 into undisturbed soybean stubble using a Hiniker no-till planted on May 5, 1988 into undisturbed soybean stubble using a Hiniker no-till planted on May 5, 1988 in

		1988
1987 treatment	Rate C	orn injury
applied to soybeans	(oz/A) -	(%)
appirous	(02/11)	
	1.25	0
Imazaquin(F)	1.5	0
Imazaquin(F)	2	0
Imazaquin(F)	1+0.5%	0
Imazaquin+X-77(EPP)	1+0.5%	0
Imazaquin+X-77(EPP)	1.25+0.5%	0
Imazaquintx //(EPP)	1.5+0.5%	0
Imazaquin+X-77(EPP)	2+0.5%	0
Imazaquin+X-77(EPP)	1	0
Imazaquin(Pre)	1.25	0
Imazaquin(Pre)	1.5	0
Imazaquin(Pre)	2	0
Imazaquin(Pre)	2	
Imazethapyr(F)	3	0
Imazethapyr(F)	4	0
Imazothanvr(t)	1.5+0.5%	0
Imazothanvrtx-//(EPP)	2+0.5%	0
ImazethanvrtX-//(EFF)	3+0.5%	0
ImazethanvrtX-//(EFF)	4+0.5%	0
Imazethapyr+X-//(EFF)	1.5	0
Imazethapyr(Pre)	2	0
Imazethapyr(Pre)	3	0
Imazethapyr(Pre)	4	0
T+hanyr(Pre)	1+1.5	0
The anguint   mazethaby   ( )	1.25+2	0
	1+1.5+0.5%	0
	1.25+2+0.59	% O
z guint mazeliduvi 'n '/ \-'	1+1.5	0
t muintimazer.liduvi ( l' c)	1 2512	0
Imazaquin+Imazethapyr(Pre)	1.25+2	0
Imazaquini inazoonaro	0	
Control	L 1	e injury to co
1	ally observabl	e illiguity co

 $\frac{\text{Summary}}{12}$ . None of the treatments caused visually observable injury to corn planted  $\frac{\text{Summary}}{12}$  to  $\frac{13}{13}$  months after application until no-till conditions.

Weed control economics in minimum till and no-till soybeans, Fargo. The experiment was established in 1988 as a multi-year study on a silty clay soil having a pH of 7.8 and organic matter of 5%. Treatments were arranged as a split plot with three tillage-row spacing combinations serving as main plots and herbicide systems constituting sub-plots. Minimum till plots were chisel plowed in late October 1987. McCall soybeans were seeded 1 to 1.5 inches deep at 58 lbs/A on May 25 using a Hini-ker no-till planter for 30-inch rows and a Haybuster drill for 7-inch rows. Herbicides were applied using a bicycle wheel sprayer delivering 8.5 gal/A for all treatments involving glyphosate or sethoxydim and 17 gal/A for all other treatments. All

Table 1. Rates and dates of herbicide applications a.

Herbicide	<u>cide treatme</u> Rate	nt Date	<u>burndow</u>	neede	tment	<u>emerqen</u>	v-up po ce trea	St- tment
	(1b/A)	Dute	Herbicid		e Date	Herbicide	Rate	Dat
Tillen				(1b/	A)		(1b/A	Dat
TILLED, 30-INCH	ROWS						(15/11)	
rif+Metr(PPI)	1+0.2	5/24	None					
Trif+Imep(PPI)	1+0.063	5/24		-		Sethoxydm	0.15	6/22
Trif+Clom(PPI)	1+0.75	5/24	None	4		Sethoxydm	0.15	6/20
Trif+Clam(PPI)	1+2.5	5/24	None	-	•	Sethoxydm	0.15	6/22
Seth(Po)/	0.2/	6/13	None	-		Sethoxydm	0.15	6/22
Bent+Acif(Po)	0.75+0.13		4.			and the state of t	0.15	6/22
HWC - Trif+	0.75+	6/20	None	•		None		
<pre>Imep+Clom(PPI)</pre>	0.05+0.5	E /04						-
	0.05+0.5	5/24	None		-	Sethoxydm	0 0	
NO-TILL, 30-INCH	ROWS					occitoxydiii	0.2	6/22
Imazethapyr(EPP)	0.063	4./07						
Cyanazine(EPP)/	3/	4/27	Glyphosat	0.19	5/26	None		
Sethoxydim(Po)	0.15	4/27				Hone		-
Metribuzin(EPP)/	0.25/	6/13	Glyphosat	0.19	5/26	Acifluorf	0.05	
Sethoxydim(Po)	0.15	4/27				ACTITUOT	0.25	6/20
Metribuzin(Pre)/		6/13	Paraquat	0.5	5/26	Acifluorf	0.05	
Sethoxydim(Po)	0.2/	5/26			0, 20	Bentazon+	0.25	6/20
Sethoxydim(Po)/	0.15	6/13	Paraquat	0.5	5/26	Acifluor	0.75+	Mary F
Bent+Acif(Po)	0.2/	6/13	Sta mailing		0/20	ACTITION	0.13	6/20
HWC - Pend+	0.75+0.13	6/20	Paraquat	0.5	5/26	None		
Imep+Clom(EPP)	2/		Coll Hardan		3/20	None	1-3/3	M.D
Imepre roll (EPP)	0.05+0.5	4/27	Glyphosat	0.19	5/26	C-41 1		
O-TILL, 7-INCH R	0110			0.15	3/20	Sethoxydm	0.2	6/22
mazethapyr(EPP)								
yanazine(EPP)/	0.063	4/27	Glyphosat	0.19	5/26	Al-		
Satharydim (D-)	3/	4/27	57	0.15	3/20	None	- 1	
Sethoxydim(Po)	0.15	6/13	Glyphosat	0.19	E /2C	4 107		
etribuzin(EPP)/	0.25/	4/27	3 p.1103 Q C	0.15	5/26	Acifluorf	0.25	6/20
Sethoxydim(Po)	0.15	6/13	Paraquat	0 5	F /06	Bentazon+	0.75 +	
etribuzin(Pre)/	0.2/	5/26	· ui uquat	0.5	5/26	Acifluor	0.13	6/20
Sethoxydim(Po)	0.15	6/13	Paraquat	0 5	F 100	Bentazon+	0.75+	-, 20
ethoxydim(Po)/	0.2/	6/13	raraquat	0.5	5/26	Acifluor	0.13	6/20
Bent+Acif(Po)	0.75+0.13	6/20	Paraquat	0 -			Anna a	0, 20
IC - Pend+	A .	4/27	i ai aquat	0.5	5/26	None	-	_
mep+Clom(EPP)			Glyphosst	0.10				
ee Table 4 for i	nfammat:	n snrav	Jazoniquin	0.19	5/26	Sethoxydm	0.2	6/22

Continued on next four pages

postemergence (planned or as-needed) and burndown treatments were applied only as required and at a rate deemed necessary by the investigator. Rates and dates of all All 30-inch row plots were cultivated herbicide applications are given in Table 1. Broadleaf weeds taller than 6 inches and foxtail plants of any size found within a plot were counted just prior to harvest. harvested on October 7 and values adjusted to 12% moisture.

A follow-up treatment of sethoxydim was needed in minimum till plots even though trifluralin had been applied. Significant amounts of foxtail emerged despite trifluralin treatment probably because of the severe drought. conditions, a follow-up postemergence treatment for foxtail following trifluraling generally would not be required. Those treatments requiring a follow-up postemergence application of acifluorfen were infested with redroot pigweed and common lambsquarters while plots receiving acifluorfen plus bentazon had significant amounts of kochia in addition to pigweed and lambsquarters.

Among no-till treatments, early preplant cyanazine or imazethapyr provided essentially complete at-planting broadleaf control such that a low rate of glyphosate sufficed as a burndown to control emerged yellow foxtail. Treatments receiving early preplant metribuzin or no early preplant herbicide were treated with a preemergence burndown of paraquat due to infestations of kochia, lambsquarters,

When comparing tilled with no-till weed control systems, no-till yields were sigtail. nificantly higher than yields obtained with tillage (under 30-inch rows) (Table 2). Minimum till plots averaged 9.3 while no-till plots averaged 11.2 bushels per acre. Although herbicide treatments were not always the same between tillage system, the two treatments that were represented in both tillages ("total post" program and the handweeded check) showed about 2.5 bushels per acre higher yields under no-till. This tillage effect on grain yield was probably due to the moisture savings achieved The drought of 1988 was quite severe and yields of through avoidance of tillage. Under moisture conditions normally received in the Red River Valley, the yield advantage of no-till soybeans could potentially be greater. Fewer weeds were present at harvest in no-till than in minimum till. Paradoxically, this may reflect the fact that soil-applied treatments in minimum provided better broadleaf control than was obtained by no-till soil-applied treatments. As a consequence, most of the no-till treatments received a follow-up broadleaf spray while infestations in minimum till were considered too light to justify a follow-up spray given the capability of cultivation. Imazethapyr plus one cultivation controlled broadleaf weeds and foxtail sufficiently to preclude a follow-up herbicide treatment.

No yield differential was evident between 30-inch and 7-inch row spacings (Table 3). The 7-inch rows had more weeds (kochia, in particular) at the end of the season than did 30-inch rows. This probably was due to the lack of cultivation in the

narrow rows and the poor canopy closure due to severe drought.

Net returns (considering variable production costs only) tended to be higher for The option of cultivating probably gave a significant no-till than minimum till. advantage to no-till 30-inch rows by keeping herbicide costs or yield losses due to weeds to a minimum. Cultivation has not been traditionally practiced in no-till yet when no-tilling into small grain or other light stubble, it should be possible to cultivate no-till row crops without significant damage. Labor costs were not part of the variable cost rates used in this economic analysis. If included, the cost of labor would tend to decrease the profitability of minimum till more so than no-till. Treatments involving imazethapyr showed highest net return. This herbicide is not yet registered, though it is expected to be priced in the vacinity of \$15.00 acre, perhaps more. A cost of \$15.00 per acre was assumed for this analysis.

Table 2. Weed control and grain yield of minimum till and no-till soybeans.

Base Herbicide treat		Wee	ds nre	u no-1	it harv	ybean:	
	Rate	Yeft	KOCZ	Den.	it narv	est	Grain
TILLED (30-inch rows)	(1b/A)			s per	Colq 100 m <sup>2</sup>	)	yield (Bu/A)
Trifluralin+Metribuzin(PPI) Trifluralin+Imazethapyr(PPI) Trifluralin+Clomazone(PPI) Trifluralin+Chloramben(PPI) Seth(Po)/Bentazon+Acifluor(Po) Hand-weeded check	1+0.2 1+0.063 1+0.75 1+2.5 0.2/0.75+0.13	9 3 0 1 0	13 5 18 14 44	32 18 57 24 141	17 7 8 7 198	71 33 83 46 383	9.0 11.1 9.1 8.4 8.2
LSD 5% (Treatment within a till NO-TILL (30-inch rows)	lage)	NS	25	NS	NS	83	10.1 NS
Imazethapyr(EPP) Cyanazine(EPP)/Sethoxydim(Po) Metribuzin(EPP)/Sethoxydim(Po) Metribuzin(Pre)/Sethoxydim(Po) Seth(Po)/Bentazon+Acifluor(Po) Hand-weeded check	0.063 3/0.15 0.25/0.15 0.2/0.15 0.2/0.75+0.13	3 3 0 8 0	1 0 1 5 22	0 5 8 30 39	1 0 4 2 7	6 8 13 44 69	12.0 11.3 10.2 10.5 10.6
LSD 5% (Treatment within a till Iillage effect		NS	- NS	- NS	- NS	-	12.5
Handweeded check was treated w 0.75 + 0.05 + 0.5 lb/A plus has Handweeded check was		NS	*	NC	NC	30	NS

Handweeded check was treated with pendimethalin + imazethapyr + clomazone (EPP) at 1.5 + 0.05 + 0.5 lb/A plus hand weeding.

Table 3. Weed control and grain yield of no-till soybeans planted in 30-inch and 7-inch rows.

<u>Base Herbicide treat</u> Herbicide	Rate	Wee	ds pres KOCZ	ent	at harv	est	Grai
80-INCH ROWS (No-till)	(1b/A)		(Plants	Rrp	100 m <sup>2</sup>	)	yiel (Bu/A
mazethapyr(EPP)  yanazine(EPP)/Sethoxydim(Po)  etribuzin(EPP)/Sethoxydim(Po)  etribuzin(Pre)/Sethoxydim(Po)  eth(Po)/Bentazon+Acifluor(Po)  andweeded checka	0.063 3/0.15 0.25/0.15 0.2/0.15 0.2/0.75+0.13	3 3 0 8 0	1 0 1 5 22	0 5 8 30 39	1 0 4 2 7	6 8 13 44 69	12.0 11.3 10.2 10.5 10.6
SD 5% (Treatment within a till	age)	NS	NS	- NS	- NS	36	12.5 NS
-INCH ROWS (Tilled) mazethapyr(EPP) yanazine(EPP)/Sethoxydim(Po) etribuzin(EPP)/Sethoxydim(Po) etribuzin(Pre)/Sethoxydim(Po) eth(Po)/Bentazon+Acifluor(Po) andweeded check	0.063 3/0.15 0.25/0.15 0.2/0.15 0.2/0.75+0.13	6 13 11 12 1	4 2 17 29 67	2 24 36 63 207	4 0 5 4 17	15 39 68 108 292	11.4 10.4 11.6 10.7 9.0
D 5% (Treatment within a tilla <u>llage effect</u> andweeded check was treated w t 1.5 + 0.05 + 0.5 lb/A plus h		NS NS	42 *	NS NS	NS NS	210	10.0 NS NS

Table 4. Herbicide and adjuvant costs for minimum till and no-till weed systems.

Base tr	eatment		Burndown	treatme	nt	Follow-up pos Herbicide		CINCTIC
Hambicida			Herbicide	Ratea	Cost	or adjuvant	Ratea	Cost
or adjuvant	Ratea	(\$/A)	or adjuvant	Nace	(\$/A)			(\$/A)
		(4/1/						
TILLED, 30-IN	ICH ROWS		None	£30 n.		Sethoxydim	0.15	8.64
Trifluralin	1	6.38 5.13	None	23.04		Oil conc.	1 qt	1.25
Metribuzin	0.2	3.13				Cathanadim	0.15	8.64
Trifluralin	1	6.38	None		œ	Sethoxydim Oil conc.	1 qt	1.25
Imazethapyr	0.063	15.00				Off Conc.		
Trifluralin	1	6.38	None	-	•	Sethoxydim	0.15	8.64
Clomazone	0.75	12.00				Oil conc.	1 qt	1.25
		c 20	None			Sethoxydim	0.15	8.64
Trifluralin Chloramben	2.5	6.38 21.83	None			Oil conc.	1 qt	1.25
CITTOT AIIDEII	2.5		9			None		111112
Sethoxydim	0.2	11.52	None	-		Hone		
Oil conc.	1 qt 0.75	1.25 9.26						
Bentazon Acifluorfen	0.13	2.84						
X-77 surf.	0.5%	.62						
NO-TILL, 30-	THEL DE	NAC .						
Imazethapyr	0.063	15.00	Glyphosate	0.19	3.78	None	-	
Imazoonapj.			X-77 surf	0.25%	.31			
Companies	3	13.88	Glyphosate	0.19	3.78	Acifluorfen	0.25	5.69
Cyanazine Sethoxydim	0.15	8.64	X-77 surf	0.25%	6 .31	X-77	0.5%	.62
Dash	1 qt	1.63						
Matwibuzin	0.25	6.41	Paraquat	0.5	10.42	Acifluorfen	0.25	5.69
Metribuzin Sethoxydim	0.15	8.64	X-77	0.5%	.62	X-77	0.5%	.62
Dash	1 qt	1.63						
Makadharda	0.2	5.13	Paraquat	0.5	10.42	Bentazon	0.75	
Metribuzin Sethoxydim	0.15	8.64	X-77	0.5%		Acifluorfen	0.25	
Dash	1 qt	1.63				X-77	0.5%	
Cakhannadam	0.2	11.52	Paraquat	0.5	10.42	None		
Sethoxydim Oil conc.	1 qt	1.25	X-77	0.5%				
Bentazon	0.75	9.26						
Acifluorfen	0.13	2.84						
X-77 surf.	0.5%	.02						
NO-TILL, 7-	INCH RO	<u>DWS</u>		h 100110	avcont :	motrihuzin ear	v prer	lant:
All treatm	nents sa	ame as no	-till, 30-1nc	II rows	except	metribuzin ear		
Metribuzin	0.2	5.13	Paraquat		10.42	Bentazon	0.75	
Sethoxydim	0.1	8.64	X-77	0.59	6 .62	Acifluorfe	0.25 0.59	
Dash	1 q						<b>3 4 3</b> /	

Table 5. Economic analysis for weed control systems in minimum till and no-till soybeans.

Base herbicide	treatment		<u>riable produ</u> Herbicide	uction co	sts		
Herbicide	Rate	Total chemical	appl. and	Culti-	Chisel	Crop	Net
	(1b/A)	(\$/A)	incorp. (\$/A)	vation	plowing	value	returi
TILLED, 30-INCH	DOUG		(4/1/	(\$/A)	(\$/A)	(\$/A)	(\$/A)
Trif+Metr(PPI)							
Trif+Imep(PPI)	1+0.2	21.40	6.04	2.00	2 45		
Trif+Clom(PPI)	1+0.063	21.38	6.04	2.00	3.45	58.50	25.6
Trif+Clam(PPI)	1+0.75	28.27	6.04	2.00	3.45	72.15	39.28
Seth(Po)/	1+2.5	38.10	6.04	2.00	3.45	59.15	19.39
Bent+Acif(Po)	0.2/		3.04	2.00	3.45	54.60	5.01
Delictricii (PO)	0.75+0.13	25.49	1.24	2.00	0 4=		
NO-TILL, 30-INCH	I DOLLO		Maria maka	2.00	3.45	53.30	21.12
Imazethapyr(EPP)							Harry Carl
Cyanazine (EPP)/		19.09	1.86	2.00			
Sethoxydim(Po)	3/		-100	2.00	0	78.00	55.05
Metribuzin(EPP)/	0.15	34.55	2.48	2 00	de la comia de la		
Sethorydim(EPP)/			2.40	2.00	0	73.45	34.42
Sethoxydim(Po)	0.15	34.03	2.48	0.00			
Metribuzin(Pre)/	0.2/		2.40	2.00	0	66.30	27.79
Sethoxydim(Po)	0.15	42.01	1.86				-1.13
Sethoxydim(Po)/	0.2/		1.00	2.00	0	68.25	22.38
Bent+Acif(Po)	0.75+0.13	36.53	1.24				22.50
IO TTI		33.05	1.24	2.00	0	68.90	29.13
10-TILL, 7-INCH F	ROWS						23.13
mazethapyr(EPP)	0.063	19.09	1 00				
yanazine(EPP)/	3/	13.03	1.86	0	0	74.10	53.15
Sethoxydim(Po)	0.15	34.55	0.40				55.15
etribuzin(EPP)/	0.25/	34.33	2.48	0	0	67.60	30.57
Sethoxydim(Po)	0.15	42.01	0.40			07.00	30.5/
etribuzin(Pre)/	0.2/	42.01	2.48	0	0	75.40	30.91
ethoxydim(Po)	0.15	42.01			3,41	, 5, 40	30.91
ethoxydim(Po)/	0.2/	42.01	1.86	0	0	69.55	25 00
Bent+Acif(Po)	0 75+0 12	26 52					25.68
ariable cost re	stoc doust	36.53	1.24	0	0 1	58 E0	20 70
milliesota value	Day tan see	Trom Univer	sity of Min	nesota va	lues rec	ducad hu	20./3
				oment).	Included	in up	30%
				ance, and	fuel	labor :	lable
as \$2.40/A per p 988 soybeans wer	ing cost was	the per t	peracion,	herbicid	A incor-	Labor i	s not
988 soybeans wer	ass, cuitiva	tion cost wa	S \$2.00/A		- meorp	oration	cost

Carryover injury to wheat by clomazone applied in soybeans. Casselton, 1988. Fall and spring treatments were applied October 17, 1986 and June 3, respectively, using bicycle wheel sprayer delivering 17 gal/A with 8002 nozzles and 40 psi. All plots were double-pass incorporated (field cultivator, 3-inch tillage depth) immediately after herbicide application except for the minimum till preemergence treatments which received one tillage pass before spraying and no tillage after spraying. Which received one tillage pass before spraying and no tillage after spraying. McCall' soybeans were planted in 22-inch rows on June 4. On June 5, the entire experimental area received metribuzin plus alachlor (0.2 + 3 lb/A) for general weed control. Soybeans were bulk-harvested October 2, 1987. Treatment tillage was done on October 22, 1987 using a moldboard plow (6-inch depth) for plowed plots and a on October 22, 1987 using a moldboard plow (6-inch depth) for plowed plots and a on October 22, 1987 using a moldboard plow (6-inch depth) for plowed plots and a language plow (25 to 40% residue remaining) for minimum till plots. On April 14, chisel plow (25 to 40% residue remaining) for minimum till plots. On April 14, chisel plow (25 to 40% residue remaining) for minimum till plots. On April 14, chisel plow (25 to 40% residue remaining) for minimum till plots. On April 14, chisel plow (25 to 40% residue remaining) for minimum till plots. On April 14, chisel plow (25 to 40% residue remaining) for minimum till plots. On April 14, chisel plow (25 to 40% residue remaining) for minimum till plots. On April 14, chisel plow (25 to 40% residue remaining) for minimum till plots. On April 14, chisel plow (25 to 40% residue remaining) for minimum till plots. On April 14, chisel plow (25 to 40% residue remaining) for minimum till plots. On April 14, chisel plow (25 to 40% residue remaining) for minimum till plots. On April 14, chisel plow (25 to 40% residue remaining) for minimum till plots. On April 14, chisel plow (25 to 40% residue remainin

	Wh	oat in	iurv	1988 <sup>a</sup>	Whe	at gra	in yield	1988 <sup>a</sup>
Clomazone rate (lb/A)	Plow	Min. PPI	till Pre (%)	No-till	Plow	Min. PPI	Pre -(bu/A)	No-till
0	0	0	0	0	15	22	20	21
0.5 0.75	-	5		- 22 8		21	es es	
1.0	-	22	6		•	20	-	
1.5	10	48	11	8	18	15 22	20	25 16
0.75 1.0	25 39	22 38	21 23	24 24	17	21	24 24 19	27 14
1.25	58	66	53	42	11 5	11 13	10 13	17 12
			13 -				- 11	
	rate (1b/A) 0 0.5 0.75 1.0 1.25 1.5 2.0 0.5 0.75 1.0 1.25 1.5 2.0	Clomazone rate Plow (lb/A)  0 0 0.5 0.75 1.0 1.25 1.5 2.0 0.5 10 0.75 1.0 39 1.25 1.5 58	Clomazone rate Plow PPI (lb/A)  0 0 0 0.5 - 2 0.75 - 5 1.0 - 13 1.25 - 22 1.5 - 25 2.0 - 48 0.5 10 11 0.75 25 22 1.0 39 38 1.25 46 46 1.5 58 66 2.0 77 75	Clomazone rate Plow PPI Pre (1b/A)	rate         Plow         PPI         Pre         No-till           (lb/A)        (%)           0         0         0         0           0.5         -         2         -           0.75         -         5         -           1.0         -         13         -           1.25         -         22         -           1.5         -         25         -           2.0         -         48         -           0.5         10         11         11         8           0.75         25         22         21         24           1.0         39         38         23         24           1.25         46         46         43         29           1.5         58         66         53         42           2.0         77         75         73         58	Min. till   Plow   PPI   Pre   No-till   Plow	Min. till   Plow   PPI   Pre   No-till   Plow   PPI   PPI     PPI   PP	Clomazone rate Plow PPI Pre No-till Plow PPI Pre (1b/A)

<sup>&</sup>lt;sup>a</sup>Tillage effect was not significant for wheat injury and grain yield; rate effect was significant for both wheat injury and grain yield.

Summary. Severe drought in 1988 limited grain yield potential and probably masked much of the effect of clomazone carryover on yield. Generally, grain yield in 1988 seemed to be reduced by clomazone applied in the spring of 1987 at rates of 1.25 lb/A and above. Such was not the case, however, for clomazone applied in the fall of 1986 (done only in minimum till with preplant incorporated clomazone). Fall-applied clomazone likewise caused less carryover injury to wheat than did spring-applied treatments. Carryover injury with spring-applied treatments ranged between about 10% for a 0.5 lb/A application rate to about 60 to 75% for a 2 lb/A rate of clomazone.

Carryover injury to wheat by clomazone applied in soybeans, Carrington, 1988. Fall and spring treatments were applied October 30, 1986 and May 29, respectively, using a bicycle wheel sprayer delivering 17 gal/A with 8002 nozzles and 40 psi. All plots were double-pass incorporated (field cultivator, 3-inch tillage depth) immediately after herbicide application except for the minimum till preemergence treatments after herbicide application except for the minimum till preemergence treatments which received two tillage passes before spraying and no tillage after spraying. 'McCall' soybeans were planted in 30-inch rows on May 30. On June 5, the entire experimental area received metribuzin plus alachlor (0.2 + 3 lb/A) for general weed control. Soybeans were bulk-harvested October 2, 1987. Treatment tillage was done on April 13, 1988 using a moldboard plow (6-inch depth) for plowed plots and a field cultivator (40 to 60% residue remaining) for minimum till plots. 'Stoa' wheat was planted 1.5 inches deep on April 13 at 63 lbs/A. The entire experimental area replanted 1.5 inches deep on April 13 at 63 lbs/A. The entire experimental area received diclofop plus DPX-M6316 (1 + 0.031 lb/A) for control of foxtail and broadleaf weeds. On May 5, 135 1bs/A of N was surface applied as urea. Visual estimates of percent wheat injury (yellowing and bleaching of foliage) were taken on May 19 when the crop was 2 to 2.5-leaf and plots were combine harvested on August 4. Plot size was 12.5 by 25 ft and the experiment was a randomized complete block design with a split-plot arrangement of treatments. Soil type was a loam with pH 6.3 and 3.0%

Clomazone applica-	Clomazone	Wh	eat i	njury till	1988 <sup>a</sup>	Wh	eat gr	ain yiel	d 1988
tion timing	rate (1b/A)	Plow	PPI	Pre (%)	No-till	Plow	Min PPI	. till Pre -(bu/A)-	No-til
Fall 1986	0	0 -	0	0	0	25	23	24	27
	0.75	-	0		9 - 70	- 1	17 21	16-6	(91 <del>-</del> /3
	1.25 1.5 2.0		0	•	- 101	•	25 22 20	- 1911 - 1911	080 <del>1</del> 48 1080 <del>1</del> 42
Spring 1987	0.5 0.75 1.0	1 2 5	3 0 1 4	1 3 4	1 1	27 25	26 20 21	25 23	26 22
	1.25 1.5 2.0	12 11 18	1 2 9	10 6 19	2 8 5 8	26 26 24 26	27 23 24 23	22 24 28	28 24 21
SD(5%) Ratex		-	6	alts 51			23	24 NS	20

<sup>&</sup>lt;sup>a</sup>Rate and tillage effects were both significant for wheat injury.

Severe drought in 1988 limited wheat yield potential and may have masked any effect of clomazone carryover on yield. Wheat yields were not affected by clomazone applied from 12 to 19 months earlier. Wheat injury by clomazone was generally greater with higher clomazone application rates, although percent chlorosis never exceeded 20% even at the 2 lb/A rate of clomazone. Greatest injury was apparent where moldboard plowing preceded planting of spring wheat. clomazone produced almost no visually detectable wheat injury.

Fall and EPP imazethapyr and BAS514 in no-till soybeans, Minot, 1988. Fall (F), early preplant (EPP), and preemergence (Pre) treatments were applied October 14, 1987, April 18, and May 18, 1988, respectively, using a bicycle wheel sprayer delivering 17 gpa with 8002 nozzles and 40 psi. At the early preplant application, kochia was 0.25 to 0.38-inch tall and greenflower pepperweed had 1 to 2-inch diameter rosettes. McCall soybeans were seeded 1.5-inches deep into durum stubble in 30inch rows on May 19 using a Buffalo Till planter. Weed control was evaluated immed-The entire experimental area was treated immediately after planting with glyphosate at 1 lb/A to control existing vegetation. Visual estimates of percent soybean injury and weed control were taken July 13. a loam having a pH of 6.2 and 2.6% organic matter. Plot size was 10 by 25 ft and the experiment was a randomized complete block design having four replications.

the experiment wa	is a randon	11Zeu	Comp									
		At planting			8	<mark>8 weeks after planting</mark> Z Grft Ruth Rrpw Colq Wimu				Udanii		
	2000	VOC7	Caft	Colq	Wimu	Gfpw	KOCZ	Grft	Ruth	Rrpw	Colq	Wimu
<u>Treatment</u> <sup>a</sup>	Rate	KOCZ	Gric	COTY	WILLIA	(%	conti	rol)				
	(1b/A)				Mary a	(,,				100	100	98
	0.000	91	37	99	100	17	100	82	87	100	100	
Imazethapyr(EPP)	0.063		71	96	100		91	86	92	100	100	100
Imazethapyr(EPP)	0.074	84	/1	-	100		87	95	68	100	100	100
Imazethapyr(Pre)	0.063	-	-	98	95	30	100	79	100	100	100	100
Imazethapyr+Pend(F	) 0.063+2	99	84		100	43	100	83	99	100	100	100
Imazethapyr+Pend(F	) 0.074+2	99	87	100		78	100	97	100	100	100	100
Imep+Pend(EPP)	0.063+2	97	98	99	100	94	100	94	99	100	100	100
Imep+Pend(EPP)	0.074+2	99	98	100	96		100	99	79	100	99	100
Imep+Pend(Pre)	0.063+2		-		-	-	100	98	99	53	90	8
Imep+rend(rie)	0.75	32	100	62	32	-	100	98	99	59	85	19
BAS514(EPP)	1	82	100	92	62	27	100		97	87	98	23
BASS14(EPP)	1.5	55	100	98	46	14	100	100	70	85	95	28
BAS514(EPP)	i		-	-	•	-	52	99		100	100	100
BASS14(Pre)	0.75+0.063	98	92	98	97	36	on .	82	100	100	100	100
BAS514+Imep(F)	0.5+0.047	93	100	95	97	-	100	95	100		100	98
BAS514+Imep(EPP)	0.370.047	87	100	97	99	-	-	98	99	100		100
BAS514+Imep(EPP)	0.75+0.063	-	-	BENEF.	-	-	66	100	81	100	100	
BAS514+Imep(Pre)	0.75+0.063	100	99	99	99	63	100	93	99		100	
BAS514+Cyan(F)	1+2	95	100			0.	100	99	100			
BAS514+Cyan(EPP)	1+2								94			
BAS514+Metr(EPP)	0.75+0.3	96	100	100	100		0		0	0	0	0
Control	0			8					W. Barrier		r	12
		17	10	7	17	47	12					13
C.V. %		21	13	100		29	15				. 6	5 14
LSD 5%	3 1 2	f hc+					ibuzi	n wer	e use	ed.		

<sup>a</sup>Dry flowable formulations of both cyanazine and metribuzin were used.

Treatments involving BAS514 and imazethapyr appeared to provide less than complete at-planting kochia control when applied EPP but fall applications generally The addition of a surfactant or oil concentrate to these EPP treatments probably would have improved performance since the kochia population was at least partially emerged at the time of application. All EPP applications of BAS514 gave 100% control of green foxtail at planting. Mid-season evaluations showed that BAS514 provided excellent control of Russian thistle, kochia, and foxtail, good to excellent control of common lambsquarters, good control of redroot pigweed, and poor control of wild mustard. Imazethapyr gave excellent control of pigweed, mustard, and lambsquarters, good to excellent control of kochia, and about 85% control of foxtail and Russian thistle. For broadleaf weeds, EPP applications were generally superior to Pre treatments, although foxtail seemed to be controlled more effectively by Pre applications. Fall treatments generally were not as effective as spring applications. No soybean injury was observed with any of the treatments.

Fall and EPP imazethapyr and BAS514 in no-till soybeans, Carrington, 1988. Fall (F), early preplant (EPP), and preemergence (Pre) treatments were applied October 21, 1987, April 19, and May 19, 1988, respectively, using a bicycle wheel sprayer delivering 17 gpa with 8002 nozzles and 40 psi. Foxtail was primarily yellow with Maple Amber soybeans were seeded 1.5-inches deep into barley stubble in 30-inch rows on May 19 using a John Deere MaxEmerge planter. Foxtail control was evaluted prior to planting. The entire experimental area was treated immediately aftimates of percent soybean injury and foxtail control were taken on July 15. Soil timates a loam with 3.6% organic matter. Plot size was 10 by 25 ft and the experiment was a randomized complete block design having four replications.

		3	, repite	CIONS
Treatment		At planting	8 weeks	afte
Treatment	Rate	Fxtl	SBInj	
	(1b/A)	(%	control)	Fxt
Imazethapyr(EPP)		(70	control	
Imazethapyr(EPP)	0.063	49	0	62
Imazethapyr(Pre)	0.074	19	Ŏ	63
Imazethanyn (Pre)	0.063		Ö	57
Imazethapyr+Pendimethalin	(F) 0.063+2	86	0	72
Imazethapyr+Pendimethalin	(F) 0.074+2	91	Ö	82
Imazethapyr+Pendimethalin	(EPP) 0.063+2	96	0	87
Imazethapyr+Pendimethalin	(EPP) 0.074+2	96		85
Imazethapyr+Pendimethalin	(Pre) 0.063+2	-	0	85
BAS514(EPP)	0.75	57		92
BAS514 (EPP)	1	33	0	56
BASS14(EPP)	1.5	64	0	67
BASS14(Pre)	1	-	0	71
BAS514+Imazethapyr(F)	0.75+0.063	54		58
BAS514+Imazethapyr(EPP)	0.5+0.047	26	0	45
AS514+Imazethapyr(EPP)	0.75+0.063	64	0	67
AS514+Imazethapyr(Pre)	0.75+0.063	-	0	75
ASSI4+LVanazine-DF/FI	1+2	55	0	68
ASSIA+LVanazine-DF/FDD1	1+2	69	0	43
ASSI4+Metribuzin-DF(FPP)	0.75+0.3	58	0	80
ontrol	0	30	0	61
V a	and the same and		0	0
.V. %		26		1503
SD 5%		22	0	18
No treatments provided	to hear he was		NS	17

Summary. No treatments provided satisfactory control of foxtail at planting time except imazethapyr plus pendimethalin. These combinations along with BAS514 + cyanazine (EPP) were the only treatments giving 80% or greater foxtail control 8 weeks after planting. Treatments applied in the fall generally provided inferior foxtail control than EPP or Pre treatments. No clear difference was observed between the performance of EPP and Pre treatments. None of the treatments appeared to injure soybeans. Foxtail plants present in the plots at the July 15 evaluation were almost entirely yellow foxtail. Though control of green and yellow foxtail trolled more readily than yellow foxtail by all treatments.

Early preplant cyanazine in no-till soybeans, Fargo, 1988. Early preplant treatments were applied April 21 (4WPP = 4 weeks preplant), April 28 (3WPP), and May 5 (2WPP), and preemergence (Pre) treatments were applied May 20 using a bicycle wheel sprayer delivering 17 gpa with 8002 nozzles and 40 psi. Kochia at the 4WPP, 3WPP, and 2WPP timings was 0.25 to 0.38 inch, 0.5, and 0.5 inch tall, respectively. Postemergence treatments of sethoxydim were applied June 16 when air temperature was 70 F and relative humidity was 35%. McCall soybeans were seeded 1.5-inches-deep into a trashy surface in 30-inch rows on May 20 using a Hiniker no-till planter. The entire experimental area was treated immediately after planting with paraquat plus diquat at 0.5 + 0.5 lb/A to control existing vegetation. Soil type was silty clay having a pH of 7.8 and 5% organic matter. Visual estimates of percent soybean injury and weed control were taken on May 20 immediately before planting and on July 8. Plot size was 10 by 25 ft and the experiment was a randomized complete block design having four replications.

laving four replications.				1 1.1	
		Eval.	Evaluat		<u>y 8</u>
		May 20	Soybean	Voft	VOC7
Turnetment a	Rate	KOCZ	injury	Teru	11
Treatment	(1b/A)	(% i	njury or	contro	1)
		0.4	0		94
Cyan-L+OC(4WPP)/Seth+OC(Po)	2.5+10/0.2+10	94	Ö	94	98
Cyan-L+OC(4WPP)/Seth+OC(Po)	3+10/0.2+10	99		27	100
Cyan-L+OC(4WPP)/Seth+OC(Po)	3.5+10/0.2+10	99	0	85	84
Cyan-L+OC(3WPP)/Seth+OC(Po)	2.5+10/0.2+10	96	0	00	04
Cyan-L+OC(3WPP)/Seth+OC(Po)	3+10/0.2+10	100	0		
Cyan-L+OC(3WPP)/Seth+OC(Po)	3.5+10/0.2+10	99	0	07	0.5
Cyan-L+OC(3WPP)/Seth+OC(Po)	2.5+10/0.2+10	99	0	87	95
Cyan-L+OC(2WPP)/Seth+OC(Po)	3+10/0.2+10	100	0	95	98
Cyan-L+OC(2WPP)/Seth+OC(Po)	3.5+10/0.2+10	99	0	88	100
Cyan-L+OC(2WPP)/Seth+OC(Po)	2.5+10/0.2+10	-	0	ca ca	97
Cyan-L+OC(Pre)/Seth+OC(Po)	3+10/0.2+10	-	0	88	99
Cyan-L+OC(Pre)/Seth+OC(Po)	3.5+10/0.2+10		0	-	100
Cyan-L+OC(Pre)/Seth+OC(Po)	3.5+14/0.2+14	Crestal Inc	0	0	0
Control	U		A STATE OF THE SECOND		
		1	0	7	6
C.V. %		2	NS	8	8
LOD FO	ion of cyanazina	· 0C = r	etroleum	oil co	ncen-
aCvan-L = flowable formulat	Toll of Cyallazine	, oo r	cre		

Cyan-L = flowable tormulation of trate containing 17% emulsifier; 10 = 1 quart per acre.

Summary. Complete preplant burndown of kochia was achieved by 3 to 3.5 lb/A of cyanazine flowable mixed with a petroleum oil concentrate. Cyanazine at 2.5 1b/A allowed a low percentage of kochia escapes when applied at the 4WPP and 3WPP timings. Drought conditions prevailed during these preplant treatments and kochia densities were high enough to completely cover the ground surface. Thus, excellent preplant burndown by cyanazine was obtained despite dense, drought-stressed plants. Cyanazine also provided excellent mid-season kochia control with the exception of the 84% control observed with cyanazine at 2.5 lb/A applied 3WPP. No soybean injury was detected.

Early preplant treatments in no-till soybeans, Fargo, 1988. Early preplant (EPP) treatments were applied April 21 when kochia was 0.25 to 0.38-inch tall in a dense to untilled soybean stubble using a Hiniker no-till planter. Visual estimates of treatments were applied May 20. All treatments were applied with a bicycle wheel sprayer delivering 17 gal/A with 8002 nozzles and 40 psi. On May 23, the entire exexisting vegetation. Soil type was silty clay having pH 7.9 and 5% organic matter. Estimates of percent crop injury and weed control were obtained on June 30. Plot ing four replications.

Treatment <sup>a</sup>	the said party	Eval. 5/20	Eval	uated	June	30
Treatment.	Rate <sup>a</sup>	KOCZ	Soybea injury		7 0	
	(1b/A)		injury	KOC:	2 Krp	w Ye
Imazethapyr+X77(EPP)	0.002.0.054		· · · · · · · · · · · · · · · · · · ·	01 (1	ontro	1)
lmazethapyr+X77(Fppi	0.063+0.25%	68	1	100	100	8
Mazethanvr(Pro)	0.078+0.25%	73	3	100	100	8
mep+Pendimethalin+Y77/FDD\	0.063	1113-	13	100	100	9.
IIICUTPENGIMETHALIALY77/FDD	0.063+2+0.25%	80	0	98	100	9
maze inapyr+pendimethalin/pun)	0.078+2+0.25%	83	0	100	99	94
722144D420A0(Fbb)	0.063+2		3	100	100	9!
AS514+BAS090(FPP)	0.75+10	-	0		100	3
AS514+BAS090(FPP)	1+10	64	1	95	36	70
ASS14(Pre)	1.5+10	55	3	97	79	
mazethapyr+BAS514+BAS000(EDD)	1	-	0	99	85	49 72
IIQ LE LII dDV T+KANN I A L RACAGA / TANN	0.047+0.5+10	87	0	99	99	74
maze unabyr+BAS514 (Dro)	0.063+0.75+10	85	3	98	100	92
erripuzin+Pendimethaliniv77/cnn	0.047+0.5	-	4	98	99	91
or IDUA INTERNATIONALISM (Due)	0.25+2+0.25%	85	0	47	13	8
"IDUZIN+Metolachlor/Drol	0.25+2	-	0	90	98	47
.3340Z+UC(EPP)	0.25+2.5	-	1	92	86	39
53482(Pre)	0.066+10	70	0	20	6	
53482+Pendimethalinioc(CDD)	0.066	-	7	19	54	1 3
JJ402+PENNIMAThaliniOC(FDD)	0.044+2+10	69	0	3	1	13
JJ402+rengimethalin/Dwol	0.066+2+10	70	0	26	11	13
53482+Pendimethalin(Pre)	0.044+2	-	0	-	11	13
ntrol	0.066+2	-	0	-		
V. %	0	•	0	0	0	6
D 5%		18	264	15		
ASO90 = BASF surfactant of prop f metribuzin was used; OC = petr			204	15	8	38

'BASO90 = BASF surfactant of proprietary composition; dry flowable formulation of metribuzin was used; OC = petroleum oil concentrate containing 17% emulsifier; 10 = 1 quart per acre.

Summary. None of the early preplant treatments provided kochia control at planting time sufficient to alleviate the need for a burndown herbicide. Control of kochia and redroot pigweed was essentially 100% for imazethapyr treatments when evaluated on June 30. BAS514 similarly provided 95 to 100% kochia control but only about 80 to 85% control of pigweed. S-53482 gave poor control of these broadleaf species. Treatments yielding good to excellent yellow foxtail control involved imazethapyr alone or in combination with either BAS514 or pendimethalin. Over 5% soybean injury was observed with imazethapyr and S-53482 treatments but these results were not consistent across comparable treatments and injury symptoms may have been complicated by drought.

Burndown treatments for no-till soybeans, Carrington, 1988. Maple Amber soybeans were planted 1.5 inches deep into barley stubble on May 19 using a John Deere Maxtemperature. Burndown treatments were applied after planting on May 19 when air temperature was 82 F with cloudy skies and 65% relative humidity. Surface soil was temperature was 82 F with cloudy skies and 65% relative humidity. Surface soil was temperature was 82 F with cloudy skies and 65% relative humidity. Surface soil was follows: foxtail (yellow with 20 to 30% green foxtail) 1 to 3-leaf, 0.25 to 1-as follows: foxtail (yellow with 20 to 30% green foxtail) 1 to 3-leaf, 0.25 to 1-as follows: foxtail (yellow with 20 to 30% green foxtail) 1 to 2-inch tall; common lambsquarinch tall; horseweed 1.5 to 3-inch tall; kochia 1 to 2-inch tall; common lambsquarinch tall; several-leaf; wild buckwheat 2 to 3-leaf, 1 to 2-inch tall. All plots received 0.2 lb/A sethoxydim plus 1 qt/A petroleum oil concentrate on June 7 when temperature was 93 F, RH was 35%, and foxtail was 2 to 4 inches tall in plots when temperature was 93 F, RH was 35%, and foxtail was 2 to 4 inches tall in plots when temperature was 93 F, RH was 35%, and foxtail was 2 to 4 inches tall in plots when temperature was 93 F, RH was 35%, and foxtail was 2 to 4 inches tall in plots when temperature was 93 F, RH was 35%, and foxtail was 2 to 4 inches tall in plots when temperature was 93 F, RH was 35%, and foxtail was 2 to 4 inches tall in plots when temperature was 93 F, RH was 35%, and foxtail was 2 to 4 inches tall in plots when temperature was 93 F, RH was 35%, and foxtail was 2 to 4 inches tall. All temperature was 93 F, RH was 35%, and foxtail was 2 to 4 inches tall; was 10 F, RH was 10 F,

complete block de	esign navin	y rou	ТСР					Evalu	ated	July	15	
Burndown treatment <sup>a</sup>	Rate (1b/A)	E Fxtl	valua Howe	ted J KOCZ	Cala	Wihw	Soybean injury ury or	n Fxtl	KOCZ			
Acif+2,4-D	.125+.5	68	96	87	100	100	0	73	92	98	100	97
+Dash+28%UAN Acif+2,4-D	+1Q+1G .25+.5	77	96	100	100	100	0	70	95	99	99	98
+Dash+28%UAN Seth+2,4-D	+1Q+1G .1+.5	90	95	94	100	80	0	73	90	74	100	99
+Dash+28%UAN	+1Q+1G			98	100	99	0	66	95	98	100	96
Seth+2,4-D+Acif +Dash+28%UAN	+1Q+1G	03	95			99	0	69	94	98	100	99
Acif+2,4-D +OC+28%UAN	.125+.5 +1Q+1G	67	97	97	100		0	73	91	58	100	100
Seth+2,4-D +OC+28%UAN	.1+.5 +1Q+1G	88	95	84	99	77			96	97	100	99
Seth+2,4-D+Acif	.1+.5+.125	88	96	99	100	99	0	69				97
+0C+28%UAN Glyt&2,4-D+AS	.28&.25+2.	5 89	98	100	98	97 0	0	62 93	99 7	99 21	99	
Control	0	0	0 2	11	1	1	0	9	6	6	2	
C.V. %		8	2	13	i	2	NS	10	8 4 D	8 - 1 an	<u>3</u> dmast	
LSD 5%	hvl ester f	ormula	ation	of 2	,4-D	was u	sed;	Tyt&2	,4-U			oncen-

The butoxyethyl ester formulation of 2,4-D was used; Glyt&2,4-D = Landmaster II herbicide; 28%UAN = 28% urea ammonium nitrate solution; OC = petroleum oil concentrate containing 17% emulsifier; AS = ammonium sulfate; Dash = BASF surfactant of

Summary. No soybean injury was evident at the June 7 evaluation. Acifluorfen at 0.125 or 0.25 lb/A did not provide foxtail burndown that was equivalent to that provided by sethoxydim or glyphosate. Foxtail burndown by sethoxydim or glyphosate was only about 90%, probably because of barley stubble and chaff which appeared to have shielded small foxtail seedlings from spray. Burndown of horseweed appeared to be primarily due to 2,4-D and was essentially complete. Burndown of lambsquarters was excellent for all treatments while effective burndown of wild buckwheat and kochia excellent for all treatments while effective burndown of mild buckwheat and kochia seemed to require acifluorfen in addition to 2,4-D. Fair to poor control of foxtail seemed to require acifluorfen foxtail control (93%) observed in the control June 7 sethoxydim treatment. Superior foxtail control (93%) observed in the control plots on July 15 was apparently due to competition from dense populations of the broadleaf species.

Cyanazine injury in no-till sunflowers and soybeans, Minot, 1988. Fall (F) treatments were applied October 14, 1987 and early preplant treatments were applied April 18 (4 weeks preplant = 4WPP), April 25 (3WPP), and May 3, 1988 (2WPP). Herbicides were applied with a bicycle wheel sprayer delivering 17 gpa with 8002 nozzles and 40 into durum stubble in 30-inch rows on May 17 using a Buffalo Till planter. Direction of planting was perpendicular to the plots and five rows of each crop were seeded into each plot. Soil type was a loam with pH 6.2 and 2.6% organic matter. Plus 2,4-D at 2.25 + 0.25 lb/A to control existing vegetation. Visual estimates of leaf and 5 inches tall and soybeans had 1 to 2 trifoliolate leaves. Plot size was replications.

<u>Treatment</u> <sup>a</sup>		Sunflower	Soybear
	Rate	injury	injury
	(1b/A)	(%	1
Cyanazine+Pendimethalin(2WPP) Cyanazine+Pendimethalin(3WPP) Cyanazine+Pendimethalin(2WPP) Cyanazine+Pendimethalin(2WPP) Cyanazine+Pendimethalin(3WPP) Cyanazine+Pendimethalin(F) Cyanazine+Pendimethalin(2WPP) Cyanazine+Pendimethalin(3WPP) Cyanazine+Pendimethalin(3WPP) Cyanazine+Pendimethalin(2WPP) Cyanazine+Pendimethalin(2WPP) Cyanazine+Pendimethalin(3WPP) Cyanazine+Pendimethalin(3WPP) Cyanazine+Pendimethalin(3WPP) Cyanazine+Pendimethalin(5WPP)	2.5+2 2.5+2 2.5+2 3+2 3+2 3+2 3.5+2 3.5+2 3.5+2 4+2 4+2 4+2 4+2	0 0 0 0 0 0 0 0 0	0 0 0 0 0 0 0 0 0 0 0
SD 5%		0	
Ory flowable formulation of cyana		NS	612 _NS

<u>Summary</u>. No injury attributable to cyanazine was observed. Low levels of rainfall experienced this year would be expected to cause only minimal movement of cyanazine into the soil profile and may explain the lack of crop injury.

Fall and EPP cyanazine and RE-40885 in no-till sunflowers, Minot, 1988. early preplant (EPP), and preemergence (Pre) treatments were applied October 14, 1987, April 18, and May 18, 1988, respectively, using a bicycle wheel sprayer delivering 17 gpa with 8002 nozzles and 40 psi. Greenflower pepperweed had 0.5 to 2-inchdiam rosettes and kochia was 0.25 to 0.38-inch tall at the time of EPP applications. Northrup King 265 sunflowers were seeded 1.5-inches deep into durum stubble in 30inch rows on May 17 using a Buffalo Till planter. Soil type was a loam with pH 6.2 and 2.6% organic matter. The entire experimental area was treated immediately after planting with glyphosate plus 2,4-D at 2.25 + 0.25 1b/A to control existing vegeta-Visual estimates of percent sunflower injury and weed control were taken on May 17 immediately before planting and on July 14. Plot size was 10 by 25 ft and the experiment was a randomized complete block design having four replications.

experiment was a ra	naomized	Comp	1600	D 1001								•	
									valua	ated J	luly 1	4	
<u>Treatment<sup>a</sup></u>	Rate (1b/A)	Grft	KOCZ	112	lay 17 Colq sunflo	CENI	Sufl inj injur	Grft	KOCZ weed	Wimu	Ruth rol)	Rrpw	Colq
Cyanazine+Pend(F) Cyanazine+Pend(F) Cyanazine+Pend(EPP Cyanazine+Pend(EPP Cyanazine+Trif(F) RE-40885+Pend(F) RE-40885+Pend(EPP)	2.5+2 3+2.5 ) 2+2 ) 2.5+2 ) 3+2.5 2.5+1.2 1+2 1.25+2. 1+1.5 1+2 1.25+2.	78 5 84 97 99 5 98	100 100 100 100 100 100 97 96 95 97 95 71	99 100 100 100 100 99 99 100 100 100	100 100 100 100 100 100 96 96 100 100 65	48 68 61 50 62 - 16 - 37 40 -	0 0 0 0 0 0 0 0 0 0 0 0 0 0	63 70 91 88 94 44 - 47 68 77 83 82 67 0	99 100 100 97 - 97 - 69 92 94 100 - 24 0	100 - 98 - 100 - 83 97 - 98 - 0	13 11 43 43 43 13 14 30 30 38 34 10 0	42 46 90 83 88 53 - 69 81 89 87 92 71 0	84 90 94 87 84 63 - 65 82 65 87 79 69
C.V. % LSD 5%  The dry flowable	formula	6 8	8 11	l NS anazi	14 19 ne was	36 26 s use	NS	20 19 the e	21 24 mulsi	11 13 fiable	79 28 e con	28 28 centr	26 <u>27</u> ate o

The dry flowar trifluralin was used.

Summary. Fall treatments of cyanazine plus pendimethalin or RE-40885 plus pendimethalin generally provided control inferior to that achieved with early preplant applications. A formulation of RE-40885 having inadequate mixing qualities was used for fall applications and may have adversely affected efficacy of these treatments. Fall-applied trifluralin (plus cyanazine) gave poor foxtail control. Essentially complete control of all species at planting time was achieved by several combinations of cyanazine plus pendimethalin or RE-40885 plus pendimethalin applied early preplant. These treatment combinations provided good to excellent mid-season control of broadleaf species except Russian thistle and mostly 80 to 90% control of foxtail. None of the treatments caused visually detectable sunflower injury.

Fall and EPP cyanazine and RE-40885 in no-till sunflowers. Carrington, 1988. Fall (F), early preplant (EPP), and preemergence (Pre) treatments were applied October 21, 1987, April 19, and May 19, 1988, respectively, using a bicycle wheel sprayer delivering 17 gpa with 8002 nozzles and 40 psi. Foxtail was primarily yellow with Interstate 897 sunflowers were seeded 1.5-inches deep into barley stubble in 30-inch rows on May 19 using a John Deere MaxEmerge planter. The entire experimental area was treated immediately after planting with glyphosate at 1 lb/A to control existing taken on July 15. Plot size was 10 by 25 ft and the experiment was a randomized complete block design having four replications.

<u>Treatment</u> <sup>a</sup>		Sunflower	
	Rate	injury	Fxt1
	(1b/A)	(%)	)
Cyanazine+Pendimethalin(F) Cyanazine+Pendimethalin(F) Cyanazine+Pendimethalin(EPP) Cyanazine+Pendimethalin(EPP) Cyanazine+Pendimethalin(EPP) Cyanazine+Trifluralin(F) RE-40885+Pendimethalin(F) RE-40885+Pendimethalin(EPP) RE-40885+Pendimethalin(EPP) RE-40885+Pendimethalin(EPP) RE-40885+Pendimethalin(EPP) RE-40885+Pendimethalin(EPP) RE-40885+Pendimethalin(EPP) Control	2.5+2 3+2.5 2+2 2.5+2 3+2.5 2.5+1.25 1+2 1.25+2.5 1+1.5 1+1.5 1+1.5 0	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	56 74 88 80 87 14 40 56 70 78 81 72 70 0
SD 5%		0	21
	L. PAS	NS	19
The dry flowable formulation the emulsifiable concentrate	n of cyana of trifle		used; used.

Summary. Fall applications provided fair to poor control of foxtail. A formulation of RE-40885 having inadequate mixing qualities was used for fall applications and may have adversely affected efficacy of these treatments. Fall-applied, non-incoreffective treatments in the study were mixtures of cyanazine plus pendimethalin applied early preplant. None of the treatments caused visually detectable sunflower experiment than yellow foxtail; nearly all the uncontrolled foxtail in the plots was yellow foxtail while the check strips between plots contained about 20 to 30% green foxtail

Early preplant treatments in no-till sunflowers, Fargo, 1988. Early preplant treatments were applied April 22 (6 weeks preplant = 6WPP), April 28 (5WPP), and May 5 (4WPP), preemergence (Pre) treatments were applied June 2, and postemergence (Po) (4WPP), preemergence (Pre) treatments were applied June 16 using a bicycle wheel sprayer delivering 17 gpa with treatments were applied treatments) or 8.5 gpa with 8001 nozzles (sethoxydim 8002 nozzles (soil-applied treatments) or 8.5 gpa with 8001 nozzles (sethoxydim 8002 nozzles (soil-applied treatments) sethoxydim was applied to 3 to 4-leaf foxtreatments. Spray pressure was 40 psi. Sethoxydim was applied to 3 to 4-leaf foxtrail (mostly yellow with some green foxtail) when air temperature was 70 F and relative humidity was 35%. Kochia was 0.5 to 1-inch tall at the 4WPP treatment but was not emerged for earlier treatments. Interstate 3001 sunflowers were seeded 1.5-inch energed for earlier treatments. Interstate 3001 sunflowers were seeded 1.5-inch energed into wheat stubble using a Hiniker no-till planter set at 30-inch rows and 22,000 seeds per acre. The entire experimental area was treated with paraquat plus diquat (0.5 + 0.5 lb/A) on June 2 to control existing vegetation. Soil type was a silty clay with pH 7.8 and 5% organic matter. Visual estimates of percent kochia control were taken on May 20 and plots were evaluated again on July 12. Plot size was 10 by 25 ft and the experiment was a randomized complete block design having four replications.

Eval
May 20 Sunfl. SF stand SF height         reatment <sup>a</sup> Rate       KOCZ injury reduction reduction Fxtl         (1b/A)      (%)(%)
reatment Rate KOCZ injury reduction Fxtl (1b/A)(%)(%)
reatment <sup>a</sup> Rate KOCZ INJURY reduction reads  (1b/A) (%)(%)
70 0 11 4 64
4 64
ar accor, Dandimothalanti/Inwrrl 176TV.5%
a = 4000 E and imperhal in (Pre) $a = 176$
nr 4000F (Dond 1 / /   DWPP)
0.75+2 - 0 0 44
RE-40003+reliata//(onit/
$C_{\text{vanazine}}(6WPP)/\text{Sethoxydim}(PO)$ 3/0.2 0 0 0 82
$C_{\text{vanazine}}(6\text{WPP})/\text{Sethoxydim}(PO)$ 3.3/0.2 3
$C_{\text{vanazine}}(SWPP)/Sethoxydim(PO)$ 3.3/0.2 3 5 8/
Cyanazine(4WPP)/Sethoxydim(Po) 3.5/0.2 100 0 3 32 84 Cyanazine(4WPP)/Sethoxydim(Po) 3.5/0.2 100 0 42 32 84
Cyanazine(Pre)/Sethoxydim(Po) $3.5/0.2$ 0 0 0 0
Control
271
20 0 23
13 NS 17 Leatherweigh word ar
LSD 5%  approx flowable formulation of cyanazine was used; cyanazine and sethoxydim were approx flowable formulation of cyanazine was used; cyanazine and sethoxydim were approx flowable formulation of cyanazine was used;

aDry flowable formulation of cyanazine was used; cyanazine and sethoxydim were applied with 1 qt/A of a petroleum oil concentrate containing 17% emulsifier.

Summary. Over 90% control of kochia at planting time was provided by early preplant applications of cyanazine, while similarly-applied mixtures of RE-40885 plus pendimethalin gave only 70% control. Reductions in sunflower stand and height were severe with preemergence applications of cyanazine and appeared to be evident with vere with preplant cyanazine treatments and the high rate of RE-40885 plus pendimethalin applied early preplant.

Fall-applied herbicides in chemical fallow, Minot, 1988. Treatments were applied October 14, 1987 into durum stubble using a bicycle wheel sprayer delivering 17 gal/A with 8002 nozzles and 40 psi. Soil type was a loam with pH 6.2 and 2.6% organic matter. Visual estimates of percent control were taken on July 14, 1988. Plot size was 10 by 25 ft and the experiment was a randomized complete block design having four replications.

<u>Treatment</u>	Rate	0. 61	We		trol	
	(1b/A)	Grft	KOCZ	Gfpw	Ruth	Cold
BAS-514		F 189		(%)-		
BAS-514	0.75	97	84	10		
BAS-514	1	97	68	19 13	92	51
BAS-514	1.5	99	99	19	94	78
BAS-514+Chlorsulfuron	2	100	99	15	96 100	92
BAS-514+Chlorsulfuron	0.75+0.015	86	100	35	99	93
BAS-514+Chlorsulfuron	1+0.015	94	100	51	100	100
BAS-514+CGA-131036	1.5+0.015	97	100	48	100	100
BAS-514+Atrazing_DE	1+0.015	90	99	35	95	78
3AS-514+Atrazine-DF	1+0.75 1.5+0.75	81	99	32	97	97
3A3-514+Atrazine-DF	2+0.75	96	100	51	97	97
hlorsulfuron	0.015	97	99	47	99	95
GA-131036	0.015	0	100	44	33	99
GA-131036	0.023	0	86	25	0	46
lomazone	0.75	Ö	98	44	0	65
lomazone	1	13	25	4	0	18
lomazone+Chlorsulfuron lomazone+Chlorsulfuron	0.75+0.015	3	74	0 31	0	23
AS-514+Clomazone	1+0.015	0	100	29	5	75
AS-514+Clomazone	1+0.75	91	93	23	77	100
ontrol	1.5+1	98	96	13	94	86
	0	0	0	0	0	95
. V. %					•	U
SD 5%		12	21	71	10	30
BAS-514 alone provide		10	24	28	9	32

Summary. BAS-514 alone provided nearly complete control of green foxtail at all rates evaluated but control was generally antagonized by mixing with other herbicides. BAS-514 alone at 1.5 and 2 lb/A provided acceptable control of kochia and Russian thistle although excellent control of these species at lower BAS-514 rates and excellent control of common lambsquarters required tank-mixing with either atracine or chlorsulfuron. Clomazone performed poorly, perhaps because of dry conditions prevailing through the duration of the experiment. None of the treatments controlled the winter annual greenflower pepperweed.

Longevity of soil-applied herbicides in fallow, Minot, 1988. Treatments were applied April 19 onto durum stubble using a bicycle wheel sprayer delivering 17 gpa tall and greenflower pepperweed had 1 to 2-inch-diameter rosettes. Soil type was a tall and greenflower pepperweed had 1 to 2-inch-diameter rosettes. Soil type was a tall and greenflower pepperweed had 1 to 2-inch-diameter rosettes. Soil type was a tall and greenflower pepperweed and 2.6% organic matter. On May 26, estimates of percent weed conloam with pH 6.2 and 2.6% organic matter. On May 26, estimates of percent weed conloam with pH 6.2 and 2.6% organic matter. On May 26, estimates of percent weed conloam with pH 6.2 and 2.6% organic matter. On May 26, estimates of percent weed conloam with pH 6.2 and 2.6% organic matter. On May 26, estimates of percent weed conloam with pH 6.2 and 2.6% organic matter. On May 26, estimates of percent weed conloam with pH 6.2 and 2.6% organic matter. On May 26, estimates of percent weed conloam with pH 6.2 and 2.6% organic matter. On May 26, estimates of percent weed conloam with pH 6.2 and 2.6% organic matter. On May 26, estimates of percent weed conloam with pH 6.2 and 2.6% organic matter. On May 26, estimates of percent weed conloam with pH 6.2 and 2.6% organic matter. On May 26, estimates of percent weed conloam with pH 6.2 and 2.6% organic matter. On May 26, estimates of percent weed conloam with pH 6.2 and 2.6% organic matter. On May 26, estimates of percent weed conloam with pH 6.2 and 2.6% organic matter. On May 26, estimates of percent weed conloam with pH 6.2 and 2.6% organic matter. On May 26, estimates of percent weed conloam with pH 6.2 and 2.6% organic matter. On May 26, estimates of percent weed conloam with pH 6.2 and 2.6% organic matter. On May 26, estimates of percent weed conloam with pH 6.2 and 2.6% organic matter. On May 26, estimates of percent weed conloam with pH 6.2 and 2.6% organic matter. On May 26, estimates of percent weed conloam with pH 6.2 and 2.6% organic matter. On May 26, estimates

Treatment <sup>a</sup>	Rate (1b/A)	Grft	valua KOCZ	Ruth	lay 26 Colq (	Gfpw %)	Eval. Grft	July KOCZ	
BAS514 BAS514 BAS514 BAS514+Chlorsulfuron BAS514+Chlorsulfuron BAS514+Chlorsulfuron Clomazone+Atrazine BAS514+Clomazone BAS514+Clomazone+Atra Control	1 1.5 2 1+0.015 1.5+0.015 2+0.015 0.5+0.5 1.5+0.5	100 100 100 100 100 100 28 100 100	74 69 71 93 92 94 100 90 100	98 100 100 98 98 100 100 99 100	87 96 98 100 100 100 100 100 100	30 33 28 71 80 77 57 37 51 0	98 98 100 96 99 100 54 99 100	100 100 100 100 100 100 100 100 100	50 85 91 100 100 100 91 91 99 0
C.V. %		2	5 6	2	3 4	31 21	9	NS	9 10

 $^{4}$ All treatments were applied with a petroleum oil concentrate containing 17% emulsifier; wettable powder formulation of atrazine was used.

Summary. BAS514 provided excellent control of green foxtail at both the May and the July evaluation dates. BAS514 also controlled Russian thistle and gave nearly complete control of common lambsquarters when applied at 1.5 to 2 lb/A. Redroot pigweed control by BAS514 was not adequate for fallow. Kochia was completely controlled by BAS514 at the July evaluation but not at the May evaluation. The only treatments giving complete kochia control in May were those containing 0.5 lb/A atrazine. The combination of clomazone plus atrazine (0.5 + 0.5 lb/A) provided complete control of kochia, Russian thistle, and common lambsquarters but failed to plete control of kochia, Russian thistle, and common lambsquarters but failed to only about 90% control of redroot pigweed which may not be adequate for a fallow only about 90% control of redroot pigweed which may not be adequate for a fallow situation. None of the treatments controlled the winter annual greenflower pepperweed.

BAS514 and clomazone for weed control in fallow, Page, 1988. Treatments were applied April 20 using a bicycle wheel sprayer delivering 17 gal/A with 8002 nozzles and 40 psi. Soil type was a loam with pH 7.4 and 8.6% organic matter. Soil surface was mostly bare with small amounts of barley residue. Visual estimates of percent green foxtail control were taken on June 7. Plot size was 20 by 25 ft and the experiment was a randomized complete block design having four replications.

<u>Treatment</u> <sup>a</sup>	Rate	Green foxtail control
	(1b/A)	(%)
BAS514		
BAS514	, 1	65
BAS514	1.5	67
BAS514+Chlorsulfuron	2	72
BAS514+Chlorsulfuron	1+0.015	62
BAS514+Chlorsulfuron	1.5+0.015	64
Clomazone+Atrazine	2+0.015	73
RASE14 Clama	0.5+0.5	37
BAS514+Clomazone	1.5+0.5	71
BAS514+Clomazone+Atrazine	1.5+0.5+0.5	74
Clomazone	0.75	40
Control	0	Ö
C.V. %		
.SD 5%		15
Wettable powder formulation		12

 $\underline{\text{Summary}}$ . None of the treatments provided acceptable control of foxtail. Conditions were very dry and the surface was quite powdery during April and much of May. Insufficent moisture for herbicide activation probably accounted for the failure of these treatments.

Postemergence treatments in fallow, Minot, 1988. Treatments were applied May 26 into durum stubble using a bicycle wheel sprayer delivering 8.5 gal/A with 8001 nozzles and 40 psi. At time of spraying (6 to 7:30 pm), air temperature was 82 F, relative humidity was 60%, and skies were sunny. Reasonably good soil surface moisture was present and weeds did not appear to be under stress. Weed stages at application were as follows: kochia was 0.5 to 3 inches tall, Russian thistle was 1 to 2.5 inches tall, and common lambsquarters was 0.5 to 2 inches tall. Visual estimates of percent control were taken on June 13. Plot size was 10 by 25 ft and the experiment was a randomized complete block design having four replications.

		Weed	contr	01
Ta	Rate	KOCZ	Ruth	
Treatment <sup>a</sup>	(1b/A)		- (%)	
219.0	0.38	98	100	91
Paraquat	0.5	99	100	98
Paraquat	0.38&0.06&0.02	99	100	95
Para&Flfp&Diur	0.5&0.08&0.027	98	100	98
Para&F1fp&Diur	0.38+0.1	98	100	94
Paraquat+Diuron	0.5+0.1	100	100	96
Paraquat+Diuron		99	100	99
Paraquat+Cyanazine	0.38+0.25	100	100	97
Paraquat+Cyanazine	0.5+0.25	-	100	99
Paraquat+Cyanazine	0.38+0.5	99	100	98
Paraquat+Cyanazine	0.38+1	99		99
Paraquat+Cyanazine	0.25+1	96	100	
Para&Flfp&Diur+Cyanazine	0.38&0.06&0.02+0.5	99	100	100
Para&Flfp&Diur+Cyanazine	0.38&0.06&0.02+1	99	100	100
Para&F1fp&Diur+Cyanazine	0.25&0.04&0.013+1	98	100	99
Para&Flfp&Diur+2,4-D	0.38&0.06&0.02+.25	100	100	100
Para&Flfp&Diur+2,4-D	0.5&0.08&0.027+.25	100	100	100
Para&Flfp&Diur+Clsu	0.38&0.06&0.02+.0125	100	100	100
Glyphosate&2,4-D	0.28&0.25	99	99	100
Glyphosateaz, 4-D	0.28&0.25+2.5	100	100	100
Glyphosate&2,4-D+AS	0.28&0.125	99	100	100
Glyphosate&Dicamba	0	0	0	0
Control				
0.14.04		1	1	2
C.V. %		2	1	2
LSD 5%	ing paraguat were mixe	d with	0.5%	X-77

All treatments containing paraquat were mixed with 0.5% X-77 surfactant; Paraquat = Cyclone; Para&Flfp&Diur = Cyclone TEC; dry flowable formulation of cyanazine was used; butoxyethyl ester formulation of 2,4-D was used; Glyphosate&2,4-D = Landmaster II; AS = ammonium sulfate; Glyphosate&Dicamba = Fallowmaster.

Summary. All treatments provided nearly complete control of kochia, Russian thistle, and common lambsquarters. Very hot weather in the days following spraying may have contributed to the high percentage control, particularly with treatments involving paraquat. Green foxtail was present in this experiment but could not be evaluated because this species had almost completely succumbed to the hot, dry weather.

Postemergence treatments in fallow, Carrington, 1988. Treatments were applied May 5 in barley stubble using a bicycle wheel sprayer delivering 8.5 gal/A with 8001 nozzles and 40 psi. Spraying conditions included relative humidity of 55%, air temperature of 85 F, sunny skies, and a low to moderate level of moisture stress apparent. Kochia was 1 to 2-inches tall, foxtail (mostly yellow with about 30% green and 0.5 to 2-inches tall, common lambsquarters was 1 to 2-inches tall, and horseweed was 2 to 5-inches tall. Spraying was done between 9 and 11 a.m. Visual estimates of percent control were taken on June 14. Plot size was 10 by 25 ft and the experiment was a randomized complete block design with four replications.

<u>Treatment</u> <sup>a</sup>	Rate	KOCZ	Wee									
	(1b/A)	NUCZ.	Fxtl	Wibw	Colq	How						
Glyphosate&2,4-D Glyphosate&2,4-D+AS Glyphosate&2,4-D Glyphosate&2,4-D Glyphosate&2,4-D Glyphosate Glyphosate Glyphosate Paraquat+Diquat Paraquat+Diquat Paraquat+Diuron+Diquat Paraquat+Diuron+Diquat Paraquat	0.188&0.17 0.188&0.17+2.5 0.25&0.22 0.25&0.22+2.5 0.375&0.33 0.25 0.375 0.188+0.188 0.25+0.25 0.188+0.1+0.188 0.25+0.1+0.25 0.375	90 86 99 95 100 72 95 71 90 91 92 90	72 80 80 82 77 82 76 61 73 68 74	70 77 70 81 90 62 80 55 73 67 84	98 100 100 100 83 97 57 94 77 90 86	85 86 87 94 94 67 75 82 94 77						
Paraquat+Diuron Paraquat+Diuron Paraquat+Cyanazine Paraquat+Cyanazine Paraquat&Fluazifop&Diuron Paraquat&Fluazifop&Diuron	0.5 0.375+0.1 0.5+0.1 0.375+0.25 0.5+0.25 0.375&0.06&0.02	98 93 93 91 88 86	69 65 68 74 74 81	64 65 59 61 73	92 95 100 98 98	97 100 97 98 97 100						
ara&Flfp&Diur+2,4-D ara&Flfp&Diur+2,4-D ontrol	0.5&0.08&0.027 0.375&0.06&0.02+0.25 0.5&0.08&0.027+0.25 0	98 100 99 0	87 73 82 0	64 57 74 0	94 - 99 0	81 99 96 96 0						
SD 5% All treatments containing and the two treatments wi	- PAR WAR CONTROL	10 13	13 13	25 23	9	7 9						

All treatments containing paraquat were applied with X-77 surfactant at 0.5% and the two treatments with glyphosate alone were applied with X-77 at 0.25%; Glyphosate&2,4-D = Landmaster II; Para&Flfp&Diur = Cyclone TEC; butoxyethylester formulation of 2,4-D was used; dry flowable formulation of cyanazine used; Paraquat = Cyclone; AS = ammonium sulfate.

<u>Comments</u>. None of the treatments gave acceptable burndown of foxtail and wild buckwheat. Foxtail control was probably limited by swaths of heavy chaff (deposited by the combine in 1987) that apparently protected smaller foxtail seedlings from spray. Diuron and ammonium sulfate did not consistently enhance the activity of paraquat and glyphosate, respectively.

Burndown of four-leaf oats, wheat, and foxtail millet, Fargo, 1988. Treatments were applied June 21 when oats (exper. variety 810814), wheat (Wheaton), and Siberian foxtail millet was 4-leaf, and redroot pigweed was 0.5 to 2-inches tall. Herbicides were applied with a bicycle wheel sprayer delivering 8.5 gal/A with 8001 nozzles and 40 psi. At time of spraying, air temperature was 94 F and relative humidity was 65%. Plants appeared to be under some degree of drought stress at time of spraying. Visual estimates of percent control were taken on July 5. Plot size was 10 by 18 ft and the experiment was a randomized complete block design with four replications.

		P	ercent c	<u>ontro</u>	<u> </u>		
b door replications	Rate	Oats	Wheat	Ftmi	Rrpw		
Treatment <sup>a</sup>	(1b/A)		(%	)			
HOE-00661+X77 HOE-00661+X77+AS HOE-00661+X77 HOE-00661+X77+AS Glyphosate&2,4-D+AS Glyphosate&2,4-D+AS Glyphosate+X77+AS Glyphosate+X77+AS Glyphosate+X77+AS Glyphosate+X77+AS Glyphosate+X77+AS Glyphosate+X77+AS Paraquat+X77 Paraquat+X77 Paraquat&Diuron+X77 Paraquat&Diuron+X77 Paraquat&Fluazifop&Diuron+X77 Paraquat&Fluazifop&Diuron+X77 Paraquat&Fluazifop&Diuron+X77	1+0.5% 1+0.5%+2.5 0.75+0.5% 0.75+0.5%+2.5 0.188&0.17+2.5 0.25&0.22+2.5 0.375&0.33+2.5 0.188+0.25%+2.5 0.25+0.25%+2.5 0.375+0.25%+2.5 0.375+0.25%+2.5 0.375+0.5% 0.375&0.015+0.5% 0.375&0.015+0.5% 0.5&0.02+0.5% 0.5&0.02+0.5% 0.5&0.083&0.027+0.5% 0.25&0.01+0.25+0.5%	100 100 98 97 98 99 100 100 100 100 87 91 95 98 97 96		100 100 99 100 97 99 100 100 99 99 100 69 74 80 91 93 88 92 86 0	100 100 98 99 100 99 100 99 100 100 100 100 100 10		
Control		2	2	3	J		
C.V. %		3	3	4			

Glyphosate&2,4-D = Landmaster II; AS = ammonium sulfate; Paraquat = Cyclone; Paraquat&Diuron = Surefire TEC; Paraquat&Fluazifop&Diuron = Cyclone TEC; dry flowable formulation of cyanazine was used.

Summary. HOE-00661 at 1 lb/A provided complete control of all species, though the 0.75 lb/A rate was not as effective on wheat. All rates of glyphosate and Landmaster II gave essentially complete control of all species. Complete burndown with paraquat was achieved only with redroot pigweed and with the 0.5 lb/A rate on wheat. The inclusion of diuron (Surefire TEC) or fluazifop plus diuron (Cyclone TEC) tended to improve control over paraquat alone, particularly with foxtail millet.

Burndown of boot-stage oats, wheat, and foxtail millet, Fargo, 1988. Treatments were applied July 11 when oats (exper. variety 810814) were 14 inches tall, late boot to early heading, wheat (Wheaton) was 12 inches tall, late boot to early heading, Siberian foxtail millet was 16 inches tall, early boot, and redroot pigweed was 6 to 12 inches tall. Herbicides were applied with a bicycle wheel sprayer delivering 8.5 gal/A with 8001 nozzles and 40 psi. At time of spraying, air temperature was 71 F and relative humidity was about 50%. Plants had grown up under significant drought stress and were under stress at time of spraying. Visual estimates of percent control were taken on July 21. Plot size was 10 by 18 ft and the experiment was a randomized complete block design with four replications.

<u>Treatment</u>	D-+	P	contro	0]	
Ga omen o	Rate	Oats	Wheat	Ftmi	Rrpw
	(1b/A)			(%)	
10E-00661+X77	1+0.5%	87	07	00	
10E-00661+X77+AS	1+0.5%+2.5		97	98	99
lyphosate&2,4-D+AS	0.188&0.17+2.5	94	97	99	99
lyphosate&2,4-D+AS	0.10000.17+2.5	42	61	75	70
lyphosate&2,4-D+AS	0.25&0.22+2.5	45	62	87	70
1 vphocato V77 AC	0.375&0.33+2.5	54	72	93	79
lyphosate+X77+AS	0.188+0.25%+2.5	35	63	78	39
lyphosate+X77+AS	0.25+0.25%+2.5	39	70	88	
lyphosate+X77+AS	0.375+0.25%+2.5	52			50
lyphosate+X77+AS	0.5+0.25%+2.5		84	95	71
araquat+X77	0.25+0.5%	71	92	95	83
araquat+X77	0.375+0.5%	89	90	87	84
raquat+X77		93	94	90	91
raquat&Diuron+X77	0.5+0.5%	98	95	93	91
araquat&Diuron+X77	0.375&0.015+0.5%	96	95	86	90
0 [ ] ( 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0.5&0.02+0.5%	97	95	88	89
ara&Flfp&Diur+X77	0.375&0.06&0.02+0.5%	98	96	89	
araaritpaujur+X//	0.5&0.083&0.027+0.5%	99	97		87
araduur+blquat+X//	0.25&0.01+0.25+0.5%	98		90	92
araquat+Cyanazine+X77	0.375+0.25+0.5%		95	89	89
ontrol		97	95	90	85
	0	0	0	0	0
.V. %		9	7		•
SD 5%		9	7	4	8
Glyphosate&2,4-D = Land	dmaster II; AS = ammon	9	8	5	9

'Glyphosate&2,4-D = Landmaster II; AS = ammonium sulfate; Paraquat = Cyclone; Paraquat&Diuron = Surefire TEC; Para&Flfp&Diur = Cyclone TEC; dry flowable formulation of cyanazine was used.

Summary. Treatments involving glyphosate did not provide adequate burndown of any species with the marginal exception of the 0.375 and 0.5 glyphosate rates on foxtail millet. HOE-00661 gave nearly complete control of wheat, foxtail millet, and redroot pigweed, and 94% control of oats when mixed with ammonium sulfate. The best paraquat treatments gave nearly complete control of oats, about 95% control of wheat, and about 90% control of both foxtail millet and redroot pigweed. Diuron did not appear to increase the activity of paraquat, though the addition of either fluazifop or diquat did improve control in some cases.

Carryover injury to wheat by clomazone applied in fallow, Minot, 1988. spring treatments were applied October 9, 1986 and May 22, respectively, in untilled durum stubble using a bicycle wheel sprayer delivering 17 gal/A with 8002 nozzles and 40 psi. Chlorsulfuron (0.33 oz/A) was applied over the entire experimental area in late fall of 1986 for weed control in 1987. 'Seward' winter wheat was seeded 1.5 inches deep (without tillage) on September 11, 1987. Fifty pounds of 11-52-0 was applied in the seed row at planting. Visual estimates of percent wheat injury (yellowing and bleaching of foliage) were taken on October 15, 1987 when winter wheat was in the 5-leaf stage. Tilled blocks were disked once (4-inches deep) 20, followed on the same day by planting of 'Stoa' spring wheat (1.5 inches deep) in both tilled and no-till plots, fertilized as above. Visual estimates of percent wheat injury were taken on May 18 when the spring wheat was 2.5 to 3-leaf and winter wheat was tillering and 5 to 8-inches tall. Winter wheat plots were combine harvested on July 20 and spring wheat was harvested July 28, 1988. Plot size was 16.5 by 25 feet and the experiment was a randomized complete block design with a splitplot arrangement of treatments. Soil type was a loam with pH 6.8 and 2.8% organic matter.

tter.								
(11)	Clemazono		Spring	wheat		Winte	r wheat	
Wheat	Clomazone applica-	Clomazone	<u>opr mg</u>	Grain	In,	jury	Stand	Grain
tillage	tion timing		Injury	yield	Fall	Spring	reduction	yield
system	tion timing	(1b/A)	(%)	(bu/A)		(%)-		(bu/A)
				27	0	0	0	18
No-till		0	0	<b>2</b> 1	Ö	i	25	21
	Fall	0.5	-	26	Ö	2	4	24
		0.75	2	24	Ö	2 5	6	21
		1.0	3	27	0	12	23	19
		1.25	4 5		0	14	17	22
		1.5	5	26	U	17	<b>.</b>	
		2.0	7	25				
	Spring	0.5	2	23			-	
		0.75	4	25				
		1.0	6	26		•		
		1.25	11	26	•	•	The state of the s	
		1.5	11	25	•			
		2.0	18	24	•	•		
			LSD (0.	05) Rate	NS	8	NS	NS
Tilled		0	0	21	-	When -	Teno-Ovu	•
IIIIeu	Fall	0.75	4	22	of many e	state - ST	- 37	
	1011	1.0	9	22	-	-	-	•
		1.25	9	24	ant-P		RT 653 T	
		1.5	17	21	•		m edf-Hilly	- 10 - 18
		2.0	32	23			10300-038	- 1
	Spring	0.5	5	20			bits 1 -basy	•
	Spr 1119	0.75	12	20				
		1.0	23	21			•	
		1.25	33	22		Inch -sh		-01
		1.5	44	25		Miles a	a lought a	0.000
		2.0	43	22		-		
100 (0	.05) Rate x		8	NS				
F2D (0	.US) Kale X	TITIAGE			<u>C</u>	ing ubas	t amain vic	Id Inn.

LSD (0.05) Rate x Tillage 8 NS

There was a significant effect of tillage for spring wheat grain yield (notill outyielded tilled plots) and a significant tillage and clomazone rate effect for spring wheat injury.

See next page for summary

## See previous page for data

Summary. Injury to spring wheat planted in 1988 was greater when clomazone was applied in the spring of 1987 than when applied in the fall of 1986. Injury by clomazone residues in the soil was also greater when spring wheat was planted under seedlings to more rapidly grow beneath the shallow zone of clomazone residues in noroots thus were exposed over a longer period. Evidence supporting this interpretation was the observation of much higher injury in one drill row in no-till plots. at a depth of about 0.5 inches instead of 1.5 inches for the other rows. Injury to not significantly affected. Low yields due to drought may have masked yield reductions attributable to clomazone residues. Winter wheat sustained low levels of viaffected.

Carryover injury to wheat by clomazone applied in fallow, Williston, 1988. Fall and spring treatments were applied October 8, 1986 and May 18, respectively, in untilled wheat stubble using a bicycle wheel sprayer delivering 17 gal/A with 8002 nozzles and 40 psi. Chlorsulfuron (0.25 oz/A) was applied over the entire experimental area area fall of 1986 for weed control in 1987. 'Seward' winter wheat was seeded 1.5 in late fall of 1986 for weed control in 1987. Fifty pounds of 18-46-0 was applied in the seed row at planting. Visual estimates of percent wheat injury (yellowing and bleaching of foliage) were taken on October 14, 1987 when winter wheat lowing and bleaching of foliage) were taken on October 14, 1987 when winter wheat was in the 5-leaf stage. Tilled blocks were tilled once with a field cultivator (3-inches deep) on April 26, followed on the same day by planting of 'Stoa' spring wheat (1.5 inches deep) in both tilled and no-till plots. The entire experimental area received diclofop (1 lb/A) for control of foxtail. Visual estimates of percent wheat injury were taken on May 17 when the spring wheat was 2.5 to 3-leaf and winter wheat injury were taken on May 17 when the spring wheat was 2.5 to 3-leaf and winter wheat was tillering and 5 to 8-inches tall. Plots were combine harvested on July 20. Plot size was 16.5 by 25 feet and the experiment was a randomized complete block design with a split-plot arrangement of treatments. Soil type was a fine sandy loam with pH 6.5 and 2% organic matter.

loam with ph	0.5 and Lie	or gains					
		6 :	· · hoota		Winte	r wheat	
Clomazone	-	Spring	wheat"	Inj		Stand	Grain
application	Clomazone		Grain		Spring	reduction	yield
timing	rate	Injury	yield	Fall	(%) <i>-</i>		(bu/A)
O TIME TO SERVICE OF THE PROPERTY OF THE PROPE	(1b/A)	(%)	(bu/A)		(10)		(
					0	0	3.1
	0	0	4.5	0	0		4.5
r-11	0.5	-	•	1	0	2	
Fall	0.75	1	6.5	2	0	4	4.0
		2	5.4	1	0	1	4.4
	1.0	4	6.0	3	1	1	3.7
	1.25	3	6.2	5	2	5	3.9
	1.5		4.2		-	•	
	2.0	11	5.8	•		•	•
Spring	0.5	2				œ	-
	0.75	4	5.5				
	1.0	11	3.7				
	1.25	17	4.4		•		
	1.5	23	4.9	•	-		
	2.0	36	4.7	-	•		
						NC	NS
1 CD (0 OE) [	Onto	10	NS	NS	NS	NS	113
LSD (0.05) F	Rate Fact was not	signifi	cant for	both in	jury and	grain y	iela; data

aTillage effect was not significant for both injury and grain yield; data are combined across tillage.

Summary. Injury to spring wheat planted in 1988 following fall 1986 applications of clomazone was low and ranged to 11% for the 2 lb/A rate of clomazone. Clomazone applied in the spring of 1987, however, caused significantly greater injury to spring wheat planted one year later. This injury, which ranged up to 36% for the 2 lb/A wheat planted one year later. This injury, which ranged up to 36% for the 2 lb/A rate of clomazone, was not reflected in reduced yield. Grain yields in 1988 were severely limited by drought, a situation that probably precluded the expression of grain yield reductions by carryover residues of clomazone. Injury and stand reductions in winter wheat were only mildly affected by clomazone residues. As with spring wheat, grain yield of winter wheat was low due to drought and was not affected by clomazone residues in the soil.

False chamomile control on fallow and noncrop land, Mohall, 1988. Postemergence herbicides were screened for winter annual false chamomile control on fallow or wetland areas near Mohall. Treatments were applied with a bicycle wheel type plot sprayer delivering 8.5 gpa at 35 psi to 8 to 14 inch and budding false chamomile on June 2 with sunny skies and 85 F. Visual estimates of percent control were taken on July 26.

<u>Treatment</u> <sup>a</sup>	Rate	False chamomile control
	(oz/A)	(%)
Chlorsulfuron+X-77 Metsulfuron+X-77 Metsulfuron+2,4-D-bee+X-77 Sulfometuron+2,4-D-bee+X-77 DPX-M6316+X-77 DPX-M6316+X-77 DPX-L5300+X-77 DPX-L5300+X-77 DPX-M6316&DPX-L5300+X-77 DPX-M6316&DPX-L5300+X-77 CGA-131036+X-77 Clopyralid+X-77 Imazethapyr+X-77	0.25+0.25% 0.06+0.25% 0.06+2+0.25% 1.5+0.25% 1.5+16+0.25% 0.25+0.25% 0.125+0.25% 0.125+0.25% 0.125+0.25% 0.25+0.25% 0.33&0.17+0.25% 0.25+0.25% 1+0.25%	91 96 91 29 51 21 28 58 75 50 71 76 28 28
S.V. %		
SD 5% bee = butoxyethanol ester: [	DPX-M6316&DPX-L5300 = DPX-R96	13 11

Summary. Chlorsulfuron, metsulfuron, and metsulfuron plus 2,4-D gave 91% or higher control of winter annual false chamomile. CGA-131036 gave 15% lower control than metsulfuron or chlorsulfuron. DPX-M6316 and sulfometuron gave less chamomile control than all other sulfonylurea herbicides. Clopyralid and imazethapyr did not adequately control false chamomile.

Paraquat and sethoxydim plus 2,4-D for no-till burndown, Fargo, 1988. Treatments were applied May 10 (early), May 26 (mid), and June 2 (late) when kochia was 0.5 to 1.5, 3 to 6, and 8 to 14 inches tall, respectively. A second flush of 0.5 to 3 inch tall kochia was also present on June 2. Wild oats were 3 to 6-leaf and tillering (4 to 10 inches tall) on May 26 and green foxtail was 2 to 3-leaf on June 2. Herbicides were applied with a bicycle wheel spayer delivering 17 gal/A with 8002 nozzles and 35 psi for treatments involving paraquat and diquat, and 8.5 gal/A with 8001 nozzles and 35 psi for all other treatments. Environmental conditions were as follows: on May 10, relative humidity (RH) was 17%, air temperature was 69 F, and wind was 10 to 12 mph; on May 26, RH was 57%, air temp was 71 F, and wind was 4 to 8 mph; on June 2, RH was 59%, air temp was 81 F, and wind was 0 to 2 mph. Visual estimates of percent control were taken on June 3 (24 days after treatment) for the early application, June 20 (25 DAT) for the mid application, and June 24 (22 DAT) for the late application. Plot size was 10 by 25 ft and the experiment was a randomized complete block design having four replications.

		Early	М	id		te
Constment a	Ratea	KOCZ	KOCZ	Wioa	KOCZ	Grft
<u> Freatment a</u>	(1b/A)		(%	contr	01)	
	0.5+0.5%	87	94	85	80	71
Paraquat+X77	0.375+0.5%	83	89	58	76	57
Paraquat+X77	0.25+0.5%	85	86	48	45	44
Paraquat+X77	0.5+0.1+0.5%	91	98	84	81	83
Paraquat+Diuron+X77	0.375+0.1+0.5%	85	88	45	76	71
Paraquat+Diuron+X77		83	90	41	59	67
Paraquat+Diuron+X77	0.25+0.1+0.5%	83	86	23	52	56
Paraquat+Diuron+X77	0.188+0.1+0.5%	94	90	38	88	71
Paraquat+2,4-D-bee+X77	0.25+0.25+0.5%	95	95	36	84	77
Para+2,4-D-bee+Diuron+X77	0.25+0.25+0.1+0.5%	25	86	13	45	46
Diquat+X77	0.5+0.5%	92	87	18	69	61
Diquat+2,4-D-bee+X77	0.25+0.25+0.5%	82	90	48	68	70
Paraguat+Diquat+X77	0.25+0.25+0.5%		86	30	63	57
Paraguat+Diguat+X77	0.188+0.188+0.5%	82	83	75	75	79
Sethoxydim+2,4-D-bee+OC	0.1+0.5+0.25G	94		62	80	81
Sethoxydim+2,4-D-bee+0C	0.05+0.5+0.25G	95	77	30	79	45
Sethoxydim+2,4-D-bee+OC	0.025+0.5+0.25G	94	89	82	77	84
Seth+2,4-D-bee+28%UAN+OC	0.1+0.5+1G+0.25G	92	80		74	76
Seth+2,4-D-bee+28%UAN+OC	0.075+0.5+1G+0.25G	96	86	75		78
Seth+2,4-D-bee+28%UAN+OC	0.05+0.5+1G+0.25G	94	84	58	78	79
Seth+2,4-D-bee+Dash	0.1+0.5+0.25G	94	77	85	75	82
Seth+2,4-D-bee+Dash	0.075+0.5+0.25G	93	75	81	78	
Seth+2,4-D-bee+Dash	0.05+0.5+0.25G	88	85	84	77	80
Control	0	0	0	0	0	0
LSD 5%		7	10	17	11	18

dX77 = X-77 surfactant; 2,4-D-bee = butoxyethyl ester; OC = petroleum oil with 17% emulsifier; 28%UAN = 28% urea ammonium nitrate solution; Dash = trade name of BASF surfactant of proprietary composition; G = gallons per acre (i.e. 0.25G = 1 quart per acre).

Summary. Paraquat and diquat were most efficacious on kochia when applied to 3 to 6-inch plants; 8 to 14-inch kochia were least controlled by these herbicides. Sethoxydim plus 2,4-D mixtures, however, gave highest kochia control when applied to the smallest plants. Control of wild oats or foxtail was inadequate on all treattreatments. Diuron tended to increase control by paraquat but enhancement was neither dramatic nor consistent. 2,4-D provided significant enhancement of kochia control by paraquat when compared to paraquat alone.

Paraquat and sethoxydim plus 2,4-D for no-till burndown, Pillsbury, 1988. ments were applied June 1 (early), June 10 (mid), and June 16 (late) when green foxtail was in the 1 to 4-leaf, 4 to 5-leaf (tillering, 6 to 9 inches tall), and 5-leaf (7 to 12 inches tall) stage, respectively. Redroot pigweed was in the 6 to 10-leaf stage on June 16. Herbicides were applied with a bicycle wheel spayer delivering 17 gal/A with 8002 nozzles and 35 psi for treatments involving paraquat and diquat, and 8.5 gal/A with 8001 nozzles and 35 psi for treatments involving paraquat and diquat, and ditions for these dates were as follows: on June 1, relative humidity (RH) was 48%, air temperature was 87 F, and wind was 6 to 10 mph; on June 10, RH was 34%, air temp was 76 F, and wind was 15 mph, gusting to 25 mph; on June 16, RH was 44%, air temp was 76 F, and wind was 10 to 12 mph. Visual estimates of percent control were taken on June 17 (16 days after treatment) for the early application, June 24 (14 DAT) for the mid application, and June 29 (13 DAT) for the late application. 10 by 25 ft and the experiment was a randomized complete block design having four Plot size was

			DIOCK	uesign	naving
<u>Treatment</u> <sup>a</sup>	Rate	Earl			ite
	(1b/A)	Grft		Grft	Rrpw
Paraquat+X77	(, ,,,		(% co	ntrol)-	
Paraquat+X77	0.5+0.5	85	71		
Paraquat+X77	0.375+0.	5% 86	71	73	-
Paraquat Dimen van	0.25+0.	5% 92	70	65	•
Paraquat+Diuron+X77	0.5+0.1+0	.5% 96	43	54	37*
Paraquat+Diuron+X77	0.375+0.1+	A Mai	76	80	-
Paraquat+Diuron+X77	0.25+0.1+	A =	73	79	56*
Paraquat+Diuron+X77	0.1875+0.1	. A	55	80	45
Paraquat+2,4-D-bee+X77	0.25+0.25+0	A Par	34	73	53
rara+2,4-D-bee+Diuron+Y77	0.25+0.25+0.		48	60	68*
Securtz, 4-D-bee+OC	0.1+0.5+0	APA	53	69	80*
Seth+2,4-D-bee+OC	0.05+0.5+0	000	63	58	65
Seth+2,4-D-bee+OC	0.03+0.5+0.	050	40	41	57*
Seth+2,4-D-bee+28%11AN+0C	0.025+0.5+0		25	39	59
3etn+2,4-U-bee+28%11AN+0C	0.1+0.5+1G+	0.25G 95	69	57	68*
Seth+2,4-D-bee+28%UAN+0C	0.075+0.5+1G	+0.25G 87	58	55	
Secritz, 4-U-bee+Dach	0.05+0.5+1G+	0.25G 81	48	53	68*
Seth+2,4-D-bee+Dash	0.1+0.5+0.	25G 85	63	64	71*
Seth+2,4-D-bee+Dash	0.075+0.5+0	.25G QA	70		55
Diquat+X77	0.05+0.5+0	.25G 82	54	58	
Diquat+2,4-D-bee+X77	0.5+0.59	% 79	31	49	54
Paraquat+Diquat+X77	0.25+0.25+0	0.5% 56	26	48	
Paraguat Diguat X77	0.25+0.25+0	0.5% 91		51	61*
Paraquat+Diquat+X77 Control	0.188+0.188+	0.5% 83	58	66	-
60111101	0	0	45	53	36
LSD 5% for companies.		U	0	0	0
LSD 5% for comparisons inve	Olving no *	12	10	10	
		12	18	12	20
					21
aX77 = X-77 surfactant; 2,4 with 17% emulsifier; 28%UA	-D-bee = butox	vethyl estare	00		23
with 17% emulsifier; 28%UA trade name of BASF surfact	N = 28% urea a	mmonium situat	OC = pe	troleum	oil
trade name of BASF surfact acre (i.e. 0.25G = 1 quart	ant of proprie	tary composition	soluti	on; Da	sh =
acre (i.e. 0.25G = 1 quart	per acre)	- TILSOUIDOS 1 £10	n; G = 0	gallons	per

acre (i.e. 0.25G = 1 quart per acre).

Summary. Treatments applied on 5-leaf and larger green foxtail (mid and late application timings) provided only fair to poor control. The early application generally gave better control although few treatments showed control percentages of mid-90s or Diuron appeared to enhance control by paraquat. Dash and 28% urea ammonium nitrate did not appear to enhance foxtail control by sethoxydim plus 2,4-D combinations. Foxtail control by diquat or diquat plus paraquat combinations was comparable to control provided by paraquat alone at equivalent rates of herbicide.

Glyphosate and HOE-00661 for no-till burndown, Fargo, 1988. Treatments were applied May 10 (early), May 26 (mid), and June 2 (late) when kochia was 0.5 to 1.5, 1.5 to 5, and 8 to 14 inches tall, respectively. A second flush of kochia was present on 5, and 8 to 14 inches tall, respectively. A second flush of kochia was present on 5 and was 0.5 to 3 inches tall. Wild oats were 3 to 4-leaf (5 to 8 inches 5) June 2 and was 0.5 to 3 inches tall. Wild oats were 3 to 4-leaf (5 to 8 inches 5) June 2 and wild green foxtail was 2 to 3-leaf on June 2. Herbicides were applied with a bicycle wheel spayer delivering 8.5 gal/A with 8001 nozzles and 35 psi for all treatments involving glyphosate, and 17 gal/A with 8002 nozzles and 35 psi for all other treatments. Environmental conditions were as follows: on May 10, relative humidity (RH) was 17%, air temperature was 69 F, and wind was 10 to 12 mph; on ative humidity (RH) was 17%, air temperature was 69 F, and wind was 10 to 12 mph; on May 26, RH was 57%, air temp was 71 F, and wind was 4 to 8 mph; on June 2, RH was 59%, air temp was 81 F, and wind was 0 to 2 mph. Visual estimates of percent control were taken on June 3 (24 days after treatment) for the early application, June 20 (25 DAT) for the mid application, and June 24 (22 DAT) for the late application. Plot size was 10 by 25 ft and the experiment was a randomized complete block design having four replications.

four replications.		Early	Mi		La	
	Rate	KOCZ	KOCZ	Wioa	KOCZ	Grft
<u>reatment<sup>a</sup></u>	(1b/A)		(%	contro	1)	
		70	00		96	84
Typhosate+X77	0.5+0.5%	72	98	97*	92	78
Typhosate+X77	0.375+0.5%	71	96	94	77	65
llyphosate+X77	0.25+0.5%	50	93		94	85
Slyphosate+AS+X77	0.5+2.5+0.5%	73	97	96	95	82
Slyphosate+AS+X77	0.375+2.5+0.5%	76	93	94	86	75
Slyphosate+AS+X77	0.25+2.5+0.5%	37	91	91*		81
Slyt&2,4-D+2,4-D-dma	0.375&0.33+0.17	94	99	an wife	97	76
Slyt&2,4-D+2,4-D-dma	0.25&0.22+0.28	93	98	74	93	86
Slyt&2,4-D+2,4-D-dma	0.188&0.17+0.33	93	98	-	93	75
Glyt&2,4-D+2,4-D-dma+AS	0.375&0.33+0.17+2.5	94	98	97	93	84
Glyt&2,4-D+2,4-D-dma+AS	0.25&0.22+0.28+2.5	95	100		96	
Glyt&2,4-D+2,4-D-dma+AS	0.188&0.17+0.33+2.5	85	77	-	96	86
61yta2,4-0+2,4-0-dilla1713	1+0.5%	51	91	48*	85	70
HOE-00661+X77	0.75+0.5%	36	85	49*	81	73 60
HOE-00661+X77	0.5+0.5%	45	64	-	67	
HOE-00661+X77	0.75+0.25+0.5%	72	86	53*		58
HOE-00661+Glyphosate+X77	0.75+0.188+0.5%	58	86	66*		6
HOE-00661+Glyphosate+X77	0.5+0.25+0.5%	34	77	46	76.	60
HOE-00661+Glyphosate+X77	0.5+0.188+0.5%	46	80	43*		6
HOE-00661+Glyphosate+X77	0.375+0.25+0.5%	19	73	34*		7
HOE-00661+Glyphosate+X77		6 34	99	95	76	6
HOE-00661+Glyphosate+AS+	0.375+0.188+0.5%	35	69	43*		6
HOE-00661+Glyphosate+X77	0	0	0	0	0	
Control				10	9	1
LSD 5% for comparisons i	nvolving no *	22	6	19	9	1
ICD 5% for comparisons 1	nyolying one "			21		2
LSD 5% for comparisons i	nvolving two *			22		

 $<sup>^{</sup>a}$ AS = ammonium sulfate; Glyt&2,4-D = Landmaster II herbicide containing 0.9 lb ae/gal glyphosate and 0.8 lb ae/gal 2,4-D; X77 = X-77 surfactant; 2,4-D-dma = dimethylamine salt of 2,4-D.

Summary. The only treatments providing greater than 90% control of 0.5 to 1.5 inch kochia were those involving glyphosate plus 2,4-D. Later kochia stages were more easily controlled and greater than 90% control was achieved with glyphosate or glyphosate plus 2,4-D. Wild oats were controlled most effectively by glyphosate. excellent foxtail burndown was not provided by any treatments. Control was often enhanced by addition of ammonium sulfate. Glyphosate applied with HOE-00661 generally gave less control than equal amounts of glyphosate applied alone.

Glyphosate and HOE-00661 for no-till burndown, Pillsbury, 1988. Treatments were applied June 1 (early) and June 9 (late) when green foxtail was 1 to 4-leaf and 4 to 5-leaf and tillering (6 to 9 inches tall), respectively. Redroot pigweed was in the delivering 8.5 gal/A with 8001 nozzles and 35 psi for all treatments involving glyronmental conditions were as follows: on June 1, relative humidity (RH) was 48%, air temperature was 87 F, and wind was 3 to 6 mph; on June 9, RH was 39%, air temp was control were taken on June 17 (16 days after treatment) for the early application experiment was a randomized complete block design having four replications.

		Wee		rol
<u>Treatment<sup>a</sup></u>	Rate	Early		ate
	(1b/A)	Grft	Grft	Rrpi
07	(10/A)		(%)	
Glyphosate+X77	0.5+0.5%	7.0		
Glyphosate+X77	0.375+0.5%	73	92	907
Glyphosate+X77	0.25+0.5%	78	87	93
Glyphosate+AS+X77	0.5+2.5+0.5%	70	73	66
Glyphosate+AS+X77	0.375+2.5+0.5%	77	96	95
Glyphosate+AS+X77	0.373+2.5+0.5%	77	94	91
Glyt&2,4-D+2,4-D-dma	0.25+2.5+0.5%	74	83	_
61yt&2,4-D+2,4-D-dma	0.375&0.33+0.17	75	83	88
61yt&2,4-D+2,4-D-dma	0.25&0.22+0.28	72	70	66*
61yt&2.4-D+2.4-D-dma+AC	0.188&0.17+0.33	71	73	60
41 y L&2, 4-D+2, 4-D-dm2+AC	0.375&0.33+0.17+2.5	79	83	90
41y Laz, 4-D+2.4-D-dma+49	0.25&0.22+0.28+2.5	72	85	83
UOE-00001+X//	0.188&0.17+0.33+2.5	73	78	75*
HOE-00661+X77	1+0.5%	71	77	69*
HOE-00661+X77	0.75+0.5%	70	65	53
10E-00661+G1 vnhosato+4777	0.5+0.5%	63	46	39
10L~00001+121Vnh0c3+01V77	0.75+0.25+0.5%	75	60	58
10[-0000]+[-1000nnca+a-077	0.75+0.188+0.5%	75	57	43
106-00001+131Vnh0ca+01V77	0.5+0.25+0.5%	72	54	45
10E-00661+Glyphosate+X77	0.5+0.188+0.5%	67	46	53
OF _00001+121ADV0053+01VC.A.	0.375+0.25+0.5%	71	58	76
OE-00661+Glyphosate+X77		68	70	, 0
ontrol	0.375+0.188+0.5%	68	43	39*
	0	0	0	0
SD 5% for comparisons inv			2 121	U
SD 5% for comparisons inv	Olving no *	5	13	22
SD 5% for comparisons inv	olving one *	1317 4	THE RESERVE	24
AS = ammonium sulfator	Clark Color			26
AS = ammonium sulfate; taining 0.9 lb ae/gal gly 77 surfactant; 2,4-D-dma	Glyt&2,4-D = Landmaster	II herh	icide d	on-
77 sunfactor to de/yal gly	phosate and 0.8 lb ae/ga = dimethylamine salt of 2	1 2 1 0.	V77	: X-

<u>Summary</u>. All treatments provided between about 65 and 80% green foxtail control at the early application timing. Greater foxtail control was achieved at the late application, although control percentages as high as the mid-90s were obtained only with the highest rates of glyphosate plus ammonium sulfate. HOE-00661 appeared to antagonize the effect of glyphosate on green foxtail. Best control of redroot pigweed was obtained with glyphosate or glyphosate plus ammonium sulfate.

Burndown efficacy of residual herbicides, Fargo and Pillsbury, 1988. At Pillsbury, cyanazine treatments were applied on June 1 when green foxtail was 1 to 4-leaf, while imazethapyr and linuron were applied June 10 when green foxtail was 4 to 5-leaf with some tillering. Environmental conditions at Pillsbury were as follows: on June 1, relative humidity (RH) was 48%, air temperature was 87 F, and wind was 10 to 12 mph; on June 10, RH was 34%, air temp was 80 F, and wind was 15 mph with gusts to 25 mph. At Fargo, cyanazine treatments were applied June 2 when green foxtail was 2 to 4-leaf, kochia was 0.5 to 8 inches tall, and redroot pigweed was 2 to 4-leaf. Imazethapyr and linuron were applied at Fargo on June 9 when green foxtail was 4 to 6-leaf, kochia was 4 to 14 inches tall, and redroot pigweed was 4 to 6-leaf. Environmental conditions at Fargo were as follows: on June 2, RH was 40%, air temp was 90 F, and wind was 0 to 2 mph; on June 9, RH was 39%, air temp was 67 F, and wind was 8 to 12 mph. All treatments were applied with a bicycle wheel sprayer delivering 17 gal/A with 8002 nozzles and 35 psi. Visual estimates of percent weed control were taken June 29 at Pillsbury and June 30 at Fargo. Plot size was 10 by 25 ft and the experiment was a randomized complete block design having four replications.

			Weed	contro	
			Fargo		Pillsbury
a a a a a a a a a a a a a a a a a a a	Rate	Grft		Rrpw	Grft
reatment <sup>a</sup>	(1b/A)			- (%)	
1.00	2+0.25G <sup>a</sup>	59	80*	72	50
yanazine-L+OC	2+0.25G	40	59*	. 47	48
yanazine-L+Dash	2.5+0.25G	78	83*	69	39
yanazine-L+OC		69	77*	70	49
yanazine-L+Dash	2.5+0.25G	67	87	87	42
yanazine-L+OC	3+0.25G	66	88*	73	61
yanazine-L+Dash	3+0.25G	84	77	87	61
[mazethapyr+X77	0.063+0.5%	85	71*	87	65
imazethapyr+OC	0.063+0.25G	90	84	85	69
[mazethapyr+Dash	0.063+0.25G	87	73	86	70
Imazethapyr+X77	0.094+0.5%		64*	69*	26
_inuron+X77	0.75+0.5%	43	88	92	25
Linuron+OC	0.75+0.25G	55		81	31
Linuron+Dash	0.75+0.25G	39	70 83	95	26
Linuron+X77	1.25+0.5%	63		90	44
Linuron+OC	1.25+0.25G	61	83		31
Linuron+Dash	1.25+0.25G	69	87*	0	0
Control	0	0	0	U	
LSD 5% for compari	cons involving NO	* 21	12	21	21
LSD 5% for compari		e *	13	23	
	sees involving th	n *	14	24	
AV77 V 77 cust	actant; OC = petr	oleum o	oil wit	h 17%	emulsifie
a l tuada n	amo of RASE SILY	rtactani	COT	ו זקטיזק	etary co
Dash = trade n position; G = gal	// /i a 0 256 =	1 quart	t per a	cre).	

Summary. None of the treatments provided acceptable burndown of green foxtail. Kochia control was generally higher than foxtail control but did not exceed 90% for any of the treatments. Redroot pigweed control from 90 to 95% was achieved with 1.25 lb/A of linuron. No clear differences were apparent between control obtained with oil adjuvant, X-77, or Dash.

Antagonism of glyphosate burndown by residual herbicides, Fargo, 1988. Treatments were applied June 2 when wild oats were 4 to 5-leaf and tillering and foxtail (mostly green with some yellow foxtail) was 2 to 3-leaf. Herbicides were applied with a bicycle wheel sprayer delivering 8.5 gal/A with 8001 nozzles and 35 psi. At time of spraying, air temperature was 94 F, relative humidity was 32%, and wind was 0 to 4 treatment). Plot size was 10 by 25 ft and the experiment was a randomized complete block design having four replications.

<u>Treatment</u> <sup>a</sup>	Data	Weed o	contro
DY TELL PROPERTY.	Rate	Wioa	Fxt
	(1b/A)	(%	6)
Glyphosate	0 275		
Glyphosate	0.375	78	83
Metolachlor+Glyphosate	0.25	56	73
Metolachlor+Glyphosate	2.5+0.375	71	78
Alachlor+Glyphosate	2.5+0.25	58	69
Alachlor+Glyphosate	2.5+0.375	77	81
Alachlor-MT+Glyphosate	2.5+0.25	63	68
Alachlor-MT+Glyphosate	2.5+0.375	33	35
Propach on Clumb and	2.5+0.25	30	34
Propachlor+Glyphosate	4+0.375	37	49
Propachlor+Glyphosate	4+0.25	52	45
Pendimethalin+Glyphosate	1.5+0.375	71	78
Pendimethalin+Glyphosate	1.5+0.25	56	69
hloramben+Glyphosate	2.5+0.375	61	
hloramben+Glyphosate	2.5+0.25	45	74
letribuzin+Glyphosate	0.375+0.375	71	60
letribuzin+Glyphosate	0.375+0.25		77
mazethapyr+Glyphosate	0.063+0.375	55	63
mazethapyr+Glyphosate	0.063+0.25	82	89
mazethapyr+Glyphosate	0.063+0.188	63	78
mazethapyr	0.063	59	69
inmethylin+Glyphosate	1.25+0.375	35	44
inmethylin+Glyphosate	1.25+0.25	81	88
ontrol	0	71	79
	0	0	0
SD 5%			
X-77 surfactant was tank-m Alachlor-MT = microanca	dyad	12	11

Alachlor-MT = microencapsulated formulation of alachlor (Lasso MicroTech); Fxtl = predominantly green foxtail with some yellow foxtail.

<u>Summary</u>. Wild oat control by glyphosate was reduced substantially by alachlor-MT and propachlor and to a lesser extent by chloramben. Similar results were obtained with foxtail control.

Common milkweed control on fallow, Minot 1988. Treatments were applied with a bicycle wheel plot sprayer delivering 8.5 gpa at 35 psi to post flowering 20 to 30 inch common milkweed on July 20 with sunny skies and 70 F and 10 to 15 mph NW winds. Evaluations were made on August 4.

Treatment	Rate	Milkweed
reacine it	(1b/A)	(% control)
2,4-D bee Dicamba 2,4-D dma + Dicamba 2,4-Dp & 2,4-D & Dicamba 2,4-Dp Clopyralid & 2,4-D Triclopyr & 2,4-D Picloram Sulfometuron + X-77 DPX-L5300 + X-77 DPX-L5300 + X-77 Fluroxypyr Fluroxypyr Glyphosate + X-77  C.V. % LSD 5%	2 1 1+0.25 2.25 1 1.2 3 0.5 0.0625+0.25% 0.015+0.25% 0.0625+0.25% 0.125 0.25 1.5+0.25%	36 71 26 64 20 13 84 86 5 4 5 4 63 56

a bee = butoxyethanol ester; Clopyralid & 2,4-D = Curtail; Triclopyr & 2,4-D = Crossbow; Fluroxypr = Starane; X-77 = surfactant;

## Summary

Evaluations in the year of treatment indicate that picloram and triclopyr & 2,4-D cause the most initial injury to common milkweed and the sulfonyl urea herbicides the least. An evaluation during the spring of 1989 will be made of the treated areas to determine herbicide effectiveness for common milkweed control the year following treatment.

