

Effect of preplant-applied Assure II on durum (2003)

The objective of the study was to define a safe replant interval from preplant applications of Assure II prior to planting spring seeded monocot crops.

Table. Effect of preplant-applied Assure II on durum

Treatment ^a	Rate	Stand reduction			Growth reduction			Yield	Test Wt
		Jun 7	Jun 19	Jul 19	Jun 7	Jun 19	Jul 19	bu/A	lb/bu
		% —————							
20 DPP									
Assure II	5 fl oz	0	0	0	0	0	0	45	63.0
Assure II	7 fl oz	0	0	0	0	0	0	50	63.1
Assure II	14 fl oz	0	0	0	0	0	0	48	62.9
Fusilade	24 fl oz	7	8	7	6	7	1	43	62.7
Poast	32 fl oz	2	1	0	1	0	0	42	63.4
Select	16 fl oz	3	3	1	2	2	0	48	63.1
7 DPP									
Assure II	5 fl oz	1	0	0	0	0	0	46	63.7
Assure II	7 fl oz	1	0	0	0	0	0	45	63.3
Assure II	14 fl oz	1	1	0	0	0	0	48	62.7
Fusilade	24 fl oz	4	2	2	3	1	0	45	63.0
Poast	32 fl oz	0	0	0	0	0	0	45	63.2
Select	16 fl oz	4	3	1	4	3	0	49	63.1
0 DPP									
Assure II	5 fl oz	0	2	0	0	1	0	43	63.6
Assure II	7 fl oz	0	1	0	0	1	0	51	63.4
Assure II	14 fl oz	2	4	0	1	3	0	46	62.9
Fusilade	24 fl oz	11	11	7	7	9	1	47	62.6
Poast	32 fl oz	3	3	1	2	2	0	48	63.1
Select	16 fl oz	13	13	12	7	7	2	48	62.8

Untreated	0	0	0	0	0	0	52	63.0
LSD (0.05)	6	6	5	4	5	1	NS	NS
CV	129	124	184	149	160	268	10	0.7

^ADPP=days prior to planting.