

Table . Forage yield of warm-season annual forages at Fargo, ND, in 2006 to 2008.

Entry	2006			2007			2008			3-yr avg.
	Cut 1	Cut 2	Total	Cut 1	Cut 2	Total	Cut 1	Cut 2	Total	
	-----tons dry matter/acre-----									
Cerise red proso millet	1.96		1.96	2.04		2.04	3.30		3.30	2.43
Japanese millet	2.13		2.13	4.29		4.29	‡		0.00	2.14
Golden German FM [†]	3.23		3.23	4.38		4.38	3.46		3.46	3.69
Manta Siberian FM [†]	2.39	1.36	3.75	2.02		2.02	2.56		2.56	2.78
102M pearl millet	2.41		2.41	2.35		2.35	4.20		4.20	2.99
Piper sudangrass (PS)	1.88	1.82	3.70	1.94	1.34	3.28	1.86	2.20	4.06	3.68
Nutri + BMR SxS [†]	1.78	1.61	3.39	1.92	1.06	2.98	1.50	2.18	3.68	3.35
Sorghum x Sudangrass	1.95	1.58	3.53	2.48	1.14	3.62	1.95	2.29	4.24	3.80
Paul oat	1.93		1.93	2.38		2.38	2.56		2.56	2.29
Haybet barley	2.17		2.17	-		-	-		-	-
Martin red milo/soybean	-		-	-		-	5.10		5.10	-
Paul oat/PS	-		-	-		-	1.57	2.55	4.12	-
Mean	2.18		2.82	2.64		3.04	2.81		3.73	
LSD (0.05)	0.53	0.30	0.56	0.58	0.21	0.57	0.55	NS	0.71	
CV,%	16.7	11.9	13.6	12.6	8.2	10.9	11.3	21.1	11.1	

[†]FM = foxtail millet; SxS = sorghum x sudangrass

[‡]Japanese millet was seeded, emerged, but never grew above 6 inches.