

Haylage and Other Fermented Forages

J.W. Schroeder
Extension Dairy Specialist

Almost any legume, grass or pasture forage can be ensiled successfully

Cutting fresh forage at the optimal stage of maturity and feeding it directly to animals year-round would supply the highest-quality and most palatable feed possible. In addition, field and storage losses would be the least of all methods of forage utilization. However, fluctuations in seasonal growth and plant maturity make harvesting and storing forages necessary to maximize quality and productivity.

Hay is the most widely used method of storing forages, but harvesting as grass or legume silage or haylage has several advantages over storing as hay, including:

- Lower field losses when harvested as silage
- More nutrients preserved for feeding
- Lower labor costs because of more complete mechanization in harvesting and feeding
- Consistent forage quality on a daily basis when properly stored
- Greater ability to harvest the crop at ideal maturity because less rain-free weather is needed for harvesting haylage

Individual crops and mixtures used for making ensiled forage differ with climate, soil type and crop rotation in North Dakota. Almost any legume, grass or pasture forage can be ensiled successfully. Select the harvest time by the growth stage (maturity) of the predominant crop in the mixture.

Definitions

Forage silages can be separated into three groups on the basis of harvest moisture levels:

- High-moisture or **direct-cut silage** at 70-plus percent moisture
- **Wilted silage** at 60 to 70- plus percent moisture
- **Low-moisture haylage** at 40 to 60 percent moisture

Direct-cut Silage


Freshly cut forage normally will contain 75 to 85 percent water. Ensiling direct-cut forages prevents weather damage and leaf shattering, and lowers harvest costs, but seepage loss from silages at this moisture level can be high.

The high moisture level in direct-cut silages can cause an abnormal low-temperature fermentation, producing conditions favorable for undesirable clostridial organism growth. This produces silage that has an unpleasant sour, butyric-acid smell that severely reduces livestock consumption. Freezing during winter months is an additional concern because it causes excessive moisture in the silage, making unloading more difficult.

Wilted Silage

In making wilted silage, the forage is cut and left to wilt to reduce the moisture content. The number of hours of wilting required can vary depending on geographic location and weather conditions. Under ideal drying temperatures of late spring and early summer, four to six hours are usually adequate to reduce moisture levels to around 65 percent. Ideal moisture levels for storage in upright silos range from 62 to 65 percent and bunker silos from 65 to 72 percent. Seepage and runoff do not normally occur at these moisture levels and nutrient preservation is maximized.

The most common problem encountered in making wilted silage is that producers allow too much wilting to occur before ensiling. Wilted silage with drier-than-recommended moisture content will heat and have reduced digestibility as well as higher harvest and storage losses (Figure 2).



Adapted from: Hoglund, 1964

Figure 1. Estimated dry-matter loss during harvest and storage of forage crops at various moisture levels.




Figure 2. Moisture management guide for preserving forages as silage or hay.

Haylage

Making haylage out of harvested legumes and grasses is an alternate option for storing and processing forage. Haylage can be made from any crop that traditionally is stored as hay (Table 1).

Ensiling grasses and legumes as haylage requires 40 to 60 percent moisture. Reductions in moisture content necessary for production of haylage are accomplished by conditioning (mowing, windrowing and drying for four to 24 hours) and depends

Table 1. Crops commonly used for haylage in North Dakota and recommended stage of maturity for harvesting.

Crop	Stage of Maturity
Legumes	
• alfalfa	pre-bud to 1/10 bloom for first cutting 1/10 to ¼ bloom for second cutting ¼ to ½ bloom for third cutting
• sweet clover (and other clovers)	½ bloom
Perennial Grasses	
• cool-season grasses – brome grass – crested wheat	before grass heads emerge boot to early head (May 15 to June 1) boot to early head (May 15 to June 1)
• wetland vegetation ¹	before heads emerge
Annual Grasses	
• sudangrass and sorghum (includes hybrids)	boot to early head
• millets	boot to early head
• small grains (oats, wheat, barley)	boot to early dough stage

¹ Wetland vegetation makes better forage stored as haylage than as baled hay at any stage of maturity.

on forage moisture and conditions. Considerably longer wilting is necessary to reduce moisture levels to 40 to 60 percent.

Haylage is the most sensitive of the ensiled forages to moisture variation. *Just a few percentage points up or down can ruin your forage stored in this manner.*

Low-moisture haylage often becomes too dry for good harvesting and storage. Leaves, as well as some of the nutrient value, are lost in the dust created by the harvesting equipment. Whenever large clouds of dust come from the wagon being loaded, the crop is too dry for haylage and is better suited to hay.

In some cases, but less common than in years past, conventional upright silos are used for low-moisture haylage, but you must pay particular attention to air exclusion. Fine chopping, rapid filling and good sealing to keep air out of the forage are critical. Allowing air into the haylage will cause heating and the growth of undesirable yeast and molds.

In addition, haylage can heat spontaneously if oxygen becomes available. Fires have been reported when the moisture in the haylage is near 40 percent. More often, the heating will form indigestible products that lower protein and energy values. A haylage that is tobacco-brown or black and has a caramelized odor has undergone some spontaneous heating.

Length of Cut

Large forage particles create difficulties in packing the silo because too much air will be trapped, causing conditions favorable for excessive heating and mold growth. Poorly chopped material with long stems is also difficult to handle mechanically, especially with silo unloading equipment and feed conveyors.

Forages can be chopped as fine as a quarter-inch if between 25 and 50 percent of the dry matter fed in the rations is from long-baled hay. With an all-silage feeding program, the theoretical length of cut (TLC) should be ¾ inch, resulting in

15 to 20 percent of the particles being more than 1.5 inches long. Chopping forages too fine decreases rumination, cud chewing and salivation and may cause rumen acidosis and low milk levels.

Feeding improperly fermented haylage can result in very adverse effects, including a decline in milk production and an increase in dead or cull cows.

Stage of Maturity

Harvesting forages at the optimum stage of maturity is crucial for top animal performance and productivity. Wisconsin Forage Council data show that alfalfa takes only five to six days to mature from mid-bud to 1/10-bloom in the Midwestern states.

Delays in harvesting grasses after heads emerge or alfalfa after the 1/10-bloom stage decrease the quality of the forage. This is because the proportion of fibrous stems increases as the percentage of leaves decreases. Each day of delay in harvesting results in the crop dropping 0.5 percent in crude protein, while increasing 0.7 percent in acid detergent fiber (ADF) and 0.9 percent in neutral detergent fiber (NDF).

Lignin, a potentially nondegradable fiber fraction, doubles in quantity from 5 to 10 percent between the early bud and full-bloom stages. Not only is lignin not digestible, but it binds with other fiber components (cellulose and hemicellulose), as well as protein, making them nondigestible as well.

Studies have shown that for each day alfalfa matures past the late-bud stage, milk production can decrease by 0.86 pounds per

cow per day, or up to 400 pounds per cow per lactation.

Pre-bud or bud-stage alfalfa leaves are high in protein and low in fiber, so the highest quality alfalfa can be obtained by harvesting when the leaf-to-stem ration is the highest. The various stages of alfalfa maturity are outlined in Table 2.

As forage matures, yield increases, but more plant fiber results in decreased digestibility. That means more grain must be fed to maintain high production levels, but has the risk of compromising ruminal health. Cows normally eat forages at 2.5 percent of their body weight daily if fed hay harvested in the bud stage but only about 1.5 percent of their weight if fed mature, full-bloom alfalfa (Table 3).

Table 2. Stages of alfalfa maturity.

Early bud	1 to 2 nodes with buds, no flowers
Late bud	more than 3 nodes with buds, no flowers
Early flower	1 node with 1 open flower
Late flower	more than 2 nodes with open flowers

Adapted from Kalu and Fick.

Table 3. Effect of stage of maturity on digestibility of alfalfa.

Stage	Percent Digestibility
Pre-bud	66.8
Bud	65.0
Early bloom	63.1
Mid-bloom	61.3
Full bloom	59.4
Late bloom	57.5
Mature	55.8

Not harvesting plants too early also is important. Immature plants, in the early bud stage, have a high proportion of leaves, and the stems have not become highly lignified, hard and unpalatable. However, by cutting alfalfa at the immature-bud stage for more than one year, carbohydrate root reserves may become depleted and plant vigor reduced. This effect is shown in Table 4.

Grass Silage

Grass as a silage crop is especially suited to areas of high rainfall where harvesting and storing of forages is made difficult because of reduced drying weather. The production of grass silage should employ the same general management practices discussed previously.

Management attention should be focused on:

- cutting when the first stems begin to head out
- wilting to less than 72 percent moisture
- chopping at ¼ to ⅜ inch length
- packing the silage well to facilitate an anaerobic environment in the storage structure

Grass silage often is harvested in wet environments not suitable for field wilting.

Ensiling direct-cut grass at high moisture levels (75 to 85 percent) presents management challenges.

These high moisture ranges may allow an undesirable clostridial fermentation in the silage that produces a high level of sour-smelling butyric acid. Clostridia also can contribute to increased silage protein breakdown.

Another challenge with direct-cut grass silage is the additional lactic acid production required to lower the pH of extremely wet silage. This lactic acid is produced by bacteria feeding on plant sugars. Proper crop maturity at harvest and the presence of efficient fermenting bacteria will help reduce potential runoff problems.

If environmental conditions permit, the most ideal management practice is to wilt grass silage to approximately 65 to 72 percent moisture. This will provide an ensiling environment favorable for bacteria that can ensure proper fermentation.

Table 4. Effects of different alfalfa cutting frequencies on three-year yield, quality, weeds and stand life at the end of the third year.

Maturity	Harvest			% Dry Matter				
	Interval	Per Year	Yield	TDN	Protein	Leaves	Weeds	Stand
	(days)		(T/A)					(%)
Pre-bud	21	9-10	7.5	62.6	29.1	58	48	29
Mid-bud	25	8-9	8.8	60.2	25.2	56	54	38
10% bloom	29	7	9.9	58.2	21.3	53	8	45
50% bloom	33	6-7	11.4	57.8	18.0	50	0	56
100% bloom	37	5-6	11.6	55.7	16.9	47	0	50

Source: V.L. Marble, 1974. Fourth California Alfalfa Symposium.

Sorghum Silage

Sorghum is a popular silage crop in many arid areas of the U.S. Sorghum adapts well to low-rainfall areas and growing conditions not well suited for corn production. Sorghums vary widely in physical characteristics but can be categorized into three main groups: forage sorghums, sorghum-sudangrass and grain sorghums.

Forage sorghum is the most common sorghum harvested for silage. It provides excellent tonnage and generally will have 80 to 90 percent of the feeding value of corn silage. Forage sorghum should be harvested when the grain is in the medium- to hard-dough stage or when it begins to lose color. The ideal moisture content should be 60 to 72 percent, depending on the type of storage structure to be used.

Sorghum typically dries slowly. Plant an early maturing hybrid if using a sealed structure and ensile at 50 to 60 percent moisture. Avoid ensiling in bunkers if moisture levels are above 75 percent. Excessive moisture may produce run-off and encourage the production of butyric acid, which produces less palatable silage. The recommended chop length is $\frac{3}{8}$ to $\frac{1}{2}$ inch. However, if the crop becomes overly dry, chop finer and consider using a recutter screen.

Sorghum adapts well to low rainfall areas and growing conditions not well suited for corn production

Sorghum-sudangrass typically has been used as a summer pasture or hay crop. However, if ensiled properly, sorghum-sudangrass can make excellent-quality silage. Total tonnage yield will be less than from forage sorghum, but if harvested early, it will contain $1\frac{1}{2}$ to two times the amount of protein, reducing the need for supplemental protein in the ration.

Sorghum-sudangrass should be harvested at the boot stage, at about 3 to 4 feet tall, just prior to head emergence. When ensiling at the recommended moisture of 60 to 72 percent, the crop likely will require field wilting.

Sorghum-sudangrass has fast regrowth and a second or third crop generally can be harvested if adequate moisture prevails. Be sure the plant is a minimum of 18 inches high before harvest to avoid possible prussic acid poisoning.

Grain sorghum has the highest grain-to-forage ratio of all sorghums and will have the greatest energy feeding value. Yield will be less because the plants are shorter. For optimum feeding value and efficient fermentation, harvest grain sorghum when the grain is in the medium- to hard-dough stage of maturity and moisture is 60 to 68 percent. Harvesting once the grain is fully developed will result in lower feed value because much of the grain will pass through the animal undigested.

Sweetclover Silage

Sweetclover sometimes is stored as silage in North Dakota. It should be harvested for silage at the 10 to 20 percent bloom growth stage. Sweetclover harvested at this growth stage will have about the same feed value as alfalfa.

Sweetclover, like alfalfa, is low in sugar content and requires field wilting to concentrate the sugars in the cell sap. The crop, when harvested at the 10 to 20 percent bloom growth stage, contains about 80 percent moisture. The crop should be wilted to an average of about 65 percent moisture when placed in the silo as high-moisture silage. If stored as haylage in oxygen-limiting silos, the moisture content should be 55 to 60 percent.

Sweetclover contains a compound called coumarin. The coumarin is converted to a toxic substance in molded or spoiled silage and may cause "sweetclover bleeding disease." The toxic substance (dicoumarol) reduces the clotting power of animals' blood. Animals consuming too much dicoumarol through time may bleed to death from slight wounds or internal hemorrhages. If animals are to be dehorned or castrated, do not feed sweetclover forage for at least three weeks prior to working cattle and at least 30 days prior to calving.

Molded or spoiled sweetclover silage can be very dangerous when fed as the only forage if high dicoumarol levels are present. Preventing mold formation in sweetclover silage is very important if high coumarin content varieties are grown.

Sorghum adapts well to low-rainfall areas and growing conditions not well suited for corn production.

Choose a low-coumarin variety of sweetclover that can be grown to eliminate the potential danger of “sweetclover bleeding disease.” Always purchase certified seed or the benefits of growing a low-coumarin variety will be lost. This is because low-coumarin varieties may revert back to a high coumarin level if more than three generations away from breeders’ seed.

Follow good silage management practices. Fill the silo rapidly, harvest at the proper moisture content, pack to exclude air from the silage mass, distribute the chopped forage uniformly in the silo when filling and cover to exclude outside air from entering the silage. Another alternative is to place a layer of another forage type over the sweetclover silage to reduce the potential for molding of sweetclover, then cover the silo.

Small-grain Silage

Silage is an excellent way to utilize small-grain crops used in companion cropping programs or grown to supplement forage supplies. Harvesting cereals as silage also can salvage a crop that has been damaged by hail, frost or insects.

Cereal silages should be harvested in the milk to soft-dough stage to maximize the yield of energy per acre. As cereal grains mature from the boot to the dough stage, the protein level drops while the energy value increases. Dairy producers can maximize protein

content by harvesting cereals in the boot stage. Little wilting is necessary when harvesting at the early dough stage.

Approximate feeding values for commonly grown cereal crops for North Dakota are found in Table 5.

Moisture levels in the range of 60 to 70 percent are best for ensiling cereal-grain silage, with 65 percent being the optimum. Small-grain silages with less than 60 percent moisture are difficult to pack. These crops dry quickly once cut and care should be taken to chop only at the recommended moisture content.

Baleage

Baleage, also known as round-bale silage, is a somewhat newer method of preserving forage. Baleage is simply forage that is baled at a higher moisture content than dry hay and then stored in sealed plastic wrap. Because of the high moisture level and air-tight environment, the forage ferments and is preserved by acid production during fermentation. This method has certain advantages and disadvantages versus other forage harvesting and preservation systems.

Advantages of Baleage

- Decreased curing time needed from cutting to baling makes weather less of a factor in forage harvesting
- Potential for more timely harvest of large quantities of forage
- Decreased need for mechanical handling and time curing to dry the forage reduces the loss of leaves, the most digestible part of the plant
- Potential for higher feed quality bale through leaf preservation and possible nitrate reduction

Disadvantages

- Increased harvest cost per bale versus conventional cured hay
- Disposal of used plastic wrap
- More likely to spoil as compared with silage in traditional silos
- Risk of forage spoilage if integrity of wrap is not maintained; birds and rodents can puncture plastic, and holes must be covered
- Transportation of bales is limited due to cost of moving high-moisture bales

Table 5. Feeding values of small-grain and corn silage.

Silage	Dry Matter	percent dry matter		
		Crude Fiber	Crude Protein	TDN
Barley	38.8	27.1	9.0	64.3
Wheat	39.4	27.9	9.6	63.8
Oats	40.2	31.2	9.8	60.7
Rye (wilted)	39.8	33.0	12.8	58.5
Corn	37.3	24.7	8.1	66.4

Source: Virginia Polytechnic Institute and State University

Making Baleage

The forage is cut as if for making hay, but it is baled at 50 to 60 percent moisture rather than at 18 to 20 percent moisture. Baling at the proper moisture content is the single most important variable. Baling haylage with too much moisture reduces the feed quality of the forage and reduces the amount of dry matter stored per bag, greatly increasing the storage cost. Baling haylage with inadequate moisture reduces fermentation and increases mold production, greatly increasing storage losses.

Storage

Successful storage depends on many factors. The storage site should be cleared of stubble and sharp objects. Some people lay a piece of plastic on the ground prior to placing the bales. Rodents can chew through the plastic wrap or bag, which will increase storage losses greatly. Spray the perimeter of the stack to kill weeds, which harbor rodents and insects. Find a shady area, preferably on a north-facing slope, to avoid temperature fluctuations that can degrade the haylage and plastic.

If you find holes in the bagged bales, patch them as soon as possible because wind causes loose plastic to bellow out, providing an air exchange that usually spoils most of the outer layer of the bale. Bags are rarely reusable because of minor pinholes.

Ensiling forage as baleage can be a management strategy for high-nitrate grass. Scientific literature suggests an average of 50 percent reduction in nitrate in ensiled forages due to the de-nitrification process.

Feeding Baleage

Feeding baleage is similar to feeding large, round bales of hay in that conventional feeding rings can be used. With the high investment in wrapping bales, controlling feeding losses is essential. Some studies have shown up to a 50 percent loss when large, round silage bales are fed to cattle without being placed in a ring feeder. This loss can be reduced to 10 to 20 percent by using a simple ring feeder. The use of an elevated hay wagon can reduce feeding losses to below 10 percent.

When feeding individually wrapped haylage bales to any species, feeding to a sufficient number of animals to eat an entire bale within one or two days is best. If multiple bales are ensiled in a plastic tube, the tube may be opened to remove individual bales and resealed without significant spoilage for up to two weeks.

Other publications in the Quality Forage series

- AS1250 “Forage Nutrition for Ruminants”
- AS1251 “Interpreting Composition and Determining Market Value”
- AS1253 “Corn Silage Management”
- AS1254 “Silage Fermentation and Preservation”
- AS1255 “Storage, Sampling and Measuring”
- AS1256 “Stressed or Damaged Crops”

References

In addition to sources cited, materials were adapted with permission from Pioneer Forage Manual, which no longer is in print.

For more information on this and other topics, see www.ag.ndsu.edu

NDSU encourages you to use and share this content, but please do so under the conditions of our Creative Commons license.

You may copy, distribute, transmit and adapt this work as long as you give full attribution, don't use the work for commercial purposes and share your resulting work similarly. For more information, visit www.ag.ndsu.edu/agcomm/creative-commons.

North Dakota State University does not discriminate on the basis of age, color, disability, gender expression/identity, genetic information, marital status, national origin, public assistance status, sex, sexual orientation, status as a U.S. veteran, race or religion. Direct inquiries to the Vice President for Equity, Diversity and Global Outreach, 205 Old Main, (701) 231-7708.

County Commissions, NDSU and U.S. Department of Agriculture Cooperating.

This publication will be made available in alternative formats for people with disabilities upon request, (701) 231-7881.

400-1-06; 200-7-13