

# Selected Management Factors for Economically Increasing Soybean Yield



**Greg Endres**

Extension Area Agronomist  
Carrington Research Extension Center

**Hans Kandel**

Extension Agronomist  
Fargo



Other researchers contributing to this study include:

Carrington Research Extension Center –  
Blaine Schatz, Mike Ostlie and Steve Metzger

Fargo/Prosper –  
Chad Deplazes and Sam Markell

Photos by Greg Endres and Hans Kandel

About 3 million acres of soybean were planted in North Dakota in 2003. That number reached 4.7 million acres a decade later in 2012. The growth in acreage can be attributed to the relative ease in producing the crop and its profitability.

Based on NDSU Extension Service crop budgets, projected return to labor and management during 2012-14 ranged from \$73 to \$114/acre in the east-central region and \$63 to \$122/acre in the southern Red River Valley.

Opportunities exist for farmers to increase soybean yield and profitability by improving plant establishment practices. Also, numerous special inputs that may add to soybean profits are being marketed.

**A study was conducted by NDSU with selected intensive management practices and inputs to examine potential increases in yield and profit.**

## Materials and Methods

Nine research trials were conducted during 2008-11 at the NDSU Carrington Research Extension Center and research sites in Prosper and Fargo to examine inputs that potentially could increase soybean seed yield and profitability. The trials at Carrington were conducted on dryland conditions during 2008 and under irrigation during the entire study period. Trials were conducted at Prosper in 2009 and 2010, and in untilled and tilled environments at Fargo in 2011.

The agronomic response of early and late-maturing varieties was examined using the production factors of planting rates, row spacings and special foliar inputs. Two Roundup Ready varieties, with maturity group ranges of 0.0 to 0.2 and 0.6 to 0.8, were planted each year (Table 1). Also, two planting dates were examined at Carrington in 2008.

Planting rates were 150,000 and 200,000 pure live seeds (pls)/acre. Row spacings were 14 and 28 inches. Special foliar inputs included:  
1) TJ Technologies micronutrient mix "Sunflower/Canola/Soybean" at 48 fluid ounces (fl oz)/acre plus Novozyme LCO Promoter (experimental product in 2008 and "Ratchet" at 4 fl oz/acre in 2009 to 2011) during second- to third-trifoliolate (V2 to V3) growth stages and 2) BASF fungicide "Headline" at 6 fl oz/acre + NIS at 0.125 percent volume/volume during flowering to early pod development (R2 to R3) growth stages.

Other best management practices were used for soybean production during the study. Seed was inoculated with *Bradyrhizobium japonicum*.

## Results and Discussion

The study's average early season (two to four weeks after planting) established soybean plant density was 130,500 plants/acre with the planting rate of 150,000 pls/acre and 163,800 plants/acre with 200,000 pls/acre (Table 2). The percentage

of established plants compared with planting rate was 87 percent with the low planting rate and 82 percent with the high planting rate.

Table 3 indicates soybean seed yield with row spacings, planting rates

and special foliar inputs for each trial. Averaged across the nine trials, the study indicates the following yield advantages:

- 1.4 bushels (bu)/acre with 14-inch versus 28-inch rows

**Table 1. Varieties and planting dates in soybean intensive management study, North Dakota, 2008-11.**

Trial	Soybean Variety		Planting Date	
	Maturity		1st	2nd
	early	late		
Carrington 2008 – dryland	DSR0401	NKS08C3	May 16	May 24
Carrington 2008 – irrigated	DSR0401	NKS08C3	May 16	May 24
Carrington 2009 – irrigated	RT0268	RT0669	May 22	x
Prosper 2009	RT0268	RT0669	May 28	x
Carrington 2010 – irrigated	PFS1000	PFS0806	May 14	x
Prosper 2010	PFS1000	PFS0806	May 20	x
Carrington 2011 – irrigated	PFS1002	DSR0747R2Y	May 23	x
Fargo 2011 – untilled	PFS1002	DSR0747R2Y	May 19	x
Fargo 2011 – tilled	PFS1002	DSR0747R2Y	May 19	x

**Table 2. Planting rate and established stand in soybean intensive management study, North Dakota, 2008-11.**

	Planting Rate (pls/acre)	
	150,000	200,000
	plants/acre <sup>1</sup>	
Carrington 2008 – dryland	143,500 <sup>b</sup>	163,100 <sup>a</sup>
Carrington 2008 – irrigated	141,700 <sup>b</sup>	182,300 <sup>a</sup>
Carrington 2009 – irrigated	151,800 <sup>b</sup>	194,200 <sup>a</sup>
Prosper 2009	119,000 <sup>b</sup>	155,600 <sup>a</sup>
Carrington 2010 – irrigated	132,900 <sup>b</sup>	167,200 <sup>a</sup>
Prosper 2010	137,200 <sup>b</sup>	177,900 <sup>a</sup>
Carrington 2011 – irrigated	147,700 <sup>b</sup>	182,600 <sup>a</sup>
Fargo 2011 – untilled	94,100 <sup>b</sup>	119,200 <sup>a</sup>
Fargo 2011 – tilled	103,400 <sup>b</sup>	135,200 <sup>a</sup>
Study average	130,500 <sup>b</sup>	163,800 <sup>a</sup>

<sup>1</sup> Different letters in the same row indicate statistical differences (LSD=0.05).

**Table 3. Yield and net return with main factors in soybean intensive management study, North Dakota, 2008-11.**

Trial	Yield (bu/acre) <sup>1</sup>					
	Row Spacing (inches)		Planting Rate (pls/acre)		Special Foliar Inputs	
	14	28	150,000	200,000	Yes	No
Carrington 2008 – dryland	36.9 <sup>a</sup>	35.5	36.2	36.2	37.1 <sup>a</sup>	35.4
Carrington 2008 – irrigated	42.7	42.0	41.9	42.7	43.6 <sup>a</sup>	41.0
Carrington 2009 – irrigated	48.3	48.2	46.5	50.0	48.8	47.7
Prosper 2009	53.8	53.7	55.5	51.9	54.7	52.8
Carrington 2010 – irrigated	63.9	62.5	62.5	63.9	64.7 <sup>a</sup>	61.7
Prosper 2010	55.2	54.5	53.6	56.0	55.7	54.0
Carrington 2011 – irrigated	52.4 <sup>a</sup>	48.1	49.6	50.9	51.5 <sup>a</sup>	49.0
Fargo 2011 – untilled	44.5	42.5	41.3	45.7 <sup>a</sup>	44.7	42.4
Fargo 2011 – tilled	51.5	49.9	48.6	52.8 <sup>a</sup>	52.0	49.4
Study average	49.9 <sup>a</sup>	48.5	48.5	49.9 <sup>a</sup>	50.3 <sup>a</sup>	48.1
Net revenue/acre <sup>2</sup>	\$499	\$485	\$435	\$432	\$466	\$481

<sup>1</sup> 'a' indicates mean within a factor is statistically significant (LSD=0.05) by trial and study average.

<sup>2</sup> Assumptions: \$10/bu soybean market price; seed costs: \$0.30/1,000 seeds; \$5/acre field application cost; \$32/acre special foliar input cost.

- 1.4 bu/acre with planting 200,000 pls/acre versus 150,000 pls/acre
- 2.2 bu/acre with special foliar inputs versus untreated check

Assumptions used for determining net revenue after costs of research factors include \$10/bu soybean market price; no difference in costs between row spacings; 30 cents/1,000 seeds with 10 percent additional seed used to reach desired pls/acre; total cost of \$32/acre for special foliar inputs plus \$5/acre for a field application cost. Averaged across the nine trials, the study indicates the following revenue advantages:

- \$14/acre with 14-inch vs. 28-inch rows
- \$3/acre with the low vs. high planting rate
- \$15/acre without special foliar inputs

The Fargo 2011 research site compared soybean performance with tilled and untilled conditions using controlled drainage structures (Table 4).

Early vigor was slightly greater, plants were taller and iron chlorosis expression was slightly lower in the tilled treatment. Seed yield with tilled was 13 percent higher than untilled yield.

**Table 4. Response to tile versus nontile environments in soybean intensive management study, North Dakota, Fargo, 2011.**

Drainage <sup>1</sup>	Plant			
	Vigor <sup>2</sup>	Height	IDC <sup>3</sup>	Yield
	(1-9)	(inch)	(1-5)	(bu/acre) <sup>4</sup>
Tiled	5.5 <sup>a</sup>	27.5 <sup>a</sup>	2.6 <sup>b</sup>	50.7 <sup>a</sup>
Untilled	4.9 <sup>b</sup>	23.6 <sup>b</sup>	2.9 <sup>a</sup>	43.9 <sup>b</sup>

<sup>1</sup> Two environments: tilled and untilled in Fargo 2011.

<sup>2</sup> Early season vigor; 1 = least vigorous and 9 = most vigorous.

<sup>3</sup> IDC = Iron Chlorosis Deficiency symptoms; 1 = tissue green and 5 = tissue yellow/brown. Scoring conducted on June 30.

<sup>4</sup> Different letters in the same column indicate statistical differences (LSD=0.05).

Soybean yield was 2.6 bu/acre (5 percent) greater with the later maturity group (50.5 bu/acre), compared with the early maturity group (47.9 bu/acre). The highest



Fourteen-inch rows and two varieties, Carrington, 2009.

yield with the combination of maturity groups and row spacing was with the later-maturing varieties grown in 14-inch rows (Table 5).

**Table 5. Yield with maturity groups and row spacing in soybean intensive management study, North Dakota, 2008-11.**

Maturity Group	Row Spacing	Yield
	(inches)	(bu/acre) <sup>1</sup>
0.6-0.8	14	51.7 <sup>a</sup>
0.6-0.8	28	49.4 <sup>b</sup>
0.0-0.2	14	48.1 <sup>c</sup>
0.0-0.2	28	47.7 <sup>c</sup>

<sup>1</sup> Letters indicate statistical differences (LSD = 0.05).

Table 6 displays average date of canopy closure, lowest pod height, and yield with row spacing and planting rates.

**Table 6. Response to row spacing and planting rate in soybean intensive management study, North Dakota, 2008-11.**

Row Spacing	Planting Rate	Canopy Closure	Lowest Pod Height <sup>2</sup>	Yield
(inches)	(pls/acre)	(Julian day) <sup>1</sup>	(cm)	(bu/acre) <sup>3</sup>
14	150,000	205 <sup>a</sup>	5.8 <sup>ab</sup>	49.3 <sup>b</sup>
14	200,000	205 <sup>a</sup>	6.2 <sup>a</sup>	50.5 <sup>a</sup>
28	150,000	241 <sup>b</sup>	5.4 <sup>b</sup>	47.8 <sup>c</sup>
28	200,000	239 <sup>b</sup>	6.1 <sup>a</sup>	49.3 <sup>b</sup>


<sup>1</sup> Seven environments (no data recorded for Fargo 2011). Julian day = day of the year.

<sup>2</sup> Six environments (no data recorded for Prosper 2010 and Fargo 2011).

<sup>3</sup> Different letters in the same column indicate statistical differences (LSD=0.05).

Averaged across seven trials, canopy closure with 14-inch rows was 35 days earlier, compared with 28-inch rows. The early canopy closure with the narrow rows provided the opportunity for increased capture of sunlight, increased crop competitiveness with weeds, decreased soil moisture loss from evaporation and ultimately yield increase, compared with the wide rows.

Canopy closure occurred sooner with 14-inch rows but was similar between planting rates. Pod height generally increased slightly with the high planting rate. Yield was highest with 14-inch rows and the high planting rate.



14-inch rows (left) and 28-inch rows (right) at 150,000 pls/A in late July 2009 at Prosper.

Soybean yield and net revenue after costs of research factors with the combination of row spacing, planting rates and special foliar inputs are listed in **Table 7**. The highest yield was obtained with foliar inputs in 14-inch

rows at both planting rates and in 28-inch rows at the high planting rate. However, the combination of 14-inch rows and 150,000 pls/acre planting rate without foliar inputs provided the highest net revenue.

**Table 7. Yield and net revenue with three factor interactions in soybean intensive management study, North Dakota, 2008-11.**

Row Spacing	Planting Rate	Special Foliar Inputs	Yield	Net Revenue After Costs of Research Factors/Acre
(inches)	(pls/acre)		(bu/acre) <sup>1</sup>	(\$)
14	150,000	yes	50.2 <sup>ab</sup>	465
14	150,000	no	48.4 <sup>c</sup>	484
14	200,000	yes	51.7 <sup>a</sup>	463
14	200,000	no	49.3 <sup>bc</sup>	476
28	150,000	yes	48.9 <sup>bc</sup>	452
28	150,000	no	46.7 <sup>d</sup>	467
28	200,000	yes	50.4 <sup>ab</sup>	450
28	200,000	no	48.2 <sup>cd</sup>	465


<sup>1</sup> Different letters in the yield column indicate statistical differences (LSD = 0.05).

<sup>2</sup> Assumptions: \$10/bu soybean market price; seed costs: \$0.30/1,000 seeds; \$5/acre field application cost; \$32/acre special foliar input cost.

## Summary

- Soybean yield increased with 14-inch versus 28-inch rows, 200,000 versus 150,000 pls/acre planting rate and special foliar inputs. The narrow rows also had higher net revenue than 28-inch rows. However, the low planting rate and no foliar inputs provided higher net revenue after costs of research factors versus the alternative choice for each factor.
- While several combinations of the three main factors in the study provided high yields, the combination of 14-inch rows and the 150,000 pls/acre planting rate without foliar inputs provided the highest net revenue after costs of research factors.
- While using pure live seed as a basis for reaching a targeted plant population is recommended, also consider that an additional 10 to 20 percent seedling loss can occur between planting and emergence.
- Management strategies that reduce the time required from planting to canopy closure will increase yield potential. This study indicated that canopy closure with narrow rows occurred more than a month earlier than with wide rows.

Researched with support from the



The NDSU Extension Service does not endorse commercial products or companies even though reference may be made to trade names, trademarks or service names.

**For more information on this and other topics, see [www.ag.ndsu.edu](http://www.ag.ndsu.edu)**

NDSU encourages you to use and share this content, but please do so under the conditions of our Creative Commons license. You may copy, distribute, transmit and adapt this work as long as you give full attribution, don't use the work for commercial purposes and share your resulting work similarly. For more information, visit [www.ag.ndsu.edu/agcomm/creative-commons](http://www.ag.ndsu.edu/agcomm/creative-commons).

North Dakota State University does not discriminate on the basis of age, color, disability, gender expression/identity, genetic information, marital status, national origin, public assistance status, sex, sexual orientation, status as a U.S. veteran, race or religion. Direct inquiries to the Vice President for Equity, Diversity and Global Outreach, 205 Old Main, (701) 231-7708.

County Commissions, NDSU and U.S. Department of Agriculture Cooperating. This publication will be made available in alternative formats for people with disabilities upon request, (701) 231-7881.