

Photo credits

All photos in the publication are by Rodney G. Lym, North Dakota State University, except as noted below.

Black henbane plant and rosette courtesy of Steve Dewey, Utah State University. Flower photograph courtesy of Washington State Noxious Weed Control Board.

False chamomile full plant from Richard Old, XID Services Inc.

Common tansy photos courtesy of Celestine Duncan, Weed Management Services, Helena, Mont.

Downy brome infestation photo courtesy of Leslie J. Mehrhoff, IPANE. Planicle photo courtesy of J.C. Shour, Biopix.dk. Ligule photo from Virginia Tech Weed Guide.

Halogeton photos courtesy of Blake Schaan, North Dakota Department of Agriculture, Bismarck, N.D.

Hoary cress flower and seed pod photos courtesy of Steve Dewey, Utah State University. Bolted plant photo from Richard Old, XID Services Inc.

Diffuse knapweed, spotted knapweed and Russian knapweed flower bract photos courtesy of Steve Dewey, Utah State University.

Kochia seedling courtesy of Richard Zollinger, NDSU.

Aphthona flea beetle photo from Biological Control of Leafy Spurge CD, USDA-APHIS and USDA-ARS, Team Leafy Spurge's IPM information series No. 1.

Scotch thistle photos of flower and leaf and whole plant courtesy of Steve Dewey, Utah State University. Flower close-up from Richard Old, XID Services Inc.

Dalmatian and yellow toadflax photos courtesy of Steve Dewey, Utah State University.

Yellow starthistle hairy weevil from Mark Schwarzländer, University of Idaho.

Composite photos of thistle plant flowers and leaves courtesy of Jason Adams, NDSU.