

Examples of Label Types

Examples of Package Insert Information

<p>COWBIOTIC</p> <p>(hydrocillin and streptazolidin in aqueous suspension) For use in beef cattle, lactating and non-lactating dairy cattle, and swine</p>	Species and Animal Class															
<p>Read entire brochure carefully <i>Before using this product.</i></p> <p>For Intramuscular Use Only</p> <p>Composition: Cowbiotic is an effective antimicrobial preparation containing hydrocillin and streptozoldin. Each ml of this suspension contains 200,000 units of hydrocillin and 250 mg of streptazolidin. The combination permits treatment of many mixed bacterial infections with the convenience of a single dosage form.</p>	Approved Uses															
<p>Indications: Cattle: Bronchitis; footrot; leptospirosis; mastitis; meritis; pneumonia; wound infections and other infection caused by or associated with hydrocillin – and streptazolidin – susceptible organisms.</p>	Approved Uses															
<p>RECOMMENDED DAILY DOSAGE</p> <p>Continue treatment for 1 to 2 days after symptoms disappear.</p> <table border="0"> <tr> <td style="text-align: left;">CATTLE</td> <td style="text-align: left;">Body Weight</td> <td style="text-align: left;">Dosage</td> </tr> <tr> <td></td> <td>Up to 100 lbs</td> <td>2 ml</td> </tr> <tr> <td></td> <td>100 to 300 lbs</td> <td>2 to 6 ml</td> </tr> <tr> <td></td> <td>300 to 700 lbs</td> <td>6 to 10 ml</td> </tr> <tr> <td></td> <td>700 lbs or more</td> <td>10 to 14 ml</td> </tr> </table>	CATTLE	Body Weight	Dosage		Up to 100 lbs	2 ml		100 to 300 lbs	2 to 6 ml		300 to 700 lbs	6 to 10 ml		700 lbs or more	10 to 14 ml	Dosage
CATTLE	Body Weight	Dosage														
	Up to 100 lbs	2 ml														
	100 to 300 lbs	2 to 6 ml														
	300 to 700 lbs	6 to 10 ml														
	700 lbs or more	10 to 14 ml														
<p>CAUTION: 1. Cowbiotic should be injected deep within the fleshy muscle of the neck. Do not inject this material in the hip or rump, subcutaneously, into a blood vessel, or near a major nerve. 2. If improvement does not occur within 48 hours, the diagnosis should be reconsidered and appropriate treatment initiated. 3. Treated animals should be closely observed for at least one-half hour. Should a reaction occur, discontinue treatment and administer epinephrine and antihistamines immediately. 4. COWBIOTIC must be stored between 2 to 8 degrees C (36 to 46 degrees F). Warm to room temperature and shake well before using. Keep under refrigeration when not in use.</p>	Route of Administration															
<p>Warning: Milk that has been taken from animals during treatment and for 48 hours (4 milkings) after the latest treatment must not be used for food. The use of this drug must be discontinued for 30 days before treated animals are slaughtered for food.</p>	Additional Information															
<p>How Supplied: Cowbiotic is available in vials of 100 ml and 250 ml.</p>	Storage Requirements															
	Withdrawal Times															
	Sizes Available															