

Hard Red Spring Wheat Recrop - 2005

Williston Research Extension Center

Cultivar	Heading Date fr plntg	Plant Height in	Test Weight lbs/b	Grain Protein %	Grain Yield			
					2003	2004	2005	3 yr avg
					----- bu/ac -----			bu/ac
<b>Conventional</b>								
Amidon	66	30	60.4	15.5	50.2	49.2	33.0	44.1
Dapps	67	29	58.8	18.1	51.3	44.3	25.5	40.4
Ernest	66	31	60.2	17.4	48.8	43.7	28.2	40.2
Glenn	65	26	62.6	16.7	54.0	44.9	32.4	43.8
Parshall	66	26	60.3	16.8	50.5	45.8	34.1	43.5
<b>Semi-dwarf</b>								
Alsen	66	25	61.5	16.1	54.8	45.4	35.3	45.2
Briggs	66	26	61.1	16.5	53.5	46.4	35.7	45.2
Choteau	65	27	60.8	15.9	--	44.4	37.8	--
Freyr	66	30	61.8	15.7	--	48.4	40.1	--
Granger	66	28	60.6	15.9	--	50.8	34.8	--
Granite	68	27	60.1	17.4	--	45.2	31.2	--
Knudson	67	25	60.9	16.9	57.0	46.7	28.9	44.2
McNeal	67	29	57.9	15.6	49.1	48.1	37.7	45.0
Norpro	67	24	60.9	16.0	55.4	49.6	37.0	47.3
Oklee	65	22	61.7	16.0	--	48.4	35.1	--
Outlook	67	29	61.0	16.8	57.1	49.0	33.8	46.6
Reeder	67	26	60.7	15.9	58.9	49.7	40.3	49.6
Steele-ND	66	29	61.1	15.4	51.7	43.4	37.1	44.1
<b>Hard White</b>								
AC Vista	66	26	58.6	14.7	--	55.0	34.9	--
Mean	66	27	60.5	16.3	53.0	46.9	34.3	--
CV%	1	8	1.6	5.7	7.3	4.1	13.9	--
LSD 0.05	1	4	2.0	NS	NS	3.2	7.8	--

Planting Date: April 21, 2005 into tilled recrop

Harvest Date: August 8, 2005

Grain protein percentages reported on a 12% moisture basis