

**NDSU Carrington Research Extension Center  
2006 Variety Trial Data**

<b>Soybean- Direct Seeded</b>	<b>Wishek</b>
-------------------------------	---------------

Brand	Variety	Mat. <sup>-1</sup> Group	Type	-- Seed Yield -								
				Pod Ht cm	Plant Ht inch	Days to PM	Seed Oil* %	Seed Protein* %	Seeds/ pound	Test Weight lb/bu	2006	2-yr. Avg. bu/ac
NDSU	Walsh	0.2	Conv.	7	13.0	120.3	17.3	39.0	2477	57.8	20.5	35.8
NDSU	Barnes	0.3	Conv.	8	14.6	119.0	18.8	38.2	2358	56.7	21.8	41.5
NDSU	LaMoure	0.7	Conv.	7	14.0	122.5	17.0	38.8	2571	57.7	18.6	36.5
Roughrider Genetics	RG6008RR	0.8	RR	6	12.2	107.0	17.6	39.7	2623	57.0	24.6	37.4
Roughrider Genetics	RG600RR	0.0	RR	8	14.4	117.3	18.5	39.0	2419	57.5	28.5	35.9
Roughrider Genetics	RG601NRR	0.1	RR	7	13.2	116.3	18.1	38.6	2621	57.8	27.4	38.2
Roughrider Genetics	RG603RR	0.2	RR	7	16.3	118.8	17.7	39.5	2548	57.5	33.6	34.0
Roughrider Genetics	RG604RR	0.3	RR	7	16.9	120.3	18.2	38.0	2532	57.2	37.1	42.3
Roughrider Genetics	RG607RR	0.8	RR	7	17.7	122.3	17.2	39.8	2354	58.0	29.5	--
Dairyland Seed	DSR-0401/RR	0.4	RR	9	15.9	119.8	18.3	38.0	2417	57.0	30.3	--
Pioneer	90M61	0.6	RR	5	13.4	122.0	18.5	38.5	2315	57.0	30.3	--
Croplan Genetics	RTO669	0.6	RR	6	16.1	119.0	18.1	38.4	2940	56.8	31.4	--
Proseed	RR 0069	0.6	RR	8	15.0	118.5	18.4	37.0	2052	56.9	35.9	--
NK Brand	S06-D9	0.6	RR	6	13.2	119.8	18.3	39.2	2313	57.3	27.6	--
NuTech Seed	NT-0660+RR	0.5	RR	7	18.7	120.0	19.0	36.6	2594	56.5	42.6	--
Kruger	K-057RR	0.5	RR	7	18.3	121.3	18.3	37.2	2471	57.1	39.1	--
Legend	LS 0624RR	0.6	RR	8	19.5	119.0	18.3	37.8	2926	56.5	40.7	--
Asgrow	AG 0604	0.6	RR	9	18.9	121.8	18.6	37.6	2562	57.4	36.9	--
Prairie Brand	PB-0636RR	0.6	RR	9	17.3	121.8	18.2	37.6	2466	57.4	41.8	--
MTX	Exp B05-J	0.5	RR	6	16.7	117.5	18.2	38.0	2591	57.5	35.2	--
MEAN				6.9	15.8	119.2	18.2	38.3	2507	57.2	32.2	--
C.V.%				26.9	15.7	1.0	1.9	1.7	3.4	1.1	18.6	--
LSD.05				NS	3.5	1.7	0.5	0.9	120	0.9	8.5	--

**Planting Date = May 23 ; Harvest Date = October 5 ; Previous Crop = Spring Wheat**

1; Maturity group based on most current company information that has been provided.

\* Seed oil and protein contents reported on a 13% moisture basis.

This trial was direct seeded into a non-tilled spring wheat residue.