

**NDSU Carrington Research Extension Center
2008 Variety Trial Data**

Hard Red Spring Wheat - Direct Seeded Dryland Recrop Elite	Carrington
---	-------------------

Variety	Days to Head	Plant Lodge	Plant Height	Leaf Disease %	-- Grain Protein --		1000 KWT gram	Test Weight lb/bu	----- Grain Yield -----		
					2008 %	3-yr. Avg. %			2008 bu/ac	3-yr. Avg. bu/ac	
Ada	76.0	0.3	32.4	12	14.9	15.4	32.0	59.7	58.2	43.8	43.8
Alsen	73.5	0.0	33.8	7	16.2	16.0	28.8	58.1	54.1	42.6	43.1
AP 604CL	74.3	0.0	32.1	17	15.9	--	26.9	59.3	50.8	41.0	--
Bigg Red	76.5	0.8	38.1	15	15.5	15.0	29.5	60.5	60.9	41.5	44.4
Briggs	72.8	0.3	34.1	12	15.8	15.8	28.8	58.6	60.6	46.4	46.7
Dapps	73.0	1.3	36.9	20	16.7	16.6	28.0	56.5	51.7	41.6	41.3
Faller	76.5	0.0	34.0	8	15.4	15.2	30.0	57.1	55.7	47.2	46.6
FBC Dylan	74.8	0.3	32.9	12	15.9	15.3	30.3	57.1	55.3	40.8	42.5
Freyr	75.3	0.0	33.8	7	15.5	15.4	27.6	57.6	57.7	45.4	46.0
Glenn	72.8	0.5	36.1	11	16.5	16.0	28.0	61.1	53.6	46.2	46.3
Granger	75.5	0.8	37.8	17	15.8	15.5	34.3	59.2	62.9	49.2	47.8
Howard	75.3	0.0	34.2	8	15.7	15.5	27.8	57.3	57.8	46.7	46.5
Kelby	73.8	0.0	32.9	22	15.6	15.8	28.2	56.9	59.0	44.8	45.5
Knudson	76.8	0.0	32.1	11	15.0	15.4	29.5	57.9	63.5	48.4	47.4
Kuntz	74.3	0.0	28.5	17	14.7	--	27.0	59.0	52.5	41.2	--
ND 901CL	75.3	0.0	37.2	12	16.4	--	29.0	58.8	55.0	41.7	--
Oklee	72.5	0.5	34.4	17	16.3	--	29.3	59.5	61.5	--	--
Parshall	74.5	0.5	38.7	14	16.3	15.9	28.3	58.4	57.3	43.3	43.6
RB07	73.3	0.0	31.2	11	15.1	--	28.3	59.3	61.7	48.3	--
Reeder	73.3	0.0	31.1	13	16.2	15.6	28.1	56.5	55.6	41.1	43.2
Rush	73.8	0.0	33.2	17	16.3	16.3	28.2	60.2	54.6	39.9	40.0
Steele-ND	74.0	0.5	34.2	12	15.7	15.7	28.4	56.9	52.0	44.1	44.9
Tom	74.3	0.3	31.4	11	15.5	--	31.3	58.8	55.0	--	--
Traverse	72.8	0.0	34.5	23	15.8	15.2	27.2	55.4	57.1	51.1	50.7
Ulen	74.0	0.3	33.1	21	15.8	15.6	29.6	58.1	58.9	43.3	44.2
MEAN	74.7	0.2	34.0	12.9	15.7	---	29.0	58.1	57.2	---	---
C.V.%	1.4	158	4.8	35.0	2.5	---	4.7	1.5	9.3	---	---
LSD.05	1.5	0.5	2.3	6.3	0.6	---	1.9	1.2	7.5	---	---

Planting Date = April 17 ; Harvest Date = August 12 ; Previous Crop = Spring Wheat

This trial was directly planted into an undisturbed wheat stubble.