

**NDSU Carrington Research Extension Center  
2012 Variety Trial Data**

<b>Winter Triticale Forage</b>																<b>Carrington</b>
Variety	Early	Jday of	Plant	Harvest <sup>3</sup>	----- Yield -----		CP	ADF	NDF	CA	P	MG	K	S	TDN	RFV
	Vigor <sup>1</sup>	Heading <sup>2</sup>	Height	Moisture	15 % M	DM										
	0-10		inch	%	----- ton/ac -----											
Brisk	3.8	154.3	56.1	64.5	4.26	3.62	10.1	43.5	65.6	0.37	0.33	0.15	3.08	0.15	55.0	78.1
Fridge	8.3	156.0	55.1	66.9	4.21	3.58	9.4	44.3	66.9	0.45	0.34	0.16	3.14	0.13	54.4	75.6
Shiveree 185	2.8	154.8	52.0	63.9	3.03	2.58	8.9	41.9	64.2	0.36	0.33	0.15	2.97	0.13	56.2	81.7
Shiveree 186	9.3	156.3	53.9	67.7	4.57	3.88	10.3	43.5	66.0	0.47	0.34	0.16	3.19	0.16	55.0	77.5
Childeric	8.3	156.5	53.5	66.3	4.00	3.40	8.8	44.1	67.5	0.37	0.34	0.15	3.04	0.13	54.5	75.1
Mean	6.5	155.6	54.2	65.9	4.01	3.41	9.5	43.5	66.0	0.40	0.34	0.15	3.08	0.14	55.1	77.6
C.V.%	30.8	0.4	4.5	2.9	27.0	27.0	9.4	2.3	1.8	13.0	2.7	6.6	2.4	11.1	1.4	3.2
LSD.05	3.1	0.9	3.7	2.9	NS	NS	1.4	1.6	1.8	0.08	0.01	NS	0.12	0.02	1.2	3.8

**Planting Date = September 15; Harvest Date = June 15; Previous Crop = Barley**

<sup>1</sup> Vigor taken May 3 with 1 = poor and 10 = excellent.

<sup>2</sup> Jday 155 = June 3

<sup>3</sup> Harvest stage: all treatments in early anthesis.

No plant lodging occurred in this trial.