

| Cultivar | Winter | Days to | Plant | . Test | Grain | 2006 | Grain Yield | | |
|------------------|----------|---------|--------|--------|---------|-----------------|-------------|------|----------|
| | Survival | Head | Height | Weight | Protein | | 2007 | 2008 | 3 yr avg |
| | % | fr 1/1 | in | lbs/bu | % | -----bu/ac----- | | | |
| RYE | | | | | | | | | |
| DR02 | 71.5 | 30.0 | 30.0 | 51.3 | 14.4 | -- | 86.2 | 20.2 | -- |
| DR0207 | 93.8 | 30.8 | 24.7 | 51.9 | 15.9 | -- | -- | 20.8 | -- |
| Remington | 55.0 | 30.0 | 29.4 | 51.7 | 16.0 | -- | 77.7 | 21.3 | -- |
| Rymin | 94.5 | 29.5 | 25.9 | 53.2 | 15.9 | -- | -- | 20.6 | -- |
| SPELT | | | | | | | | | |
| Frank | 66.3 | 49.8 | 27.6 | 27.9 | 16.3 | 32.6 | 71.4 | 30.6 | 44.9 |
| PI348145 | 71.3 | 50.0 | 27.4 | 24.4 | 15.7 | 31.2 | 70.0 | 25.0 | 42.1 |
| PI348159 | 75.0 | 50.0 | 28.9 | 28.3 | 16.5 | 38.1 | 68.8 | 27.4 | 44.8 |
| PI348377 | 48.8 | 49.3 | 23.2 | 24.0 | 15.9 | -- | 68.0 | 22.7 | -- |
| TRITICALE | | | | | | | | | |
| Boreal | 70.0 | 38.5 | 37.7 | 43.6 | 24.7 | 31.5 | 76.5 | 19.2 | 42.4 |
| HR003 | 86.3 | 35.5 | 23.2 | 50.8 | 25.9 | -- | 72.0 | 20.7 | -- |
| NE426GT | 86.3 | 33.5 | 22.9 | 52.8 | 24.2 | -- | -- | 22.7 | -- |
| Windrift | 56.3 | 36.5 | 33.6 | 48.6 | 24.0 | 34.4 | 73.6 | 14.7 | 40.9 |
| EXP MEAN | 72.9 | 38.6 | 27.9 | 42.4 | 18.8 | 32.8 | 69.4 | 22.2 | -- |
| C.V. % | 28.9 | 2.8 | 12.4 | 3.8 | 5.1 | 9.1 | 12.0 | 22.6 | -- |
| LSD 5% | NS | 1.6 | 5.0 | 3.5 | 2.1 | 4.2 | 12.7 | 7.2 | -- |

Planting Date: September 13, 2007 into durum stubble.

Harvest Date: July 22

Bushels/acre figured on 56, 50 and 40 pounds/bushel test weights for rye, triticale, and spelt respectively.

Analyzed in lbs/ac and converted to bu/ac. No proteins for Spelt cultivars.