## NDSU Carrington Research Extension Center 2005 Variety Trial Data

Rye							Carrington	
						Grain Yield		
	Plant	Plant	Grain	1000	Test		3-yr.	
Variety	Lodge	Height	Protein	KWT	Weight	2005	Avg.	
	0 - 9	inch	%	gram	lb/bu	bu/ac		
AC Remington	2.5	47.9	10.8	21.0	53.6	62.9		
AC Rifle	0.0	37.3	11.3	17.1	51.3	42.4	69.0	
Dacold	1.3	46.1	11.0	21.2	51.8	61.9	81.7	
Hancock	1.8	50.1	11.3	25.3	55.4	66.3		
Musketeer	1.5	49.7	11.3	23.7	55.3	71.9	86.7	
Prima	1.3	49.6	10.9	23.4	54.2	69.8	82.7	
Spooner	1.5	49.1	11.9	23.7	55.2	63.7		
Wheeler	0.5	52.4	16.4	33.8	50.4	28.0		
Mean	1.3	47.8	11.9	23.7	53.4	58.4		
C.V. %	52.9	3.7	2.6	4.2	0.9	7.0		
LSD .05	1.0	2.6	0.5	1.4	0.7	6.0		

<sup>\*\*</sup> At harvest, the variety 'Wheeler' had noticeable ergot in all reps of the harvested sample.

Planting Date = October 8; Harvest Date = July 28; Previous Crop = Flax