

**NDSU Carrington Research Extension Center  
2007 Variety Trial Data**

<b>Camelina</b>	<b>Carrington</b>
-----------------	-------------------

Variety	Days to Bloom	Plant Height inch	Plant Lodge 0-9	Oil Content %	Test Weight lb/bu	----- Grain Yield -----	
						2007 ----- lb/ac -----	3-yr. Avg.
CO46	50.0	23.6	2.3	34.8	48.4	405	737.9
CO54	50.5	23.0	3.8	34.3	49.5	259	540.0
CO54-97	49.0	26.0	0.0	35.3	52.0	1116	927.0
Celine	51.0	27.8	1.8	36.2	51.9	955	--
Galena	50.3	25.4	0.3	36.8	52.3	1097	--
Gold of Pleasure	50.3	24.8	3.5	34.1	50.3	395	823.3
Ligena	49.0	26.2	2.5	36.1	51.0	1030	--
Experimental A	49.8	24.4	0.3	36.0	50.8	935	--
Experimental B	48.0	27.8	1.3	34.5	51.7	999	--
Robby	50.3	24.6	0.0	36.8	52.7	849	--
Robinson	50.5	22.4	1.8	34.0	52.2	582	801.7
MEAN	49.9	25.1	1.6	35.3	51.3	785.5	--
C.V.%	1.4	5.9	59.3	3.1	1.0	27.9	--
LSD.05	1.0	2.1	1.3	1.6	0.8	316	--

**Planting Date = May 4 ; Harvest Date = August 13 ; Previous Crop = Spring Wheat**

\* The varieties within this trial required on average, 83 days from planting to reach physiological maturity.

Lodging Score: 0 = no lodging; 9 = plants lying flat.