NDSU Carrington Research Extension Center 2007 Variety Trial Data

| Crambe | | | | | | | Carrington | |
|---------|------------|---------|---------|--------|---------|------------|------------|--|
| | | | | | | Seed Yield | | |
| | Weathering | Days to | Plant | Test | Oil | | 3-yr. | |
| Variety | Tolerance* | Bloom | Lodge** | Weight | Content | 2007 | Avg. | |
| | 0 to 9 | | 0-9 | lb/bu | % | 1b/ac | | |
| BelAnn | 2.8 | 47.0 | 1.5 | 22.8 | 27.4 | 1851 | 2130 | |
| Meyer | 3.8 | 44.3 | 0.8 | 24.1 | 29.9 | 1571 | 1966 | |
| MEAN | 3.9 | 45.1 | 1.1 | 28.5 | 23.7 | 1705 | | |
| C.V.% | 55.2 | 1.4 | 98.6 | 2.7 | 3.1 | 7.0 | | |
| LSD.05 | 3.1 | 0.9 | NS | 1.1 | 1.0 | NS | | |

Planting Date = May 9; Harvest Date = August 13; Previous Crop = Spring Wheat

^{*} Weathering Tolerance: 0 = no plant breakdown and good color to 9 = plants breaking down and discolored. Differences existed among lines after plants were exposed to a prolonged period of rain.

^{**} Lodging Score: 0 = no lodging; 9 = plants lying flat.