## NDSU Carrington Research Extension Center 2011 Variety Trial Data

Organic Potato - Dryland	Carrington

										Total Yield			
			Days to		Plant	Days to	Tuber	Yield	Yield		3-yr.		
Variety	Type	Skin/flesh color	Emergence <sup>1</sup>	Vigor <sup>2</sup>	Disease <sup>3</sup>	Brown Vines <sup>4</sup>	Weight <sup>5</sup>	В	$#1^{6}$	2011	Avg.		
				1-10	1-10		gram	cwt/ac	cwt/ac	cwt/ac			
Burbank	russet	brown russet/white	33.0	6.3	6.8	115.8	127.7	51.5	131.0	182.5	167.4		
Butte	russet	brown russet/white	35.3	4.5	9.5	123.8	124.1	56.6	116.2	172.8	173.5		
Caribe	specialty	purple/white	32.5	7.5	4.0	106.0	137.6	23.4	167.1	190.5	175.8		
Defander	white	white/white	32.0	6.8	8.3	123.5	95.0	100.6	88.9	189.4			
Kennebec	white	white/white	31.8	8.3	6.8	118.0	186.5	16.0	234.1	250.1	206.0		
Missaukee	white	white netted/white	33.3	5.5	9.8	123.5	94.9	64.7	194.7	259.3			
Russian Banana	fingerling	golden/golden	34.5	1.8	9.0	116.3	42.2		104.1	104.1			
Prairie Blush	yellow	golden/golden	31.8	8.8	2.3	103.8	138.5	13.5	165.5	179.0			
Yukon Gem	yellow	golden/golden	31.3	6.5	4.3	111.3	105.1	25.4	100.5	126.0			
Yukon Gold	yellow	golden/golden	32.3	8.5	3.3	103.8	156.3	10.1	194.3	204.4	173.1		
Jacqueline Lee	yellow	golden/golden	37.5	3.8	9.8	122.0	110.9	112.7	147.3	260.0			
Sangre	red	red/white	34.8	2.5	8.0	120.5	145.6	39.2	158.2	197.4	180.1		
Dark Red Norland	red	red/white	32.0	7.0	1.8	103.3	134.4	17.0	161.5	178.5			
Red Cloud	red	red/white	35.8	2.0	7.5	118.8	160.8	17.9	255.8	273.7			
Red Norland	red	red/white	29.5	8.0	1.0	98.5	123.0	22.2	152.4	174.6	172.9		
Red Pontiac	red	red/white	30.3	6.3	4.0	113.8	194.7	16.5	162.1	178.6			
Mean			32.9	5.9	6.0	113.9	129.8	39.2	158.4	195.1			
C.V. (%)			3.6	27.8	19.4	3.0	16.1	34.1	18.6	15.1			
LSD 0.05			1.7	2.3	1.7	4.9	29.9	19.1	41.9	42.1			

Planting Date = May 10; Harvest Date = September 19; Previous Crop = Cover crop mixture (sorghum/sudan, italian ryegrass, pearl millet, berseem & crimson clover, black lentil, turnip, and rape)

<sup>&</sup>lt;sup>1</sup> Days to emergence = days from planting to 75% plants emerged

<sup>&</sup>lt;sup>2</sup> Vigor indicates early season plant growth, 1=least vigor while 10=greatest

 $<sup>^{3}</sup>$  Trial was infected with an unidentified disease. This disease caused the plants to dry down and die early. Plants were rated on August 16 with 1 = most disease while 10 = least disease.

<sup>&</sup>lt;sup>4</sup> Days to brown vines = days from planting to 95% of vines dry.

<sup>&</sup>lt;sup>5</sup> Tuber weight = weight in grams of an average tuber.

<sup>&</sup>lt;sup>6</sup> Fingerlings were not sized, therefore all yield is reported as #1

<sup>\*</sup> This trial experienced a hail storm on July 24