NDSU Carrington Research Extension Center 2018 Variety Trial Data

Winter Rye	Carrington

										Grain Yield		
		Winter	Jday	Early	Plant	Plant	Ergot	1000	Test		3-yr.	
Variety	Vigor ¹	Survival	of Head	Height ²	Height ²	Lodge	Sclerotia ³	KWT	Weight	2018	Avg.	
	1-10	%		inch	inch	0-9	%	gram	lb/bu	bu	/ac	
Hancock	2.3	66.3	155.3	8.3	36.9	1.3	0.70	27.5	51.1	44.1	51.6	
Spooner	1.0	38.8	154.3	9.0	38.2	2.3	0.42	27.0	49.7	36.6	46.4	
Rymin	8.0	91.3	155.5	8.9	34.5	0.8	0.32	30.7	52.2	61.2	61.1	
ND Dylan	8.5	95.0	154.5	8.3	38.0	2.0	0.18	24.9	51.0	60.2	62.5	
Dacold	6.5	85.8	159.0	8.5	38.6	1.8	0.29	24.6	48.4	52.9	54.6	
Aroostok	3.8	71.0	149.3	11.2	37.8	5.3	0.81	20.8	50.0	29.1	36.4	
Hazlet	8.0	90.8	155.8	8.4	32.9	1.3	0.29	29.7	51.9	55.1	58.1	
Wheeler	4.8	81.3	158.0	9.2	39.9	0.0	5.03	33.0	46.3	18.0	17.3	
Brasetto	8.3	91.8	155.3	8.4	28.8	0.0	0.35	23.2	48.8	58.8		
Mean	6.0	80.8	154.4	9.1	36.3	1.8	0.88	26.4	50.0	46.3		
C.V. (%)	18.4	10.8	0.5	14.0	3.7	30.3	45.2	4.1	0.9	11.7		
LSD 0.10	1.3	10.5	1.0	1.5	1.6	0.6	0.48	1.3	0.6	6.5		
LSD 0.05	1.6	12.7	1.2	1.8	2.0	0.7	0.58	1.6	0.7	7.8		

Planting Date = September 21, 2017; Harvest Date = July 23; Previous Crop = Oats

¹ Vigor: 1 = worst and 10 = best

² Early height taken May 17 and plant height taken at harvest.

³ Ergot = ergot sclerotia bodies in a processed sample expressed as percent by weight.