

North Central Research Extension Center—Minot
Emmer Variety Trial

Variety	Days						Grain Yield			2 Year	3 Year
	to Head	Plant Height	Lodge	Kernel Weight	Test Weight	Protein %	2007	2008	2009		
Bowman	DAP ¹	in	0-9 ²	g/1000	lb/bu	%				bu/A	
Bowman	63	40	6.3	62.7	36.2	14.0	--	75.8	80.4	75.8	--
Common-H	63	40	5.3	63.5	36.3	13.9	--	88.1	103.3	88.1	--
Common-M	64	41	6.0	66.4	36.0	14.1	--	79.2	99.7	79.2	--
Common-R	63	40	7.3	63.1	37.2	14.2	--	75.4	98.4	75.4	--
Lucille	64	42	5.3	75.8	36.0	13.8	26.1	85.9	115.2	56.0	75.7
ND Common	63	41	6.3	73.1	37.3	14.0	37.2	86.5	121.0	61.9	81.6
Red Vernal	64	42	4.3	78.3	35.1	13.9	--	88.7	114.5	88.7	--
LSD 5%	NS	NS	1.8	4.8	1.5	NS	NS	7.2	13.0	--	--
C.V.%	1.5	4.1	20.3	4.7	2.7	2.7	26.8	5.8	8.4	--	--
Mean	63	41	5.8	69.0	36.3	14.0	31.7	82.8	104.6	--	--

Planted on May 6 with a seeding rate of 1 million PLS and harvested on August 21.

2007 Evaluation had severe heat stress during the heading period.

¹*DAP = Days after planting.*

²*Lodging score based on scale 0-9 (0 = upright, 9 = flat).*