

**North Central Research Extension Center—Minot
Rye Variety Trial on No-till Recrop (winter)**

Variety	Days to Head	Lodge	Plant Height	Kernel Weight	Test Weight	Protein	Grain Yield			2 Year	3 Year
							2007	2008	2009		
	DAJ ¹	0-9 ²	in	g/1000	lb/bu	%	-----bu/A-----				
Aroostok	20	6.0	51	26.2	55.3	14.8	--	--	67.9	--	--
Dacold	18	1.5	45	31.9	55.3	9.4	--	--	105.3	--	--
Hancock	19	2.0	50	35.7	56.0	12.3	--	--	82.0	--	--
Rymin	21	3.3	49	33.8	55.5	12.0	--	--	83.7	--	--
Spooner	22	3.5	51	36.7	55.8	11.9	--	--	78.9	--	--
Wheeler	19	1.5	52	42.5	52.8	14.9	--	--	42.6	--	--
LSD 5%	NS	1.0	3	2.9	0.5	0.9	--	--	16.5	--	--
C.V.%	14.9	21.0	4.1	5.6	0.6	5.2	--	--	13.6	--	--
Mean	20	3.1	50	34.5	55.3	12.1	--	--	81.6	--	--

Planted on September 26 with a seeding rate of 1.2 million PLS and harvested on August 18.

This trial was initiated in 2009.

¹ DAJ = Days after June 1.

² Lodging score based on scale 0-9 (0 = upright, 9 = flat).