

SMALL GRAIN PRE-HARVEST

Herbicide	Product/A (ai/A)	Weeds	When to Apply	Remarks and Paragraphs																									
Glyphosate ⁹ For HRS, Durum and Winter Wheat and Feed Barley Only.	Up to 0.75 lb ae See Remarks.	Annual and perennial grass and broadleaf weeds including Canada thistle.	Wheat and barley: Hard-dough stage, 30% or less grain moisture. Allow a 7 day PHI.	<table border="0"> <tr> <td>lb ae/gal</td> <td>lb ai/gal</td> <td>0.38 ae</td> <td>0.57 ae</td> <td>0.75 ae</td> </tr> <tr> <td>3</td> <td>4</td> <td>= 16 fl oz</td> <td>24 fl oz</td> <td>32 fl oz</td> </tr> <tr> <td>4/4.17</td> <td>5.4/5.1</td> <td>= 12 fl oz</td> <td>18 fl oz</td> <td>24 fl oz</td> </tr> <tr> <td>4.5</td> <td>5.5</td> <td>= 11 fl oz</td> <td>16 fl oz</td> <td>22 fl oz</td> </tr> <tr> <td>5</td> <td>6.1</td> <td>= 10 fl oz</td> <td>15 fl oz</td> <td>20 fl oz</td> </tr> </table> <p>Do not apply more than 0.75 lb ae/season. Do not apply on wheat or barley grown for seed because reduced germination/vigor may occur. Apply 0.75 lb ae/A or more for Canada thistle control. May be applied with 2,4-D or dicamba for improved broadleaf weed control. Add AMS fertilizer at 8.5 lb/100 gal. Refer to label for adjuvant use and application information. A3-7 B2 B8 D8 T3-10</p>	lb ae/gal	lb ai/gal	0.38 ae	0.57 ae	0.75 ae	3	4	= 16 fl oz	24 fl oz	32 fl oz	4/4.17	5.4/5.1	= 12 fl oz	18 fl oz	24 fl oz	4.5	5.5	= 11 fl oz	16 fl oz	22 fl oz	5	6.1	= 10 fl oz	15 fl oz	20 fl oz
lb ae/gal	lb ai/gal	0.38 ae	0.57 ae	0.75 ae																									
3	4	= 16 fl oz	24 fl oz	32 fl oz																									
4/4.17	5.4/5.1	= 12 fl oz	18 fl oz	24 fl oz																									
4.5	5.5	= 11 fl oz	16 fl oz	22 fl oz																									
5	6.1	= 10 fl oz	15 fl oz	20 fl oz																									
2,4-D ⁴ ester For HRS, Durum, and Winter Wheat, Barley, and Rye	1.5 to 3 pt 4EC/SL (0.75 to 1.5 lb)	Broadleaf weeds.	Wheat and oat: Dough stage to harvest. Allow a 14 day PHI.	Do not feed straw to livestock. Use only 2,4-D brands labeled for preharvest application. Drift to broadleaf crops is especially hazardous at this time. B3 T5-7 T11																									
Dicamba ⁴ + 2,4-D ⁴ For HRS, Durum, and Winter Wheat Only	0.5 to 1 pt + 1 to 2 pt 4EC/SL (0.25 to 0.5 lb + 0.5 to 1 lb)		Wheat: Hard-dough stage and green color is gone from the nodes (joints) of the stem. Allow a 7 day PHI.	Do not feed treated straw to livestock. Drift to broadleaf crops is especially hazardous at this time. A3 A7 B6 C4 T1-3 T5-7 Y9																									
Sharpen (saflufenacil ¹⁴) For HRS, Durum, and Winter Wheat Barley and Triticale Only	1 to 2 fl oz (0.36 to 0.72 oz)	Annual broadleaf weeds.	Wheat: Hard-dough stage and grain with less than 30% moisture. Allow a 3 day PHI.	Do not apply Sharpen to cereals grown for seed because reduced germination/vigor may occur. Apply with MSO adjuvant at 1.5 pt/A + AMS at 8.5 to 17 lbs/100 gal or 28% N at 1.25 to 2.5 gal/100 gal. Apply with glyphosate for additional weed control weed and desiccation. Sharpen has no grass activity. Refer to label for crop rotation intervals. Caution: MRL's have not been established for cereal grain exportation to international markets. B12																									